

Pacific Journal of Mathematics

SHARPENED POLYNOMIAL APPROXIMATION

ALAN GREENWELL LAW AND ANN L. MCKERRACHER

SHARPENED POLYNOMIAL APPROXIMATION

ALAN G. LAW AND ANN L. MCKERRACHER

In 1951, W. Wolibner showed that a real continuous function on a closed interval can be uniformly approximated by a polynomial which interpolates at prescribed points and which has a uniform norm agreeing with the function's. This fit can be sharpened to include matching of some relative extrema as well. The paper characterizes functions that permit Simultaneous Approximation and Interpolation which is Norm-Preserving and Extrema-Matching over the entire interval except, perhaps, for a subset of arbitrarily-small diameter.

1. Introduction. The classical Weierstrass Theorem of (uniform) polynomial approximation to real, continuous functions has been generalized in many ways [2, 3]; one type of extension concerns approximations in various normed linear spaces [3]. However, if the element to be approximated remains in the space of continuous functions on a closed and finite interval, and if the uniform approximation is by polynomial, then the generalizations involve "closeness" of approximation. As an example of this second type, the Walsh result of 1935 [2] shows existence of a uniformly-good polynomial approximation to a function that also interpolates the function at prescribed points. Deutsch and Morris [3] have pointed out that W. Wolibner [4] extended this last result in 1951; he, in essence, proved existence of a polynomial which (uniformly) approximates a continuous function, and which interpolates the function, and whose (uniform) norm is equal to that of the function on the prescribed interval. Theorem 1 shows that such a fit can easily be sharpened to include matching of relative extrema too; that is, finitely-many relative extrema of the function are relative extrema of the approximating polynomial (where a minimum corresponds to a minimum and a maximum to a maximum, of course).

If f has an infinite number of relative extrema, no polynomial can match at all of them—otherwise, the derivative of the polynomial would have an infinite number of zeros. However, some continuous functions have approximating polynomials that match all extreme points except those over a closed set of arbitrarily-small diameter; Theorem 2 shows that a function has such approximating polynomials when, and only when, the points where the function achieves its extrema constitute a (countable) convergent sequence.

Throughout the discussion, the space considered is the set $C[a, b]$ of continuous, real-valued functions endowed with the max norm,

$\| \cdot \|$, on the closed and finite interval $[a, b]$.

2. **Polynomial approximations which match extrema.** Wolibner's theorem actually shows that a finite number of points in the plane, for which the ordinates of any two adjacent points are distinct, can be interpolated by a polynomial that preserves monotonicity of the piecewise-linear function, ϕ , connecting the points. It is clear in the proof of this theorem that the polynomial constructed not only is monotonically increasing or decreasing where ϕ is, but is a uniform approximation to ϕ as well. Now, any member of $C[a, b]$ can be approximated arbitrarily closely by the piecewise linear function which connects certain points on the graph (a sufficient number with equally-spaced abscissae, for instance). If to such a set of points there are adjoined, if necessary, (i) a point $(x_0, f(x_0))$ where $|f(x_0)| = \|f\|$, and (ii) intermediate points so that any two adjacent points in the final set will have distinct ordinates, then Wolibner's development for the corresponding polygonal function gives:

THEOREM 1. *Suppose f is a member of $C[a, b]$ and that $t_i (1 \leq i \leq m)$ are m given values in $[a, b]$. Let $s_j (1 \leq j \leq k)$ be k points at which f has relative extrema. Then, for any positive ε , there exists a polynomial p such that:*

- (i) $\|f - p\| < \varepsilon$,
- (ii) $p(t_i) = f(t_i)$ for $1 \leq i \leq m$,
- (iii) $\|p\| = \|f\|$, and
- (iv) $p(s_j) = f(s_j)$ and p has a relative extremum at each s_j for $1 \leq j \leq k$ (a maximum of p corresponds to a maximum of f , and a minimum to a minimum).

Thus, any f in $C[a, b]$ has a polynomial that simultaneously approximates, interpolates, preserves norm and, further, matches extrema at a finite number of points.

DEFINITION. Suppose f is a member of $C[a, b]$, and let X be a given subset of $[a, b]$. f is said to have property S on X if for any $\varepsilon > 0$ and for any prescribed finite subset, T , of $[a, b]$ there exists a polynomial p such that:

- (i) $\|f - p\| < \varepsilon$,
- (ii) $p(t) = f(t)$ for $t \in T$,
- (iii) $\|p\| = \|f\|$, and
- (iv) at each point of X where f has a relative extremum, p interpolates f and p has a relative extremum (of the same type) there.

Wolibner's result is that any f has property S on the null set;

Theorem 1 can easily be used to show that f has property S on $[a, b]$ if, and only if, the set of points in $[a, b]$ at which f has relative extrema is finite. The next theorem characterizes those functions in $C[a, b]$ which have property S on sets $X = [a, b] - V$, where V is a closed set with arbitrarily-small diameter (that is, $d(V) = \sup_{x, y \in V} |x - y|$ is arbitrarily small).

THEOREM 2. *For any member, f , of $C[a, b]$, the following two assertions are equivalent:*

- (a) *corresponding to any given $\varepsilon > 0$, there exists a closed subset V , with $d(V) < \varepsilon$, for which f has property S on $[a, b] - V$;*
- (b) *the points of $[a, b]$ where f has relative extrema constitute a (countable) convergent sequence.*

Proof. Suppose, first, that the relative extrema of f occur only at $\{x_1, x_2, x_3, \dots\}$ and let x_0 , in $[a, b]$, denote the limit of this sequence. For $\varepsilon > 0$ the set $V = [x_0 - \varepsilon/3, x_0 + \varepsilon/3] \cap [a, b]$ has diameter less than ε and, by Theorem 1, f has property S on $[a, b] - V$.

For the converse, let E denote the set of points in $[a, b]$ at which f has relative extrema. If E is finite, there is nothing to prove; so, suppose that E is infinite. Let $\{V_n\}$, $n \geq 1$, be a sequence of closed subsets of $[a, b]$ such that f has property S on $[a, b] - V_n$, and such that $d(V_n) < 1/n$ for $n \geq 1$. Since E is infinite, $\{V_n\}$ has the finite intersection property; thus, $V = \bigcap_{n=1}^{\infty} V_n$ is nonempty. But $V \subset V_n$ implies that $d(V) = 0$ - consequently, $V = \{x_0\}$ for some x_0 in $[a, b]$. The relation

$$\begin{aligned} E &= \{E \cap V\} \cup \left\{ E \cap \left([a, b] - \bigcap_{n=1}^{\infty} V_n \right) \right\} \\ &= \{E \cap V\} \cup \bigcup_{n=1}^{\infty} \{E \cap ([a, b] - V_n)\} \end{aligned}$$

shows that E is a countable union of finite sets and, hence, is countable. Finally, since each V_n contains both x_0 and infinitely-many elements of E , and because $d(V_n) \rightarrow 0$, x_0 must be the limit of the sequence E .

It is, perhaps, not surprising that Theorem 2 no longer holds if diameter is replaced by (Lebesgue) measure μ . Of course, (b) would still imply (a) since $\mu(V) = d(V)$; however, the following example shows that the converse would fail. For each $n = 1, 2, 3, \dots$, let (a_n, b_n) be an open, middle-third interval removed from $[0, 1]$ in the construction of the Cantor ternary set K . Let $g(x)$ be defined as follows: $g(x) = 0$ for $x \in K$, $g(x) = x - a_n$ if $a_n < x < (a_n + b_n)/2$ and $g(x) = b_n - x$ if $b_n > x \geq (a_n + b_n)/2$, for $n \geq 1$. It is easy to check that g is in $C[0, 1]$, that it attains its relative minima on K and that it has a relative maximum at $(a_n + b_n)/2$ for each $n \geq 1$. Thus, the

corresponding set at which g has relative extrema is uncountable. On the other hand, for any $N \geq 1$, g has only a finite number of extrema on $\bigcup_{n=1}^N (a_n, b_n)$; hence, with $V_N = [0, 1] - \bigcup_{n=1}^N (a_n, b_n)$, g has property S on $X = [0, 1] - V_N$ and $\mu(V_N)$ can be made arbitrarily small.

REFERENCES

1. K. A. Bush, *Pathological functions, crinkly and wild*, J. Math. Anal. Appl., **35** (1971), 559-562.
2. P. J. Davis, *Interpolation and Approximation*, Blaisdell, New York, 1963.
3. F. Deutsch and P. D. Morris, *On simultaneous approximation and interpolation which preserves norm*, J. Approximation Theory, **2** (1969), 355-373.
4. W. Wolibner, *Sur un polynôme d'interpolation*, Colloq. Math., **2** (1951), 136-137.

Received February 1, 1973. Work by the first author was supported in part by National Research Council of Canada Grant A-7186.

UNIVERSITY OF SASKATCHEWAN, REGINA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific of Journal Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 51, No. 2

December, 1974

Robert F. V. Anderson, <i>Laplace transform methods in multivariate spectral theory</i>	339
William George Bade, <i>Two properties of the Sorgenfrey plane</i>	349
John Robert Baxter and Rafael Van Severen Chacon, <i>Functionals on continuous functions</i>	355
Phillip Wayne Bean, <i>Helly and Radon-type theorems in interval convexity spaces</i>	363
James Robert Boone, <i>On k-quotient mappings</i>	369
Ronald P. Brown, <i>Extended prime spots and quadratic forms</i>	379
William Hugh Cornish, <i>Crawley's completion of a conditionally upper continuous lattice</i>	397
Robert S. Cunningham, <i>On finite left localizations</i>	407
Robert Jay Daverman, <i>Approximating polyhedra in codimension one spheres embedded in S^n by tame polyhedra</i>	417
Burton I. Fein, <i>Minimal splitting fields for group representations</i>	427
Peter Fletcher and Robert Allen McCoy, <i>Conditions under which a connected representable space is locally connected</i>	433
Jonathan Samuel Golan, <i>Topologies on the torsion-theoretic spectrum of a noncommutative ring</i>	439
Manfred Gordon and Edward Martin Wilkinson, <i>Determinants of Petrie matrices</i>	451
Alfred Peter Hallstrom, <i>A counterexample to a conjecture on an integral condition for determining peak points (counterexample concerning peak points)</i>	455
E. R. Heal and Michael Windham, <i>Finitely generated F-algebras with applications to Stein manifolds</i>	459
Denton Elwood Hewgill, <i>On the eigenvalues of a second order elliptic operator in an unbounded domain</i>	467
Charles Royal Johnson, <i>The Hadamard product of A and A^*</i>	477
Darrell Conley Kent and Gary Douglas Richardson, <i>Regular completions of Cauchy spaces</i>	483
Alan Greenwell Law and Ann L. McKerracher, <i>Sharpened polynomial approximation</i>	491
Bruce Stephen Lund, <i>Subalgebras of finite codimension in the algebra of analytic functions on a Riemann surface</i>	495
Robert Wilmer Miller, <i>TTF classes and quasi-generators</i>	499
Roberta Mura and Akbar H. Rhemtulla, <i>Solvable groups in which every maximal partial order is isolated</i>	509
Isaac Namioka, <i>Separate continuity and joint continuity</i>	515
Edgar Andrews Rutter, <i>A characterization of QF - 3 rings</i>	533
Alan Saleski, <i>Entropy of self-homeomorphisms of statistical pseudo-metric spaces</i>	537
Ryōtarō Satō, <i>An Abel-maximal ergodic theorem for semi-groups</i>	543
H. A. Seid, <i>Cyclic multiplication operators on L_p-spaces</i>	549
H. B. Skerry, <i>On matrix maps of entire sequences</i>	563
John Brendan Sullivan, <i>A proof of the finite generation of invariants of a normal subgroup</i>	571
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, VI</i>	573
Ronson Joseph Warne, <i>Generalized $\omega - \mathcal{L}$-unipotent bisimple semigroups</i>	631
Toshihiko Yamada, <i>On a splitting field of representations of a finite group</i>	649