

Pacific Journal of Mathematics

**A PROOF OF THE FINITE GENERATION OF INVARIANTS OF
A NORMAL SUBGROUP**

JOHN BRENDAN SULLIVAN

A PROOF OF THE FINITE GENERATION OF INVARIANTS OF A NORMAL SUBGROUP

JOHN BRENDAN SULLIVAN

A fundamental theorem in the development of the quotient theory of an affine algebraic group G shows that the coordinate functions invariant under a normal subgroup form a finitely generated algebra. We show that this theorem follows from the finite field generation of the quotient field of the algebra of invariant coordinate functions in the connected case.

Notation. Let K be an algebraically closed field and A an integral domain Hopf algebra over K . Denote by $[A]$ the field of fractions of A , and by $G(A)$ the group of K -algebra morphisms from A to K . There is the natural left action (translation) of $G(A)$ on A , denoted by (\cdot) .

For the sake of completeness, we include a proof of a known proposition for fields:

PROPOSITION 0. *$K \subset L \subset E$ fields. If E is finitely generated as a field over K , then L is finitely generated as a field over K .*

Proof. Let x_1, \dots, x_t be a transcendence basis for L over K . We will see that L is a finite extension of $\mathcal{L} = K(x_1, \dots, x_t)$. Let y_1, \dots, y_n be a transcendence basis for E over \mathcal{L} ; since E is finitely generated over K , E is a finite extension of $\mathcal{L}(y_1, \dots, y_n)$. Since $\mathcal{L}(y_1, \dots, y_n)$ is purely transcendental over \mathcal{L} and L is algebraic over \mathcal{L} , $\mathcal{L}(y_1, \dots, y_n)$ and L are linearly disjoint over \mathcal{L} . Therefore, the dimension of E over $\mathcal{L}(y_1, \dots, y_n)$ is at least as large as the dimension of L over \mathcal{L} . So L is finite over \mathcal{L} and L is finitely generated over K .

PROPOSITION 1. *Let $A_1 \subset A$ be domain Hopf algebras. Then $[A_1] \cap A = A_1$.*

Proof. Let $f = a/b$ be an element of $[A_1] \cap A$, where $a, b \in A_1$. For the purpose of demonstrating this proposition, we may suppose that A is generated as a Hopf algebra by a, b , and f and that A_1 is generated by a and b . Let M be the K -linear span of $G(A) \cdot f$; M is a finite-dimensional $G(A)$ -invariant subspace of $[A_1] \cap A$. Let $I \subset A_1$ be the ideal $\{c \in A_1 \mid cM \subset A_1\}$; since M is finite-dimensional, $I \neq (0)$. If $I = A_1$, then $f \in A_1$, as was to be shown. Otherwise, from $g \cdot M = M$ and $g \cdot A_1 = A_1$ for $g \in G(A)$, it follows that $g \cdot I = I$; so, I is a

$G(A)$ -module. Moreover, since $I \subset A_1$ and $G(A)$ separates the points of A_1 , we have $G(A_1) \cdot I = I$. Since $I \neq A_1$, by the Nullstellensatz there is an element x of $G(A_1)$ which vanishes on I . Therefore, $0 = x(I) = x(G(A_1) \cdot I) = (x \cdot G(A_1))(I) = G(A_1)(I)$. By the Nullstellensatz, $I = (0)$. This contradicts $I \neq (0)$.

PROPOSITION 2. *If $A_1 \subset A$ are domain Hopf algebras and $[A_1] = [A]$, then $A_1 = A$.*

Proof. $A_1 = [A_1] \cap A = A$.

THEOREM. *For $A \subset B$ domain Hopf algebras, if B is a finitely generated K -algebra, then A is a finitely generated K -algebra.*

Proof. Since B is finitely generated as a K -algebra, $[B]$ is finitely generated as a field over K . By Proposition 0, $[A]$ is finitely generated as a field over K . Let a_1, \dots, a_n be field generators for $[A]$ where $a_i \in A$. Let A_1 be the sub-Hopf algebra of A generated by a_1, \dots, a_n ; so, $[A_1] = [A]$. By Proposition 2, $A_1 = A$. Therefore, A is finitely generated as a K -algebra.

For $G(B)$ a connected affine algebraic group with domain Hopf algebra B of coordinate functions and H a normal subgroup, the subalgebra B^H of B of H -invariant elements is a sub-Hopf algebra; since B is finitely generated, so is B^H by the theorem.

From this point, the extension to the nonconnected case is given in [1, p. 38].

REFERENCE

1. G. Hochschild, *Introduction to Affine Algebraic Groups*, Holden-Day, San Francisco, 1971.

Received February 2, 1973.

UNIVERSITY OF CALIFORNIA, BERKELEY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI*
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

* C. R. DePrima California Institute of Technology, Pasadena, CA 91109, will replace J. Dugundji until August 1974.

Pacific Journal of Mathematics

Vol. 51, No. 2

December, 1974

Robert F. V. Anderson, <i>Laplace transform methods in multivariate spectral theory</i>	339
William George Bade, <i>Two properties of the Sorgenfrey plane</i>	349
John Robert Baxter and Rafael Van Severen Chacon, <i>Functionals on continuous functions</i>	355
Phillip Wayne Bean, <i>Helly and Radon-type theorems in interval convexity spaces</i>	363
James Robert Boone, <i>On k-quotient mappings</i>	369
Ronald P. Brown, <i>Extended prime spots and quadratic forms</i>	379
William Hugh Cornish, <i>Crawley's completion of a conditionally upper continuous lattice</i>	397
Robert S. Cunningham, <i>On finite left localizations</i>	407
Robert Jay Daverman, <i>Approximating polyhedra in codimension one spheres embedded in S^n by tame polyhedra</i>	417
Burton I. Fein, <i>Minimal splitting fields for group representations</i>	427
Peter Fletcher and Robert Allen McCoy, <i>Conditions under which a connected representable space is locally connected</i>	433
Jonathan Samuel Golan, <i>Topologies on the torsion-theoretic spectrum of a noncommutative ring</i>	439
Manfred Gordon and Edward Martin Wilkinson, <i>Determinants of Petrie matrices</i>	451
Alfred Peter Hallstrom, <i>A counterexample to a conjecture on an integral condition for determining peak points (counterexample concerning peak points)</i>	455
E. R. Heal and Michael Windham, <i>Finitely generated F-algebras with applications to Stein manifolds</i>	459
Denton Elwood Hewgill, <i>On the eigenvalues of a second order elliptic operator in an unbounded domain</i>	467
Charles Royal Johnson, <i>The Hadamard product of A and A^*</i>	477
Darrell Conley Kent and Gary Douglas Richardson, <i>Regular completions of Cauchy spaces</i>	483
Alan Greenwell Law and Ann L. McKerracher, <i>Sharpened polynomial approximation</i>	491
Bruce Stephen Lund, <i>Subalgebras of finite codimension in the algebra of analytic functions on a Riemann surface</i>	495
Robert Wilmer Miller, <i>TTF classes and quasi-generators</i>	499
Roberta Mura and Akbar H. Rhemtulla, <i>Solvable groups in which every maximal partial order is isolated</i>	509
Isaac Namioka, <i>Separate continuity and joint continuity</i>	515
Edgar Andrews Rutter, <i>A characterization of QF - 3 rings</i>	533
Alan Saleski, <i>Entropy of self-homeomorphisms of statistical pseudo-metric spaces</i>	537
Ryōtarō Satō, <i>An Abel-maximal ergodic theorem for semi-groups</i>	543
H. A. Seid, <i>Cyclic multiplication operators on L_p-spaces</i>	549
H. B. Skerry, <i>On matrix maps of entire sequences</i>	563
John Brendan Sullivan, <i>A proof of the finite generation of invariants of a normal subgroup</i>	571
John Griggs Thompson, <i>Nonsolvable finite groups all of whose local subgroups are solvable, VI</i>	573
Ronson Joseph Warne, <i>Generalized $\omega - \mathcal{L}$-unipotent bisimple semigroups</i>	631
Toshihiko Yamada, <i>On a splitting field of representations of a finite group</i>	649