

# Pacific Journal of Mathematics

## **GLOBAL REFLECTION FOR A CLASS OF SIMPLE CLOSED CURVES**

JAMES MCLEAN SLOSS

# GLOBAL REFLECTION FOR A CLASS OF SIMPLE CLOSED CURVES

JAMES M. SLOSS

Global reflection is considered for a class of closed Jordan curves  $\Gamma: [x(\theta), y(\theta)]$ ,  $0 \leq \theta < 2\pi$  where  $x(\theta)$  and  $y(\theta)$  are trigonometric polynomials. Every curve of this form is algebraic and global reflection across it reduces to investigating an algebraic function and its critical points. The reflection function is picked to be that solution of the algebraic equation that maps  $\Gamma: [x(\theta), y(\theta)]$  pointwise into  $[x(\theta), -y(\theta)]$ . This function is defined and analytic except on a finite set of points inside  $\Gamma$ , and at each of these points it is continuous.

1. Introduction. Reflection across an analytic arc which is a generalization of inversion in a circle and reflection across a straight line, goes back to Schwarz. Because of the current interest, see e.g. [1], [3], [4], [5], in reflection of solutions of plane elliptic differential equations across analytic arcs, it seems appropriate to analyze the global reflection across a fairly general class of closed Jordan curves.

We shall investigate the class of Jordan rectifiable curves  $\Gamma$  of the form

$$(1.1) \quad x(\theta) = \sum_{k=0}^n a_k \cos k\theta + b_k \sin k\theta$$

$$0 \leq \theta < 2\pi, n \geq m$$

$$(1.2) \quad y(\theta) = \sum_{k=0}^m \alpha_k \cos k\theta + \beta_k \sin k\theta$$

with  $x'^2(\theta) + y'^2(\theta) \neq 0$ ,  $(a_n, b_n) \neq (0, 0) \neq (\alpha_m, \beta_m)$  and if  $m = n$  then either  $\alpha_n^2 + \beta_n^2 \neq a_n^2 + b_n^2$  or  $\alpha_n a_n + \beta_n b_n \neq 0$ .

The investigation will be reduced to analyzing a certain algebraic equation  $M[z, \zeta] = 0$  arising from (1.1) and (1.2) (see (2.5)–(2.8)).

Let  $R$  be a simply-connected region bounded by a curve  $\Gamma$  of the form (1.1) and (1.2). Let  $S$  be the finite set of points made up of zeros of the resultant polynomials

$$P(z) = R[M[z, \zeta], M_{\bar{\zeta}}[z, \zeta]] = 0$$

and

$$Q(z) = R[M[z, \zeta], M_z[z, \zeta]] = 0.$$

Let  $L_i$  be a rectifiable Jordan arc in  $R$  containing  $\{e_1, \dots, e_s\} = R \cap S$ . Then for one of the functions

$$\zeta = G(z)$$

defined by  $M[z, \zeta] = 0$  we have

- 1  $G(z)$  is defined and analytic on  $R - \{e_1, \dots, e_s\}$ ,
- 2  $G(z)$  is single-valued on  $R \setminus L_i \cup \Gamma$ ,
- 3  $G'(z) \neq 0$  on  $R \setminus L_i \cup \Gamma$ .
- 4 For  $z$  on  $\Gamma$ ,  $\bar{z} = G(z)$ .
- 5  $\overline{G[R \setminus L_i]} \cap R \setminus L_i = \emptyset$ .
- 6 About each  $e_j$  ( $1 \leq j \leq s$ ) we have either

(i)  $G(z)$  is defined, single-valued and analytic on a neighborhood of  $e_j$  with

$$G(z) = (z - e_j)^2 G^*(z),$$

$G^*(z)$  analytic and thus  $G'(e_j) = 0$  or

(ii) on some neighborhood of  $e_j$

$$G(z) = \sum_{k=0}^{\infty} f_k [\sqrt[p]{z - e_j}]^k, \quad 1 \leq p \leq 2n \quad (n \text{ of (1.1)}),$$

$f_k$  constant, or

(iii)  $G(z)$  is defined, single-valued and analytic on a neighborhood of  $e_j$  and  $G'(e_j) \neq 0$ .

In the event  $M[z, \zeta]$  is irreducible 6(iii) is excluded. We shall denote for  $z$  in  $R \setminus L_i$

$$\hat{z} = \overline{G(z)}.$$

$G(z)$  is the reflection function and  $\hat{z}$  is the reflection of  $z$  across  $\Gamma$ .

7  $G(z)$  can be extended to be defined and analytic and single-valued on

$$\{R \setminus L_i\} \cup \Gamma \cup \{\overline{G(R \setminus L_i)}\} = \{R \setminus L_i\} \cup \Gamma \cup \{\widehat{R \setminus L_i}\}$$

with

$$\hat{\hat{z}} = z \quad \text{for } z \text{ in } \{R \setminus L_i\} \cup \Gamma \cup \{\widehat{R \setminus L_i}\}.$$

It is the proof of 6(ii) that gives the most difficulty.

**2. Geometrical reflection.** To begin our investigation of reflection across a rectifiable Jordan curve  $\Gamma$  of the form

$$(2.1) \quad x(\theta) = \sum_{k=0}^n a_k \cos k\theta + b_k \sin k\theta \quad 0 \leq \theta < 2\pi, n \geq m$$

$$(2.2) \quad y(\theta) = \sum_{k=0}^m \alpha_k \cos k\theta + \beta_k \sin k\theta$$

we let

$$t = e^{i\theta} = \cos \theta + i \sin \theta$$

and express (2.1) and (2.2) in terms of  $t$  and  $\bar{t}$ , then (2.1) and (2.2) become:

$$(2.3) \quad 2x = 2\alpha_0 + \bar{c}_1 t + c_1 \bar{t} + \bar{c}_2 t^2 + c_2 \bar{t}^2 + \cdots + \bar{c}_n t^n + c_n \bar{t}^n$$

and

$$(2.4) \quad 2y = 2\alpha_0 + \bar{\gamma}_1 t + \gamma_1 \bar{t} + \bar{\gamma}_2 t^2 + \gamma_2 \bar{t}^2 + \cdots + \bar{\gamma}_m t^m + \gamma_m \bar{t}^m$$

with

$$c_k = a_k + i b_k, \quad \gamma_k = \alpha_k + i \beta_k.$$

If we multiply (2.3) by  $\bar{t}^n$  and (2.4) by  $\bar{t}^m$  we see that (2.3) and (2.4) are equivalent respectively to:

$$f(t) \equiv \bar{c}_n + \bar{c}_{n-1} \bar{t} + \cdots + \bar{c}_1 \bar{t}^{n-1} + 2(a_0 - x) \bar{t}^n + c_1 \bar{t}^{n+1} + \cdots + c_n \bar{t}^{2n} = 0$$

and

$$g(t) \equiv \bar{\gamma}_m + \bar{\gamma}_{m-1} \bar{t} + \cdots + \bar{\gamma}_1 \bar{t}^{m-1} + 2(\alpha_0 - y) \bar{t}^m + \gamma_1 \bar{t}^{m+1} + \cdots + \gamma_m \bar{t}^{2m} = 0.$$

Thus the curve  $\Gamma$  is given by exactly those  $t$  for which  $f(t) = 0$  and  $g(t) = 0$ , i.e., by the common roots of  $f(t)$  and  $g(t)$ . But a necessary and sufficient condition for  $f(t)$  and  $g(t)$  to have common roots is that Sylvester's determinant  $D(f, g)$  of order  $(2n + 2m) \times (2n + 2m)$  vanish. If we let

$$\alpha(x) = 2(a_0 - x), \quad \beta(y) = 2(\alpha_0 - y)$$

then

$$(2.5) \quad D(f, g) = \det \begin{bmatrix} \bar{c}_n & \bar{c}_{n-1} & \cdots & \bar{c}_1 & \alpha & c_1 & \cdots & c_{n-1} & c_n & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \bar{c}_n & \bar{c}_{n-1} & \cdots & \bar{c}_1 & \alpha & c_1 & \cdots & c_{n-1} & c_n & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & \bar{c}_n & \bar{c}_{n-1} & \cdots & \bar{c}_1 & \alpha & c_1 & \cdots & c_{n-1} & c_n & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \cdots & \bar{c}_n & \bar{c}_{n-1} & \cdots & \bar{c}_1 & \alpha & c_1 & \cdots & c_{n-1} & c_n & 0 & 0 \\ \bar{\gamma}_m & \bar{\gamma}_{m-1} & \cdots & \bar{\gamma}_1 & \beta & \gamma_1 & \cdots & \gamma_{m-1} & \gamma_m & 0 & 0 & \cdots & 0 & 0 \\ \bar{\gamma}_m & \bar{\gamma}_{m-1} & \cdots & \bar{\gamma}_1 & \beta & \gamma_1 & \cdots & \gamma_{m-1} & \gamma_m & 0 & \cdots & 0 & 0 \\ \bar{\gamma}_m & \bar{\gamma}_{m-1} & \cdots & \bar{\gamma}_1 & \beta & \gamma_1 & \cdots & \gamma_{m-1} & \gamma_m & \cdots & 0 & 0 \\ 0 & 0 & 0 & \cdots & \bar{\gamma}_m & \bar{\gamma}_{m-1} & \cdots & \bar{\gamma}_1 & \beta & \gamma_1 & \cdots & \gamma_{m-1} & \gamma_m & 0 & 0 \end{bmatrix}$$

provided  $c_n \neq 0 \neq \gamma_m$ . Since  $f$  and  $g$  are fixed, we define

$$\Delta[\alpha(x), \beta(y)] = D(f, g).$$

Then  $\Gamma$  is given by the algebraic equation:

$$\Delta[\alpha(x), \beta(y)] = 0, \quad \alpha = 2(a_0 - x), \quad \beta = 2(\alpha_0 - y).$$

We now consider the algebraic equation of order  $2n$  in  $\zeta$ :

$$M[z, \zeta] = A\left[\alpha\left(\frac{z+\zeta}{2}\right), \beta\left(\frac{z-\zeta}{2i}\right)\right] \\ = g_{2n}(z)\zeta^{2n} + g_{2n-1}(z)\zeta^{2n-1} + \cdots + g_2(z)\zeta^2 + g_1(z)\zeta + g_0(z) = 0$$

and investigate the Riemann surface that this equation defines over the  $z = x + iy$  plane.

First we prove:

LEMMA 2.1.

(i) If  $n > m$  and  $c_n \neq 0 \neq \gamma_m$ , then

$$g_{2n}(z) = \pm \frac{1}{(2n)!} |c_n|^{2m} = \text{constant} \neq 0.$$

(ii) If  $n = m$  and  $c_n \neq 0 \neq \gamma_n$  and  $c_n + i\gamma_n \neq 0 \neq \bar{c}_n + i\bar{\gamma}_n$ , then

$$g_{2n}(z) = \pm \frac{1}{(2n)!} (c_n + i\gamma_n)^n (\bar{c}_n + i\bar{\gamma}_n)^n = \text{constant} \neq 0.$$

*Proof.* First we note that

$$g_{2n}(z) = \frac{1}{(2n)!} \frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta]_{\zeta=0}.$$

In order to differentiate  $M[z, \zeta]$ , it will be convenient to introduce the notation:

$$C = C^{m \times m} = \begin{bmatrix} c_n & c_{n-1} & c_{n-2} & \cdots & c_{n-m} \\ 0 & c_n & c_{n-1} & \cdots & c_{n-m+1} \\ 0 & 0 & c_n & \cdots & c_{n-m+2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & c_n \end{bmatrix}, \quad I = I^{m \times m} = \begin{bmatrix} \gamma_m & \gamma_{m-1} & \gamma_{m-2} & \cdots & \gamma_1 \\ 0 & \gamma_m & \gamma_{m-1} & \cdots & \gamma_2 \\ 0 & 0 & \gamma_m & \cdots & \gamma_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \gamma_m \end{bmatrix}.$$

Then

$$M[z, \zeta] = \begin{array}{c|c|c|c|c|c} & m & n-m & 2m & n-m & m \\ \hline m & \bar{C} & A_1 & \alpha & A_2 & 0 \\ & & & \alpha & & \\ m & 0 & B_1 & & B_2 & C^\top \\ & & & & & \alpha \\ \hline m & \bar{I} & \beta & & & \\ & & \beta & & & \\ 2n-m & 0 & & & & 0 \\ & & & & & 2n-m \\ & & & & & \beta \\ & & & & & I^\top \\ \hline m & & & 2n & & m \end{array} \quad \begin{array}{l} \alpha = 2a_0 - (z + \zeta) \\ \beta = 2\alpha_0 + i(z - \zeta) \end{array}$$

where, as indicated,  $A_1, B_1, A_2, B_2$  are matrices of size  $n - m \times m$ . Thus

$$\frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta] =$$

$\bar{C}$	0	$-I$	0	0	
0	0		0	$C^\top$	
$\bar{F}$	$-iI$			0	
0				$F^\top$	

  

$$\frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta] =$$

$\bar{C}$	0	$I$	0	0	
0	0		0	$C^\top$	
$\bar{F}$	$iI$			0	
0				$F^\top$	

Next we perform the following set of operations

(i)<sub>1</sub> multiply the  $m + 1$  column by  $i\bar{\gamma}_m$  and add it to the first column

(ii)<sub>1</sub> multiply the  $m + 1$  column by  $i\bar{\gamma}_{m-1}$  and add it to the second column

(iii)<sub>1</sub> multiply the  $m + 1$  column by  $i\bar{\gamma}_{m-2}$  and add it to the third column etc. to  $m$

(i)<sub>2</sub> multiply the  $m + 2$  column by  $i\bar{\gamma}_m$  and add it to the second column

(ii)<sub>2</sub> multiply the  $m + 2$  column by  $i\bar{\gamma}_{m-1}$  and add it to the third column etc. to  $m - 1$

⋮

(i)<sub>m</sub> multiply the  $2m$  column by  $i\bar{\gamma}_m$  and add it to the  $m$ th column. This yields the following result:

$$\frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta] =$$

$\tilde{C}$	0	$I$	0	0	
0	0		0	$C^\top$	
0	$iI$			0	
0				$F^\top$	

where if  $n - m \leq m$ , i.e.,  $n \leq 2m$  then

$$\tilde{C} = \bar{C} + i \begin{pmatrix} \begin{array}{c|c} n-m & \\ \hline & \end{array} & \begin{array}{cccc} \bar{\gamma}_m & \bar{\gamma}_{m-1} & \bar{\gamma}_{m-2} & \cdots \\ 0 & \bar{\gamma}_m & \bar{\gamma}_{m-1} & \cdots \\ 0 & 0 & \bar{\gamma}_m & \cdots \\ 0 & 0 & 0 & \cdots 0 \\ 0 & 0 & 0 & \cdots 0 \end{array} \end{pmatrix} \quad m \times m$$

and if  $n - m > m$

$$\tilde{C} = \bar{C}.$$

Next we perform the following set of operations

(i)<sub>1</sub> multiply column  $2n + m$  by  $i\gamma_m$  and add it to column  $2n + 2m$

(ii)<sub>1</sub> multiply column  $2n + m$  by  $i\gamma_{m-1}$  and add it to column  $2n + 2m - 1$

(iii)<sub>1</sub> multiply column  $2n + m$  by  $i\gamma_{m-2}$  and add it to column  $2n + 2m - 2$  etc. to  $m$

(i)<sub>2</sub> multiply column  $2n + 2m - 1$  by  $i\gamma_m$  and add it to column  $2n + 2m - 1$

(ii)<sub>2</sub> multiply column  $2n + m - 1$  by  $i\gamma_{m-1}$  and add it to column  $2n + 2m - 2$  etc. to  $m - 1$

⋮

(i)<sub>m</sub> multiply column  $2n + 1$  by  $i\gamma_m$  and add it to column  $2n + m + 1$ . This yields the following result:

$$\frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta] = \begin{array}{|c|c|c|c|c|} \hline \bar{C} & 0 & & 0 & 0 \\ \hline 0 & 0 & I & 0 & C^* \\ \hline 0 & & & 0 & \\ \hline 0 & & iI & & 0 \\ \hline & & & & 0 \\ \hline \end{array}$$

where if  $n - m \leq m$ , i.e.,  $n \leq 2m$  then

$$C^* = C^T + i \begin{pmatrix} \begin{array}{cccccc} 0 & 0 & 0 & \cdot & \cdot & \cdot & 0 \\ 0 & 0 & 0 & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \gamma_m & 0 & 0 \\ \cdot & \cdot & \cdot & \gamma_{m-1} & \gamma_m & 0 \\ \cdot & \cdot & \cdot & \gamma_{m-1} & \gamma_{m-1} & \gamma_m \end{array} & \begin{array}{c|c} n-m & \\ \hline & \\ \hline 0 & \end{array} \end{pmatrix} \quad m \times m$$

and if  $n - m > m$

$$C^* = C^\top .$$

The next set of operations is as follows:

- (i) multiply row  $n + m + 1$  by  $i$  and add to row 1
- (ii) multiply row  $n + m + 2$  by  $i$  and add to row 2
- (iii) multiply row  $n + m + 3$  by  $i$  and add to row 3
- (2*m*) multiply row  $n + 2m$  by  $i$  and add to row  $2m$ .

This yields:

$$\frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta] = \begin{array}{|c|c|c|c|c|} \hline \tilde{C} & 0 & & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & C^* \\ \hline 0 & & & & 0 \\ \hline 0 & & iI & & 0 \\ \hline & & & & 0 \\ \hline \end{array} .$$

Since  $c_n \neq 0$  and if  $n = m$ ,  $\bar{c}_n + i\gamma_n \neq 0 \neq c_n + i\gamma_n$  the above determinant, with appropriate column operations, is also given by:

$$\frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta] = (i)^{2n} \begin{array}{|c|c|c|c|c|} \hline \tilde{D} & 0 & & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & D^* \\ \hline 0 & & & & 0 \\ \hline 0 & & I & & 0 \\ \hline & & & & 0 \\ \hline \end{array}$$

where

$$\begin{aligned} \tilde{D} &= \begin{array}{ll} \bar{c}_n I^{m \times m} & \text{if } n > m \\ (\bar{c}_n + i\bar{\gamma}_n) I^{n \times n} & \text{if } n = m \\ c_n I^{m \times m} & \text{if } n > m \end{array} \\ D^* &= \begin{array}{ll} (c_n + i\gamma_n) I^{n \times n} & \text{if } n = m . \end{array} \end{aligned}$$

Thus

$$\begin{aligned} \frac{\partial^{2n}}{\partial \zeta^{2n}} M[z, \zeta] &= \pm (\bar{c}_n)^m (c_n)^m \quad \text{if } n > m \\ &= \pm (\bar{c}_n + i\bar{\gamma}_n)^n (c_n + i\gamma_n)^n \quad \text{if } n = m \end{aligned}$$

which proves the lemma.



If we write

$$\begin{aligned} M[z, \zeta] &= 4 \left[ \alpha \left( \frac{z + \zeta}{2} \right), \beta \left( \frac{z - \zeta}{2i} \right) \right] \\ &= h_{2n}(\zeta) z^{2n} + h_{2n-1}(\zeta) z^{2n-1} + \dots + h_1(\zeta) z + h_0(\zeta) = 0, \end{aligned}$$

then we also have:

LEMMA 2.2.

(i) If  $n > m$  and  $c_n \neq 0 \neq \gamma_m$  then

$$h_{2n}(\zeta) = \pm \frac{1}{(2n)!} |c_n|^{2m} = \text{constant} \neq 0.$$

(ii) If  $n = m$  and  $c_n \neq 0 \neq \gamma_n$  and  $c_n - i\gamma_n \neq 0 \neq \bar{c}_n - i\bar{\gamma}_n$  then

$$h_{2n}(\zeta) = \pm \frac{1}{(2n)!} (c_n - i\gamma_n)^n (\bar{c}_n - i\bar{\gamma}_n)^n.$$

*Proof.* As in Lemma 2.1

$$h_{2n}(\zeta) = \frac{1}{(2n)!} \frac{\partial^{2n}}{\partial z^{2n}} M[z, \zeta]_{z=0}$$

and the proof proceeds as in Lemma 2.1.

We now recall some well-known facts from the theory of algebraic functions and Riemann surfaces, see e.g., [2].

We restrict ourselves to the case where  $M[z, \zeta]$  is irreducible.

A point  $z_0$  is called a critical point of  $M[z, \zeta]$  if either

(i)  $g_{2n}(z_0) = 0$  or

(ii)  $M[z_0, \zeta] = 0$  has multiple roots.

THEOREM (i). If  $z_0$  is a point such that  $g_{2n}(z_0) \neq 0$  and  $\zeta_0$  is a root of  $M[z_0, \zeta]$  of multiplicity  $l$ ,  $1 \leq l \leq 2n$ , then there exists an  $\varepsilon > 0$  and  $\delta(\varepsilon) > 0$  such that if  $z_1 \neq z_0$  lies in the disc  $D(z_0, \delta)$  of radius  $\delta$  about  $z_0$  then  $M(z_1, w) = 0$  has exactly  $l$  distinct roots in  $D(\zeta_0, \varepsilon)$ . [1], p. 122.

If  $z_0$  is a point such that  $g_{2n}(z_0) \neq 0$  and  $M[z_0, \zeta]$  has no multiple roots,  $M[z_0, \zeta_0] = 0$ , then from the above theorem  $l = 1$  and for every  $z = z_1$  in  $D(z_0, \delta)$ , there is exactly one root of  $M[z_1, \zeta] = 0$ . Thus on  $D(z_0, \delta)$   $M[z, \zeta] = 0$  defines a single-valued continuous function  $\zeta = g_1(z)$  for which  $\zeta_0 = g_1(z_0)$ .

THEOREM (ii).  $g_1(z)$  is an analytic function of  $z$  on  $D(z_0, \delta)$ .

In the case  $g_{2n}(z) = \text{constant} \neq 0$  then there are at most a finite number of critical points.

Let  $e_1, e_2, \dots, e_r$  be the critical points and join these and  $\infty$  by  $L_0$ , any nonintersecting smooth arc and half line. Cut the plane along  $L_0$ . Then on the cut plane,  $M[z, \zeta] = 0$  defines  $2n$  single-valued analytic functions  $G_1(z), \dots, G_{2n}(z)$ .

**THEOREM (iii).** *About each critical point  $e$  we have the following expansion in a neighborhood of  $e$*

$$G(z) = \sum_{m=-\infty}^{\infty} f_m(\sqrt[p]{z-e})^m$$

where

- (1)  $1 \leq p \leq 2n$
- (2) *there are at most a finite number of negative powers*
- (3) *if  $g_{2n}(e) \neq 0$  there are no negative powers.*

We shall also need

**LEMMA 2.3.** *Let*

- (1)  $x'^2(\theta) + y'^2(\theta) \neq 0$
- (2)  $(a_n, b_n) \neq 0 \neq (\alpha_n, \beta_n)$
- (3) *either  $\alpha_n^2 + \beta_n^2 \neq b_n^2 + a_n^2$  or  $\alpha_n a_n + b_n \beta_n \neq 0$*

*then for no point  $z$  of  $\Gamma$  (i) is  $\zeta = \bar{z}$  a multiple root of  $M[z, \zeta]$ ; (ii) is  $g_{2n}(z) = 0$ .*

*Proof.* Since

$$\begin{aligned}(\gamma_n - ic_n) &= (\alpha_n + i\beta_n) - i(a_n + ib_n) = \alpha_n + b_n + i(\beta_n - a_n) \\(\bar{\gamma}_n - i\bar{c}_n) &= (\alpha_n - i\beta_n) - i(a_n - ib_n) = \alpha_n - b_n - i(\beta_n + a_n)\end{aligned}$$

then

$$(\gamma_n - ic_n)(\bar{\gamma}_n - i\bar{c}_n) = \alpha_n^2 + \beta_n^2 - (a_n^2 + b_n^2) - 2i[\alpha_n a_n + b_n \beta_n] \neq 0 \text{ by (2).}$$

Thus we see by Lemma 2.1 that  $g_{2n}(z_0)$  is never zero. For  $z$  on  $\Gamma$ , i.e., for  $z = x(\theta) + iy(\theta)$ ,  $0 \leq \theta < 2\pi$ , we have

$$M[x(\theta) + iy(\theta), x(\theta) - iy(\theta)] \equiv 0, \quad 0 \leq \theta < 2\pi.$$

Thus if we differentiate with respect to  $\theta$ , we see since (1) holds, that

$$\frac{(x' + iy')^2}{x'^2 + y'^2} = \frac{M_\zeta[z, \bar{z}]}{M_z[z, \bar{z}]} \neq 0 \quad \text{on } \Gamma.$$

Thus  $M_\zeta[z_0, \zeta] \neq 0$  for  $\zeta$  on  $\Gamma$ , i.e., for  $\zeta = \bar{z}_0$  and the proof is complete.

**LEMMA 2.4.** *Let*

- (1)  $(a_n, b_n) \neq 0 \neq (\alpha_n, \beta_n)$  and

(2) either  $\alpha_n^2 + \beta_n^2 \neq a_n^2 + b_n^2$  or  $\alpha_n a_n + \beta_n b_n \neq 0$   
 then there are at most a finite number of  $z$  for which

$$M[z, \zeta] = 0 \quad \text{and} \quad M_z[z, \zeta] = 0.$$

*Proof.*  $z$  is a point at which the result of the lemma holds if and only if for  $\zeta$  fixed

$$M_1[z] = M[z, \zeta]$$

has a multiple root. Thus

$$\begin{aligned} M_1[z] &= h_{2n}(\zeta)z^{2n} + h_{2n-1}(\zeta)z^{2n-1} + \cdots + h_0(\zeta) \\ M'_1[z] &= 2nh_{2n}(\zeta)z^{2n-1} + \cdots + h_1(\zeta) \end{aligned}$$

and from Lemma 2.2 and the assumption of the lemma

$$h_{2n}(\zeta) = \text{constant} \neq 0.$$

But a necessary and sufficient condition for  $M_1[z]$  to have a multiple root is that the resultant

$$R[M_1, M'_1] = 0.$$

As this is a polynomial in  $\zeta$ , the conclusion of the lemma follows.

LEMMA 2.5. *Let the hypotheses of Lemma 2.3 hold, then for no point  $z$  of  $\Gamma$  do we have for  $\zeta = \bar{z}$*

$$M[z, \zeta] = 0 \quad \text{and} \quad M_z[z, \zeta] = 0.$$

*Proof.* From the proof of Lemma 2.3

$$\frac{M_z[z, \bar{z}]}{M_\zeta[z, \bar{z}]} = \frac{(x' - iy')^2}{x'^2 + y'^2} \neq 0$$

which is the conclusion.

We shall assume that for  $\Gamma$  we have

$$(1) \quad x'^2(\theta) + y'^2(\theta) \neq 0$$

$$(2) \quad (a_n, b_n) \neq 0 \neq (\alpha_n, \beta_n)$$

$$(3) \quad \text{either } \alpha_n^2 + \beta_n^2 \neq (b_n^2 + a_n^2) \text{ or } \alpha_n a_n + b_n \beta_n \neq 0.$$

Assume, moreover, that  $M[z, \zeta]$  is irreducible. Let  $e_1, e_2, \dots, e_r$  be the set of critical points of  $M[z, \zeta] = 0$  and let  $e_{r+1}, e_{r+2}, \dots, e_{s_0}$  (by Lemma 2.4) be the set of  $z$  for which

$$M[e_j, \zeta] = 0 \quad \text{and} \quad M_z[e_j, \zeta] = 0$$

and let

$$S = \{e_j: 1 \leq j \leq s_0\}.$$

Let  $z_0$  be a point of  $\Gamma$  and  $M[z_0, \bar{z}_0] = 0$ . By Lemma 2.3,  $\zeta = \bar{z}_0$  is not a multiple root of  $M[z_0, \zeta]$  and thus  $M[z_0, \zeta] = 0$  defines a single-valued function of  $z$ ,  $\zeta = G(z)$  in some neighborhood  $N_0$  of  $z_0$  with  $G(z_0) = \bar{z}_0$ . Moreover, for each point  $z$  on  $\Gamma \cap N_0$  we have

$$\bar{z} = G(z)$$

by Lemma 2.3. Analytically continuing  $G(z)$  around  $\Gamma$  we return to  $G(z)$ . Since if we arrive at  $G_1(z)$  where  $G(z)$  and  $G_1(z)$  are defined on a common neighborhood of  $z_0$ , then for  $z$  on  $\Gamma$ ,  $z(\theta) = x(\theta) + iy(\theta)$  and  $\zeta(\theta) = \overline{z(\theta)}$  is a periodic function and thus

$$G(z) = G_1(z) \quad \text{on } \Gamma.$$

Therefore, they agree on the common neighborhood. From this it follows that  $G(z)$  is single-valued and analytic on a neighborhood of  $\Gamma$ .

Let  $S_i = \{e_i, \dots, e_s\}$  be that subset of  $S$  (renumber if necessary) that is contained in the interior of  $\Gamma$  and let  $S_e = S \setminus S_i$ , then  $G(z)$  is analytic on  $R \setminus S_i$ . Moreover, if we join each  $e_j$  of  $S_i$  by a Jordan arc  $L_i$  then  $G(z)$  is analytic and single-valued on

$$R_0 = R \setminus L_i \cup \text{neighborhood of } \Gamma.$$

Since on  $R_0$  we have  $G(z)$  is single-valued and analytic then

$$G'(z) = - \frac{M_z[z, \zeta]}{M_\zeta[z, \zeta]} \Big|_{\zeta=G(z)}$$

which is  $\neq 0$  for  $z$  on  $\Gamma$  and  $\zeta = \bar{z}$  on  $\Gamma$  by Lemma 2.5, and also  $\neq 0$  for  $z$  on  $R \setminus L_i$  since all of the points  $M_z[z, \zeta] \Big|_{\zeta=G(z)} = 0$  lie on  $L_i$  by construction. Thus we have partially proved the

**THEOREM.**  $M[z, \zeta] = 0$  defines a function

$$\zeta = G(z)$$

which is determined by having  $z_0$  on  $\Gamma$  correspond to  $\bar{z}_0 = G(z_0)$ . For this  $G(z)$  we have

(1)  $G(z)$  is defined and analytic on  $R - \{e_1, \dots, e_s\} \cup \text{neighborhood of } \Gamma$ .

(2)  $G(z)$  is single-valued on  $R \setminus L_i \cup \text{neighborhood of } \Gamma$ .

(3)  $G'(z) \neq 0$  on  $R \setminus L_i \cup \text{neighborhood of } \Gamma$ .

(4) For  $z$  on  $\Gamma$ ,  $\bar{z} = G(z)$ .

(5)  $\overline{G[R \setminus L_i]} \cap R \setminus L_i = \emptyset$ .

(6) About each  $e_j$  ( $1 \leq j \leq s$ ) we have either

(i)  $G(z)$  is defined, single-valued and analytic on a neighborhood of  $e_j$  with

$$G(z) = (z - e_j)^2 G^*(z)$$

$G^*(z)$  analytic and thus  $G'(e_j) = 0$  or

(ii) on some neighborhood of  $e_j$

$$G(z) = \sum_{k=0}^{\infty} f_k [\sqrt[p]{z - e_j}]^k, \quad 1 \leq p \leq 2n,$$

$f_k$  constant, or

(7) If we let  $\hat{z} = G(z)$  then  $G(z)$  can be extended to be defined analytic and single-valued on

$$(i) \quad \{R \setminus L_i\} \cup \Gamma \cup \{\widehat{R \setminus L_i}\}$$

with

$$(ii) \quad \hat{\hat{z}} = z \text{ there.}$$

*Proof.* (1)-(4) have been proved.

(5) Since  $\overline{G(\Gamma)} = \Gamma$  and since  $G$  is continuous on  $R \setminus L_i$  we know either  $\overline{G[R \setminus L_i]} \subset R$  or  $\overline{G[R \setminus L_i]} \cap R \setminus L_i = \emptyset$ . We shall have proved the result if we can show that for one point  $z \in R \setminus L_i$ ,  $\overline{G(z)} \notin R$ .

Since  $x(\theta)$  and  $y(\theta)$  are analytic functions for real  $\theta$  with  $x'(\theta) + y''(\theta) \neq 0$  they can be continued as analytic functions  $x(\tau)$  and  $y(\tau)$  of the complex variable  $\tau = \theta + i\eta$  on some circle  $|\tau| < \rho$  for which, on  $|\tau| < \rho$ ,  $x'(\tau) + iy'(\tau) \neq 0$ . Then

$$g(\tau) = x(\tau) + iy(\tau) \quad |\tau| < \rho$$

maps  $\tau = \theta + i\eta$ ,  $\eta = 0$  onto a subarc  $\Gamma_0$  of  $\Gamma$  and thus maps  $|\tau| < \rho$  1-1 onto a neighborhood of  $\Gamma_0$ . Consider

$$H(\tau) = \overline{G[g(\tau)]}$$

for  $\tau$  such that  $g(\tau) \in \text{domain of } G$ . Since  $G(z)$  is defined and  $G'(z) \neq 0$  on a neighborhood of  $\Gamma$  then  $H(\tau)$  is defined on a neighborhood  $N$  of  $|\tau| < \rho$ ,  $\eta = 0$  with  $\eta = 0$  mapping onto  $\Gamma_0$  and  $H(\tau)$  establishes a 1-1 correspondence between points of  $N$  and  $H(N)$ . Thus that portion  $N$  of  $N$  for which  $\eta < 0$  maps onto the region  $R_-$  of one side of  $\Gamma_0$  and  $N_+$ , that portion of  $N$  for which  $\eta > 0$  maps onto the other side  $R_+$  of  $\Gamma_0$ . Without loss of generality let  $R_- \cap R \setminus L_i \neq \emptyset$ ,  $R_+ \cap R \setminus L_i = \emptyset$  and let  $z_0 \in R_- \cap R \setminus L_i$  be such that if  $g(\tau_0) = z_0$  then  $\bar{\tau}_0 \in N_+$ . Then  $g(\bar{\tau}_0) \in R_+$ . Note that on that neighborhood of  $\eta = 0$  where everything is defined

$$\overline{g^{-1}[\overline{G[g(\tau)]}]},$$

is analytic with

$$\overline{g^{-1}[\overline{G[g(\tau)]}]} = \overline{g^{-1}[g(\tau)]} = \bar{\tau} = \tau$$

for  $\tau$  on  $\eta = 0$  and thus is the identity map. Hence

$$g(\bar{\tau}_0) = \overline{G[g(\tau_0)]} = \overline{G[z_0]}$$

and  $\overline{G(z_0)} \in R_+$  where as  $z_0 \in R \setminus L_i$ . This completes the proof of (5).

(6) (i) and (ii) follow immediately from Theorem (iii) and Lemma 2.1.

(7) (i) follows from (3) and (5). (ii) follows from the fact that

$$\widehat{z} = \overline{G[\overline{G(z)}]}$$

is an analytic function on  $\{R \setminus L_i\} \cup \widehat{\Gamma \cup \{R \setminus L_i\}}$  with

$$\widehat{z} = z \quad \text{on } \Gamma$$

and thus

$$\widehat{z} = z \quad \text{on } \{R \setminus L_i\} \cup \Gamma \cup \widehat{\{R \setminus L_i\}}$$

and the theorem is proved.

In the event  $M[z, \zeta]$  is not irreducible then the analysis and the theorem will hold provided we decompose  $M[z, \zeta]$  into its irreducible factors and (1) study that factor which determines  $\Gamma$  and (2) prove that for this factor we have the coefficient of the highest order term in  $\zeta$  is constant and the coefficient of the highest order term in  $z$  is constant. We shall be possibly excluding an unnecessary number of points  $e_j$  where  $G(z)$  may be analytic single-valued and  $G'(z) \neq 0$ . To see that the coefficient of the highest order term of  $\zeta$  and  $z$  are constants we let

$$M[z, \zeta] \equiv Q_1(z, \zeta)Q_2(z, \zeta) \cdots Q_r(z, \zeta)$$

where the  $Q_j(z, \zeta)$  are irreducible. Then if

$$Q_j(z, \zeta) = q_{js_j}(z)\zeta^{s_j} + q_{js_{j-1}}(z)\zeta^{s_{j-1}} + \cdots + q_{j0}(z)$$

we have  $s_1 + s_2 + \cdots + s_r = 2n$ . Moreover,

$$q_{js_j}(z) = \text{constant} \neq 0 \quad \text{for all } j = 1, 2, \dots, r,$$

since

$$q_{1s_1}(z) \cdot q_{2s_2}(z) \cdots q_{rs_r}(z) \equiv g_{2n}(z) = \text{constant}.$$

Similarly if we write

$$Q_j(z, \zeta) \equiv p_{js_j}(\zeta)z^{s_j} + p_{js_{j-1}}(\zeta)z^{s_{j-1}} + \cdots + p_{j0}(\zeta)$$

we see that

$$p_{js_j}(\zeta) = \text{constant} \neq 0 \quad \text{for all } j = 1, 2, \dots, r.$$

It would be of interest to find conditions on the  $c_k$  and  $\gamma_k$  so

that  $M[z, \zeta]$  is irreducible. This would eliminate the calculation of an unnecessary number of points.

#### REFERENCES

1. P. R. Garabedian, *Partial Differential Equations*, John Wiley & Sons, 1964, (see references).
2. K. Knopp, *Theory of Functions*, Part II. Transl. by F. Bagemihl, Dover Pub., 1949.
3. H. Lewy, *On the reflection laws of second order differential equations on two independent variables*, Bull. Amer. Math. Soc., **65** (1959), 37-58.
4. J. M. Sloss, *Reflection laws of second order systems of elliptic equations in two independent variables with constant coefficients*, Pacific J. Math., **24** (1968), 541-575.
5. ———, *Reflection laws of high order elliptic differential equations in two independent variables with constant coefficients and unequal characteristics across analytic boundary conditions*, Duke Math. J., **35** (1968), 415-434.

Received January 17, 1973.

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER



# Pacific Journal of Mathematics

Vol. 52, No. 1

January, 1974

David R. Adams, <i>On the exceptional sets for spaces of potentials</i> .....	1
Philip Bacon, <i>Axioms for the Čech cohomology of paracompacta</i> .....	7
Selwyn Ross Caradus, <i>Perturbation theory for generalized Fredholm operators</i> .....	11
Kuang-Ho Chen, <i>Phragmén-Lindelöf type theorems for a system of nonhomogeneous equations</i> .....	17
Frederick Knowles Dashiell, Jr., <i>Isomorphism problems for the Baire classes</i> .....	29
M. G. Deshpande and V. K. Deshpande, <i>Rings whose proper homomorphic images are right subdirectly irreducible</i> .....	45
Mary Rodriguez Embry, <i>Self adjoint strictly cyclic operator algebras</i> .....	53
Paul Erdős, <i>On the distribution of numbers of the form <math>\sigma(n)/n</math> and on some related questions</i> .....	59
Richard Joseph Fleming and James E. Jamison, <i>Hermitian and adjoint abelian operators on certain Banach spaces</i> .....	67
Stanley P. Gudder and L. Haskins, <i>The center of a poset</i> .....	85
Richard Howard Herman, <i>Automorphism groups of operator algebras</i> .....	91
Worthen N. Hunsacker and Somashekhar Amrith Naimpally, <i>Local compactness of families of continuous point-compact relations</i> .....	101
Donald Gordon James, <i>On the normal subgroups of integral orthogonal groups</i> ....	107
Eugene Carlyle Johnsen and Thomas Frederick Storer, <i>Combinatorial structures in loops. II. Commutative inverse property cyclic neofields of prime-power order</i> .....	115
Ka-Sing Lau, <i>Extreme operators on Choquet simplexes</i> .....	129
Philip A. Leonard and Kenneth S. Williams, <i>The septic character of 2, 3, 5 and 7</i> ...	143
Dennis McGavran and Jingyal Pak, <i>On the Nielsen number of a fiber map</i> .....	149
Stuart Edward Mills, <i>Normed Köthe spaces as intermediate spaces of <math>L_1</math> and <math>L_\infty</math></i> .....	157
Philip Olin, <i>Free products and elementary equivalence</i> .....	175
Louis Jackson Ratliff, Jr., <i>Locally quasi-unmixed Noetherian rings and ideals of the principal class</i> .....	185
Seiya Sasao, <i>Homotopy types of spherical fibre spaces over spheres</i> .....	207
Helga Schirmer, <i>Fixed point sets of polyhedra</i> .....	221
Kevin James Sharpe, <i>Compatible topologies and continuous irreducible representations</i> .....	227
Frank Siwiec, <i>On defining a space by a weak base</i> .....	233
James McLean Sloss, <i>Global reflection for a class of simple closed curves</i> .....	247
M. V. Subba Rao, <i>On two congruences for primality</i> .....	261
Raymond D. Terry, <i>Oscillatory properties of a delay differential equation of even order</i> .....	269
Joseph Dinneen Ward, <i>Chebyshev centers in spaces of continuous functions</i> .....	283
Robert Breckenridge Warfield, Jr., <i>The uniqueness of elongations of Abelian groups</i> .....	289
V. M. Warfield, <i>Existence and adjoint theorems for linear stochastic differential equations</i> .....	305