

Pacific Journal of Mathematics

**A CHARACTERIZATION OF NORMAL ANALYTIC SPACES BY
THE HOMOLOGICAL CODIMENSION OF THE STRUCTURE
SHEAF**

ANDREW GUY MARKOE

A CHARACTERIZATION OF NORMAL ANALYTIC SPACES BY THE HOMOLOGICAL CODIMENSION OF THE STRUCTURE SHEAF

ANDREW MARKOE

In 1951 K. Oka proved that if X is a hypersurface in C^n whose singular set $\Sigma(X)$ has codimension at least 2 in X , then X is a normal analytic space. This result was subsequently generalized by S. Abhyankar and (independently) W. Thimm to the case of a complete intersection.

The main result of the present work is the following **criterion for normality**: *If $\dim [\Sigma(X) \cap \{x \in X: \text{codim}_x \mathcal{O}_x \leq k + 2\}] \leq k$ for all integers $k \geq -1$, then X is a normal analytic space. This is the best possible criterion following the lines of Oka, Abhyankar, and Thimm, and is, in fact, a characterization; the converse is true. The criterion implies that whenever \mathcal{O}_X has Cohen-Macaulay stalks and $\Sigma(X)$ has codimension at least 2, then X is normal. Finally, the techniques used in proving the criterion are used to obtain a vanishing theorem for the first cohomology group of the complement of a subvariety A of suitably high codimension in a Stein manifold, with coefficients in the ideal sheaf of a normal subvariety containing A .*

It is appropriate to sketch Oka's proof. Oka looked at the exact cohomology sequence (where \mathcal{I} = ideal sheaf of X in C^n)

$$\Gamma(C^n - \Sigma(X), {}_n\mathcal{O}) \xrightarrow{\rho} \Gamma(X - \Sigma(X), \mathcal{O}_X) \xrightarrow{\partial} H^1(C^n - \Sigma(X), \mathcal{I}) \xrightarrow{\alpha} H^1(C^n - \Sigma(X), {}_n\mathcal{O}).$$
 Since X is a hypersurface, $\mathcal{I} \approx {}_n\mathcal{O}$ and Cartan's three annuli theorem implies that $H^1(C^n - \Sigma(X), \mathcal{I})$ vanishes (since $\Sigma(X)$ is of codimension at least 3 in C^n). Thus ρ is surjective and the Riemann 2nd removable singularity theorem in C^n shows that holomorphic functions on $X - \Sigma(X)$ extend holomorphically to X . Therefore, X is normal.

Since I is not generally free for an arbitrary complete intersection, Abhyankar and Thimm do not use Cartan's three annuli theorem. In this paper however, the spirit of Oka's proof is followed by reintroducing Cartan's three annuli theorem in a very general version due to G. Trautmann. This result is Theorem T in §2.

An inspection of the above exact sequence shows that the best possible result using Oka's technique occurs when α is injective. It is surprising therefore that α is injective only when (and, of course, when) $H^1(C^n - \Sigma(X), \mathcal{I}) = 0$. This is an immediate consequence of the vanishing theorem (Theorem 4) presented in the following section.

The few results needed about normal analytic spaces may be

found in R. Narasimhan [4]. In particular:

(1) *Riemann's 2nd removable singularity theorem.* If X is a normal analytic space and if A is a closed subvariety with $\text{codim } A \geq 2$, the restriction $\Gamma(X, \mathcal{O}_X) \rightarrow \Gamma(X - A, \mathcal{O}_X)$ is isomorphic.

(2) If X is normal then $\text{codim } \Sigma(X) \geq 2$.

In the sequel $\Sigma = \Sigma(X)$ will denote the singular subvariety of X and dimension or codimension statements will be taken pointwise (i.e., $\text{codim } A \geq 2$ means $\text{codim}_x A \geq 2$ for all $x \in X$).

We require the notions of *homological codimension*, codh , *mth singularity subvariety of the analytic sheaf* \mathcal{F} , $S_m(\mathcal{F})$ and *profondeur (depth)*, prof . The definition and basic properties of these concepts may be found in [6]. For the present it suffices to recall that $S_m(\mathcal{F}) = \{x \in X: \text{codh}_x \mathcal{F} \leq m\}$ and that $S_m(\mathcal{F})$ is a subvariety of X , if \mathcal{F} is coherent.

1. **Characterization of normal analytic spaces.** The following theorem is a modification of a result due to G. Trautmann [8]. It follows easily from Theorem 1.14 [6] and will not be proved here.

THEOREM T. *Let (X, \mathcal{O}_X) be an analytic space, A a closed subvariety of X , \mathcal{F} a coherent analytic sheaf and q an integer ≥ 0 . Then the following conditions are logically equivalent:*

(1) *For every Stein open set $U \subset X$ the restriction*

$$\Gamma(U, \mathcal{F}) \longrightarrow \Gamma(U - A, \mathcal{F})$$

is isomorphic and

$$H^i(U - A, \mathcal{F}) = 0$$

for $1 \leq i \leq q - 1$.

(2) $\dim [A \cap S_{k+q+i}(\mathcal{F})] \leq k$ for all integers $k \geq -1$.

(3) $\text{prof}_A \mathcal{F} \geq q + 1$.

An implication of the form (2) \Rightarrow (1) can be considered as a combination of Riemann's 2nd removable singularity theorem with a generalization of Cartan's three annuli theorem (or as a general version of Frenkel's lemma).

The following theorem, which characterizes normal analytic spaces, is the most general version of the results of Oka, Abhyankar, and Thimm, [5], [1], and [7], on the normality of subvarieties of manifolds.

THEOREM 1. *A reduced analytic space X is normal if and only if*

$$\dim [\Sigma(X) \cap S_{k+2}(\mathcal{O}_X)] \leq k$$

for all integers $k \geq -1$.

Proof. If the dimension estimate holds, then (2) of Theorem T is fulfilled for $q = 1$. Hence the 2nd Riemann removable singularity theorem holds and, as in the introduction, X is normal.

Conversely, if X is normal, the second Riemann removable singularity theorem holds, so that (1) of Theorem T is verified for $q = 1$. Hence (2) of Theorem T holds, giving the dimension estimate.

There is a characterization of normal Noetherian rings due to J. P. Serre [cf. 3, page 125]: A Noetherian ring \mathcal{O} is normal if and only if $\mathcal{P} \in \text{Spec}(\mathcal{O})$ implies that $\text{prof}(\mathcal{O}_{\mathcal{P}}) \geq 2$ when $\text{ht } \mathcal{P} \geq 2$ and that $\mathcal{O}_{\mathcal{P}}$ is regular when $\text{ht } \mathcal{P} < 2$. An application of (3) in Theorem T and Theorem 1 gives the following analytic version of Serre's characterization.

COROLLARY. *An analytic space X is normal if and only if $\text{prof}_x \mathcal{O}_x \geq 2$.*

Another immediate consequence of Theorem 1 is the classical result that the nonnormal points of an analytic space form an analytic subvariety.

THEOREM 2. *If X is a reduced analytic space then $\{x \in X: X \text{ is not normal at } x\}$ is an analytic subvariety X .*

Proof. This follows immediately from Theorem 1 and the fact that for a subvariety V of X $\{x \in X: \dim_x V \geq k\}$ is a subvariety of X , too.

A local ring \mathcal{O} is said to be a *Cohen-Macaulay ring* if $\text{codh}_m \mathcal{O} = \text{Krull dim } \mathcal{O}$ (where $m = \text{maximal ideal of } \mathcal{O}$). The next result is a stronger generalization of the Oka-Abhyankar-Thimm theorems than Theorem 1.

THEOREM 3. *If X is a reduced analytic space and $x \in X$ is such that $\mathcal{O}_{x,x}$ is a Cohen-Macaulay ring and $\text{codim}_x \Sigma(X) \geq 2$, then X is normal at x .*

Proof. If $\mathcal{O}_{x,x}$ is Cohen-Macaulay, then this is true at nearby points also, therefore, $\text{codh}_y \mathcal{O}_x = \dim_y X$ for y near x . Since $y \mapsto \text{codh}_y \mathcal{O}_x$ is lower semi-continuous and $y \mapsto \dim_y X$ is upper semi-continuous, it follows that $\text{codh}_y \mathcal{O}_x$ is constant in a neighborhood of x . Let $n = \dim_x X$.

Then $\Sigma \cap S_{k+2}(\mathcal{O}_x)$ is empty if $k < n - 2$ and is Σ if $k = n - 2$. But we observe that by hypothesis, $\dim \Sigma \leq n - 2$, so the dimension estimate of Theorem 1 holds.

COROLLARY (theorems of Oka-Abhyankar-Thimm). *If X is a complete intersection such that $\text{codim } \Sigma \geq 2$, then X is normal.*

Proof. A complete intersection is Cohen-Macaulay ([2], Proposition 3, page 200).

Oka [5] proved Theorem 3 in the case that X is a hypersurface. The key idea in the proof is to show that $H^1(\mathbb{C}^n - \Sigma(X), \mathcal{I}) = 0$, where I is the ideal sheaf of X in \mathbb{C}^n . We now establish a general vanishing theorem for normal analytic spaces embedded in Stein manifolds.

THEOREM 4. *If Z is a Stein manifold and X a proper normal analytic subvariety of Z with ideal sheaf \mathcal{I} , then for any subvariety A of X with*

$$\dim A \leq \dim X - 2$$

we have

$$H^1(Z - A, \mathcal{I}) = 0.$$

Proof. We have $\mathcal{O}_X = \mathcal{O}_Z/I$ and hence we have the following exact sequence

$$\begin{aligned} \Gamma(Z - A, \mathcal{O}_Z) &\xrightarrow{\rho} \Gamma(X - A, \mathcal{O}_X) \xrightarrow{\partial} H^1(Z - A, \mathcal{I}) \\ &\longrightarrow H^1(Z - A, \mathcal{O}_Z). \end{aligned}$$

Therefore, it suffices to prove that ρ is surjective and that $H^1(Z - A, \mathcal{O}_Z) = 0$.

To show ρ surjective, consider the commutative diagram

$$\begin{array}{ccc} \Gamma(Z - A, \mathcal{O}_Z) & \xrightarrow{\rho} & \Gamma(X - A, \mathcal{O}_X) \\ \uparrow r & & \uparrow r' \\ \Gamma(Z, \mathcal{O}_Z) & \xrightarrow{r''} & \Gamma(X, \mathcal{O}_X) \end{array}$$

where r , r' , and r'' are restrictions.

Since X and Z are normal, the 2nd Riemann removable singularity theorem implies that r and r' are isomorphic. Also Cartan's Theorem B implies that r'' is surjective. Hence ρ is surjective.

Next, $H^1(Z - A, \mathcal{O}_Z) = 0$. To see this we observe that since the stalks of \mathcal{O}_Z are regular rings, $\text{codh}_x(\mathcal{O}_Z) = \dim_x Z$ for all x . Hence $A \cap S_{k+3}(\mathcal{O}_Z)$ is empty if $k < \dim Z - 3$ and is A if $k = \dim Z - 3$. Since A is of codimension ≥ 2 in X and since X is a proper subvariety of Z , $\dim A \leq \dim Z - 3$. Therefore, the dimension estimate in (2)

of Theorem T holds for $q = 2$. Hence (1) of Theorem T implies that $H^1(Z - A, \mathcal{O}_Z) = 0$, completing the proof.

The next result gives a characterization of embedded normal analytic spaces. The proof is similar to the proof of Theorem 1, so we omit the derivation.

THEOREM 5. *If Z is a Stein manifold and X is a proper analytic subvariety with \mathcal{I} the ideal sheaf of X in Z , then X is normal if and only if*

$$\dim [\Sigma(X) \cap S_{k+3}(\mathcal{I})] \leq k$$

for all $k \geq -1$.

REFERENCES

1. S. Abhyankar, *Concepts of order and rank on a complex space and a condition for normality*, Math. Ann., **141** (1960), 171-192.
2. A. Andreotti and H. Grauert, *Théorèmes de finitude pour la cohomologie des espaces complexes*, Bull. Soc. math. France, **90** (1962), 193-259.
3. H. Matsumura, *Commutative Algebra*, W. A. Benjamin, New York, 1970.
4. R. Narasimhan, *Introduction to the Theory of Analytic Spaces*, Lect. Notes Math., Springer-Verlag, Berlin, 1966.
5. K. Oka, *Lemme Fondamental*, J. Math. Soc. Japan, **3** (1951), 204-214.
6. Y. T. Siu and G. Trautmann, *Gap Sheaves and Extension of Coherent Analytic Subsheaves*, Lect. Notes Math., Springer-Verlag, Berlin, 1971.
7. W. Thimm, *Über Moduln und Ideal von holomorphen Funktionen mehrerer Variablen*, Math. Ann., **139** (1959), 1-13.
8. G. Trautmann, *Ein Endlichkeitssatz in der analytischen Geometrie*, Invent. Math., **8**, (1969), 143-174.

Received July 3, 1973 and in revised form February 8, 1974.

UNIVERSITY OF WASHINGTON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 52, No. 2

February, 1974

Harm Bart, <i>Spectral properties of locally holomorphic vector-valued functions</i>	321
J. Adrian (John) Bondy and Robert Louis Hemminger, <i>Reconstructing infinite graphs</i>	331
Bryan Edmund Cain and Richard J. Tondra, <i>Biholomorphic approximation of planar domains</i>	341
Richard Carey and Joel David Pincus, <i>Eigenvalues of seminormal operators, examples</i>	347
Tyrone Duncan, <i>Absolute continuity for abstract Wiener spaces</i>	359
Joe Wayne Fisher and Louis Halle Rowen, <i>An embedding of semiprime P.I.-rings</i>	369
Andrew S. Geue, <i>Precompact and collectively semi-precompact sets of semi-precompact continuous linear operators</i>	377
Charles Lemuel Hagopian, <i>Locally homeomorphic λ connected plane continua</i>	403
Darald Joe Hartfiel, <i>A study of convex sets of stochastic matrices induced by probability vectors</i>	405
Yasunori Ishibashi, <i>Some remarks on high order derivations</i>	419
Donald Gordon James, <i>Orthogonal groups of dyadic unimodular quadratic forms. II</i>	425
Geoffrey Thomas Jones, <i>Projective pseudo-complemented semilattices</i>	443
Darrell Conley Kent, Kelly Denis McKennon, G. Richardson and M. Schroder, <i>Continuous convergence in $C(X)$</i>	457
J. J. Koliha, <i>Some convergence theorems in Banach algebras</i>	467
Tsang Hai Kuo, <i>Projections in the spaces of bounded linear operations</i>	475
George Berry Leeman, Jr., <i>A local estimate for typically real functions</i>	481
Andrew Guy Markoe, <i>A characterization of normal analytic spaces by the homological codimension of the structure sheaf</i>	485
Kunio Murasugi, <i>On the divisibility of knot groups</i>	491
John Phillips, <i>Perturbations of type I von Neumann algebras</i>	505
Billy E. Rhoades, <i>Commutants of some quasi-Hausdorff matrices</i>	513
David W. Roeder, <i>Category theory applied to Pontryagin duality</i>	519
Maxwell Alexander Rosenlicht, <i>The nonminimality of the differential closure</i>	529
Peter Michael Rosenthal, <i>On an inversion theorem for the general Mehler-Fock transform pair</i>	539
Alan Saleski, <i>Stopping times for Bernoulli automorphisms</i>	547
John Herman Scheuneman, <i>Fundamental groups of compact complete locally affine complex surfaces. II</i>	553
Vashishtha Narayan Singh, <i>Reproducing kernels and operators with a cyclic vector. I</i>	567
Peggy Strait, <i>On the maximum and minimum of partial sums of random variables</i>	585
J. L. Brenner, <i>Maximal ideals in the near ring of polynomials modulo 2</i>	595
Ernst Gabor Straus, <i>Remark on the preceding paper: "Ideals in near rings of polynomials over a field"</i>	601
Masamichi Takesaki, <i>Faithful states on a C^*-algebra</i>	605
R. Michael Tanner, <i>Some content maximizing properties of the regular simplex</i>	611
Andrew Bao-hwa Wang, <i>An analogue of the Paley-Wiener theorem for certain function spaces on $SL(2, \mathbb{C})$</i>	617
James Juei-Chin Yeh, <i>Inversion of conditional expectations</i>	631