

Pacific Journal of Mathematics

MAXIMAL QUOTIENT RINGS OF GROUP RINGS

EDWARD WILLIAM FORMANEK

MAXIMAL QUOTIENT RINGS OF GROUP RINGS

EDWARD FORMANEK

Let $F[G]$ be the group ring of a group G over a field F , and A the subgroup of G consisting of those elements with only finitely many conjugates. Let $Q(R)$ denote the maximal (Utumi) quotient ring of a ring R . This paper proves: (1) If H is a subnormal subgroup of G , $Q(F[H])$ is naturally embedded as a subring of $Q(F[G])$. (2) $Q(F[A])$ contains the center of $Q(F[G])$. (3) If $F[G]$ is semiprime with center C , $Q(C)$ is the center of $Q(F[G])$. These results are analogues of theorems of M. Smith and D.S. Passman for the classical (Ore) quotient ring.

1. Introduction. Rings are associative and have a unit, and modules are unitary. Group rings will always be over fields, and we follow the definitions and notation of [5] for group rings and of [3] for quotient rings. In particular, if $F[G]$ is the group ring G over F , then

$$A = A(G) = \{g \in G : g \text{ has finitely many conjugates}\};$$

$$A^+ = A^+(G) = \text{torsion subgroup of } G;$$

$$\theta: F[G] \rightarrow F[A] \text{ is the natural projection.}$$

If R is a ring, $Q = Q(R)$ is the maximal quotient ring of R .

There are many quotient rings which can be associated with a ring R . The two which have received the greatest attention are the classical (Ore) quotient ring and the maximal (Utumi) quotient ring. The classical quotient ring has a relatively straightforward description, but it is only defined for rings which satisfy the so-called Ore condition. In contrast the maximal quotient ring is less easy to describe but is defined for all rings. In both cases there are distinct notions of left and right quotient rings and we will always consider left quotient rings.

For group rings the classical quotient ring has been studied by Herstein and Small [2], Passman [5, 6], M. Smith [7], and P. F. Smith [8], and the maximal quotient ring has been studied by Burgess [1].

This paper investigates the relationship of the maximal quotient rings of group rings, subgroup rings, and the centers of group rings. The object is to obtain for the maximal quotient ring analogues of theorems of Passman and M. Smith on the classical quotient ring. Their techniques are used for the group ring arguments while the quotient ring arguments reflect the formalism of the maximal quotient ring.

If R is a subring of S , there is in general no relation between $Q(R)$ and $Q(S)$. Thus to say that $Q(R)$ is a subring of $Q(S)$ for a given R and S has little meaning unless accompanied by a precise interpretation, and this will be given in the body of the paper. Modulo this interpretation, the main results are summarized by the following theorem.

THEOREM. *Let $F[G]$ be a group ring with center C .*

- (1) *If H is a subnormal subgroup of G , $Q(F[H])$ is a subring of $Q(F[G])$.*
- (2) *$Q(F[\Delta])$ contains the center of $Q(F[G])$.*
- (3) *If $F[G]$ is semiprime, $Q(C)$ is the center of $Q(F[G])$.*

I do not know if the hypothesis that $F[G]$ be semiprime is required in (3). Passman [6] has proved the analogue of (3) for the classical quotient ring without this hypothesis.

2. Dense ideals and the maximal quotient ring. With each ring R is associated a larger ring $Q = Q(R)$, called the maximal quotient ring of R . There are several equivalent constructions of Q . We will use the original one which is based on dense ideals and is due to Utumi [9, see 3, p. 96-99].

A left ideal D of R is *dense* if for each $a \in R$ the right annihilator of Da^{-1} is zero, where $Da^{-1} = \{r \in R: ra \in D\}$. (Note that if a is invertible, Da^{-1} has the usual meaning.) Some of the basic properties of dense left ideals are (see [3, p. 96-98]):

- (1) If D_1 is dense and $D_1 \subseteq D_2$ then D_2 is dense.
- (2) If D is dense and $a \in R$, then Da^{-1} is dense.
- (3) If D_1 and D_2 are dense, so is $D_1 \cap D_2$.
- (4) If D_1 and D_2 are dense and $f: D_2 \rightarrow R$ is a homomorphism, then $f^{-1}(D_1)$ is dense.

- (5) If R is commutative, D is dense iff it has zero annihilator.

The *maximal (left) quotient ring* of R is the set of all pairs (f, D) where D is a dense left ideal of R and $f: D \rightarrow R$ is a homomorphism of left R -modules, modulo the equivalence relation $(f_1, D_1) \sim (f_2, D_2)$ if f_1 and f_2 agree on $D_1 \cap D_2$. The sum and product of (f_1, D_1) and (f_2, D_2) are represented by the homomorphisms $f_1 + f_2: D_1 \cap D_2 \rightarrow R$, $f_1 f_2: f_2^{-1}(D_1) \rightarrow R$. Each $a \in R$ defines a homomorphism $T_a: R \rightarrow R$ by $T_a(r) = ra$ and the map $a \mapsto T_a$ identifies R with a subring of Q .

If R is a subring of S , then $Q(R)$ is not in general a subring of $Q(S)$ and in general there is little relation between $Q(R)$ and $Q(S)$. However, there is a natural attempt to define a homomorphism $Q(R) \rightarrow Q(S)$ and when it succeeds it is automatically an injection

of rings and then $Q(R)$ can be considered a subring of $Q(S)$. Namely, if $f: D \rightarrow R$ represents an element of $Q(R)$, one tries to extend f to an S -homomorphism $f_1: SD \rightarrow S$. f_1 is unique if it exists but in general it does not exist. Even if f_1 exists, SD may not be a dense ideal of R . If it happens that for every $(f, D) \in Q(R)$, SD is dense in S and the extension f_1 exists, then $(f_1, SD) \in Q(S)$ and the map $(f, D) \rightarrow (f_1, SD)$ identifies $Q(R)$ with a subring of $Q(S)$. It turns out that this procedure works at least for some subrings of group rings.

3. Dense ideals in group rings.

THEOREM 1. *Let H be a normal subgroup of G . If D is a dense ideal of $F[H]$, then $F[G]D$ is a dense left ideal of $F[G]$.*

Proof. Let $a = a_1g_1 + \cdots + a_kg_k \in F[G]$ where $a_i \in F$, $a_i \neq 0$, $g_i \in G$. We have to show that the right annihilator of $(F[G]D)a^{-1}$ in $F[G]$ is zero. But for each $a_i g_i$

$$(F[G]D)(a_i g_i)^{-1} = F[G]Dg_i^{-1} \supseteq g_i Dg_i^{-1}.$$

D is dense in $F[H]$ so each $g_i Dg_i^{-1}$ is dense in $F[H]$ since conjugation by g_i is an automorphism of $F[H]$. Thus

$$(F[G]D)a^{-1} \supseteq \bigcap (F[G]D)(a_i g_i)^{-1} \supseteq \bigcap g_i Dg_i^{-1} = J,$$

where J is dense in $F[H]$ since it is a finite intersection of dense left ideals of $F[H]$. J has zero right annihilator in $F[H]$ so it has zero right annihilator in $F[G]$ since $F[G]$ is a free left $F[H]$ -module. Hence $(F[G]D)a^{-1}$ also has zero right annihilator which shows that $F[G]D$ is dense.

REMARK. Theorem 1 is false if H is not normal in G . For example, let G be the free group generated by g and h and let H be the subgroup generated by h . Then $D = F[H](h - 1)$ is dense in $F[H]$ but $D' = F[G]D = F[G](h - 1)$ is not dense in $F[G]$. E.g. $D'(g - 1)^{-1} = 0$ since $g - 1$ and $h - 1$ do not have a common left multiple. In this case $F[H]$ is a commutative domain and $Q(F[H])$ is just its classical quotient ring, a field. But no nonunit of $F[H]$ becomes invertible in $Q(F[G])$.

Assume now that H is normal in G and let $\{g_i\}$ be a set of coset representatives of H in G . If $f: D \rightarrow F[H]$ represents an element of $Q(F[H])$, then $F[G]D = \oplus g_i D$, a direct sum of abelian groups, so defining $\bar{f}: F[G]D \rightarrow F[G]$ by $\bar{f}(\sum g_i d_i) = \sum g_i f(d_i)$ gives a well-defined map. To verify that \bar{f} is in fact $F[G]$ -linear it suffices to show that

if $a \in F[G]$, g_i is a coset representative, and $d \in D$, then $\bar{f}(ag_id) = a\bar{f}(g_id)$. Letting $ag_i = \sum g_j a_j$, where $a_j \in F[H]$ (a finite sum), we have

$$\begin{aligned}\bar{f}(ag_id) &= \bar{f}(\sum g_j a_j d) = \sum g_j \bar{f}(a_j d) \\ &= \sum g_j a_j f(d) = ag_i f(d) = a\bar{f}(g_id) .\end{aligned}$$

It is clear that $f \rightarrow \bar{f}$ defines a ring monomorphism of $F[H]$ into $F[G]$ which is natural. We summarize this below, noting that it is enough for H to be subnormal in G .

THEOREM 2. *Let H be a subnormal subgroup of G . Then $Q(F[H])$ is naturally identified with a subring of $Q(F[G])$ via the map $(f, D) \rightarrow (\bar{f}, F[G]D)$, where $f: D \rightarrow F[H]$ represents an element of $Q(F[H])$.*

From now on we will consider $Q(F[H])$ a subring of $Q(F[G])$ when Theorem 2 applies. If G is abelian (or more generally, nilpotent) this means that $Q(F[G])$ contains $Q(F[H])$ for every subgroup H of G .

4. The center of $Q(R)$. Suppose $f: D \rightarrow R$ represents a central element of $Q(R)$. Then f commutes with the image of R in $Q(R)$, namely with all the homomorphisms $T_a, a \in R$, where $T_a: R \rightarrow R$ is defined by $T_a(r) = ra$. fT_a is defined on $T_a^{-1}(D) = Da^{-1}$ and $T_a f$ is defined on D . Since f is central fT_a and $T_a f$ agree on $D \cap Da^{-1}$. Hence for any $a \in R$ and $d \in D \cap Da^{-1}$

$$f(da) = fT_a(d) = T_a f(d) = f(d)a .$$

LEMMA 3. *Suppose $f: D \rightarrow R$ represents a central element of $Q(R)$. Then f can be extended to a map $f: DR \rightarrow R$ by*

$$f(d_1 a_1 + \cdots + d_n a_n) = f(d_1) a_1 + \cdots + f(d_n) a_n$$

for $d_1, \dots, d_n \in D, a_1, \dots, a_n \in R$. Hence every central element of $Q(R)$ is represented by a map $f: D \rightarrow R$ where D is a two-sided ideal and f is a homomorphism of R -bimodules—i.e., $f(rds) = rf(d)s$ for $r, s \in R, d \in D$.

Proof. The only problem is to show that the extension of f is well-defined—we must show that if $\sum d_i a_i = 0$, then $\sum f(d_i) a_i = 0$. Suppose $\sum d_i a_i = 0$ and let E be the dense left ideal $E = \bigcap D(d_i a_i)^{-1}$. If $b \in E$, then $bd_i \in D, bd_i a_i \in D$, so

$$0 = f(\sum bd_i a_i) = \sum f(bd_i a_i) = \sum f(bd) a_i = b \sum f(d_i) a_i .$$

Hence $\sum f(d_i) a_i = 0$, since it is a right annihilator of the dense left ideal E .

Returning to group rings, the center of $F[G]$ is the set of finite sums $\sum x_g g$ which are constant on conjugacy classes and hence is a subring of $F[\Delta]$. Since Δ is normal in $F[G]$, $Q(F[\Delta])$ is a subring of $Q(F[G])$ and it is reasonable to suppose that it contains the center of $Q(F[G])$. We will show this but first we need some preliminaries on $\theta: f[G] \rightarrow F[\Delta]$.

Let $\{g_i\}$ be a set of coset representatives of Δ in G , with $g_1 = 1$. If $a = a_1 + g_2 a_2 + \cdots + g_k a_k$, where $a_i \in F[\Delta]$, then $\theta(g_i^{-1}a) = a_i$. From this the following lemma is routine (see [1, 4.5-4.6] for more general results).

LEMMA 4. *Let D be a left ideal of $F[G]$. Then*

- (1) *$\theta(D)$ is a left ideal of $F[\Delta]$.*
- (2) *$F[G]\theta(D) \supseteq D$.*
- (3) *If D is dense in $F[G]$, $\theta(D)$ is dense in $F[\Delta]$.*
- (4) *If D is a two-sided ideal, so are $\theta(D)$ and $F(G)\theta(D)$.*

The next result has had widespread use in the study of group rings.

LEMMA 5. (M. Smith [7, Lemma 2.3], [5, Lemma 1.3]). *Suppose $a, b, c, d \in F[G]$ and $agb = cgd$ for all $g \in G$. Then $a\theta(b) = c\theta(d)$.*

LEMMA 6. *Any central element of $Q(F[G])$ can be represented by a map $f: D \rightarrow F[G]$ where D is a two-sided ideal of $F[G]$, $\theta(D) \subseteq D$, and $f(\theta(D)) \subseteq F[\Delta]$.*

Proof. By Lemma 3, any central element can be represented by a bimodule homomorphism $f: D \rightarrow F[G]$, where D is a two-sided ideal of $F[G]$, so we will be done if we can extend f to a homomorphism $f_1: D_1 \rightarrow F[G]$, where $D_1 = F[G]\theta(D)$ and $f_1(\theta(D)) \subseteq F[\Delta]$.

Suppose $a \in D$, and let $a = a_1 + g_2 a_2 + \cdots + g_k a_k$, $f(a) = b_1 + g_2 b_2 + \cdots + g_k b_k$ where $a_i, b_i \in F[\Delta]$, (possibly some a_i, b_i are zero). $\theta(D)$ is the set of all such a_i , as a varies over D , so if we can define $f_1: F[G]\theta(D) \rightarrow F[G]$ by $f_1(ga_i) = gb_i$ for any $g \in G$, this f_1 will be the required extension. The only difficulty is to verify that f_1 is well-defined—it will then automatically be an $F[G]$ -module homomorphism, extend f , and map the dense ideal $\theta(D)$ of $F[\Delta]$ into $F[\Delta]$.

This amounts to showing that if $a_1 = \theta(a) = 0$, then $b_1 = \theta(f(a)) = 0$. To see this, suppose $\theta(a) = 0$ and let $d \in D$. Then for any $g \in G$

$$dgf(a) = f(dga) = f(d)ga \text{ since } f \text{ is a bimodule homomorphism.}$$

$$\therefore d\theta(f(a)) = f(d)\theta(a) = 0, \text{ by Lemma 5.}$$

$$\therefore D\theta(f(a)) = 0, \text{ so } \theta(f(a)) = 0 \text{ since } D \text{ is dense in } F[G].$$

Since Δ is normal in G , Theorem 2 says that $Q(F[\Delta])$ is (identified with) a subring of $Q(F[G])$. In the notation of Lemma 6, $f_1|_{\theta(D)}: \theta(D) \rightarrow F[\Delta]$ is identified with $f_1: F[G]\theta(D) \rightarrow F[G]$ which represents the same element of $Q(F[G])$ as $f: D \rightarrow F[G]$. Thus we have shown:

THEOREM 7. *The center of $Q(F[G])$ is a subring of $Q(F[\Delta])$.*

5. Semiprime group rings. In this section, the following data is fixed. F is a field, G is a group, $\Delta = \Delta(G)$. We assume that $\Delta^+(G)$ has no elements of order p if F has characteristic p . This is equivalent to assuming that $F[G]$ is semiprime by a theorem of Passman [5, Theorem 3.7]. It implies that $F[H]$ is semiprime whenever H is a subgroup of Δ . Let C denote the center of $F[G]$.

Passman used the following lemma in his work on the classical quotient ring of group rings. It plays a similar role with respect to the maximal quotient ring. Because we have the additional hypothesis that $F[H]$ is semiprime we get the additional conclusion (over [6]) that $F[Z]^{-1}F[H]$ is semisimple.

LEMMA 8. (Passman [6, Lemma 1]). *Let $H \subseteq \Delta$ be a finitely generated normal subgroup of G . Then*

(1) *H has a torsion-free central subgroup Z of finite index which is normal in G .*

(2) *The ring of fractions $F[Z]^{-1}F[H]$ obtained by inverting the nonzero elements of the central domain $F[Z]$ is a finite-dimensional semisimple algebra over the field $F[Z]^{-1}F[Z]$.*

LEMMA 9. *Let $I \neq 0$ be a G -invariant ideal of $F[\Delta]$. Then $I \cap C \neq 0$.*

Proof. Let $H \subseteq \Delta$ be a finitely generated normal subgroup of G with $I_1 = I \cap F[H] \neq 0$, and let Z be as in Lemma 8, $A = F[Z]^{-1}F[H]$, $J = AI_1$. G acts on A which is semisimple Artinian and J is a G -invariant ideal of A , so J is generated as an A -module by a G -invariant idempotent $e = a/b$, where $a \in I_1$, $b \in F[Z]$, $b \neq 0$. Let $b_1 = b, b_2, \dots, b_n$ be the finitely many G -conjugates of b . Then

$$e = a/b = ab_2 \cdots b_n/b_1 \cdots b_n.$$

e and $b_1 \cdots b_n$ are centralized by G , so $ab_2 \cdots b_n$ is central in $F[G]$. Thus $0 \neq ab_2 \cdots b_n \in I \cap C$, as required.

LEMMA 10. *Suppose D is a dense ideal of C and $f: D \rightarrow C$ is a C -homomorphism. Then*

(1) *$F[G]D$ is dense in $F[G]$.*

(2) f has a unique extension to an $F[G]$ -homomorphism

$$\bar{f}: F[G]D \rightarrow F[G].$$

(3) \bar{f} represents a central element of $Q(F[G])$.

Proof. (1) Since D is central $(F[G]D)a^{-1} \supseteq F[G]D$ for all $a \in F[G]$ so to show that $F[G]$ is dense in $F[G]$ it suffices to show that $A = \text{Ann}_{F[G]}(D) = 0$. But if $A \neq 0$, Lemma 9 says that $A \cap C \neq 0$, which contradicts the hypothesis that D is dense in C .

(2) If f has such an extension \bar{f} it is clearly unique and is defined by

$$\bar{f}(a_1 d_1 + \cdots + a_k d_k) = a_1 f(d_1) + \cdots + a_k f(d_k)$$

for $a_1, \dots, a_k \in F[G]$, $d_1, \dots, d_k \in D$.

The only problem is to show that \bar{f} is well-defined—we must show that if $\sum a_i d_i = 0$, then $\sum f(a_i) d_i = 0$. Suppose $\sum a_i d_i = 0$ and let $d \in D$. Since d is central

$$\begin{aligned} d(\sum a_i f(d_i)) &= \sum a_i d f(d_i) = \sum a_i f(dd_i) \\ &= \sum a_i f(d_i d) = (\sum a_i d_i) f(d) = 0. \end{aligned}$$

Hence $\sum a_i f(d_i) = 0$, since it annihilates the dense ideal $F[G]D$ on the right.

(3) First note that $\bar{f}: F[G]D \rightarrow F[G]$ is a bimodule homomorphism and suppose $g: D_1 \rightarrow F[G]$ represents any element of $Q(F[G])$. Then for any $d \in D$ and $d_1 \in D_1$,

$$\bar{f}g(dd_1) = \bar{f}(dg(d_1)) = \bar{f}(d)g(d_1) = g(\bar{f}(d)d_1) = g\bar{f}(dd_1).$$

Thus $\bar{f}g$ and $g\bar{f}$ are defined and agree on DD_1 . It is easy to see that DD_1 is a dense left ideal of $F[G]$, so $\bar{f}g$ and $g\bar{f}$ represent the same element of $Q(F[G])$ and so \bar{f} is central in $Q(F[G])$.

As in §3, $(f, D) \rightarrow (\bar{f}, F[G]D)$ is a ring homomorphism and we obtain a result analogous to Theorem 2.

THEOREM 11. *Let $F[G]$ be a semiprime group ring with center C . Then $Q(C)$ is naturally identified with a central subring of $Q(F[G])$ via the map $(f, D) \rightarrow (\bar{f}, F[G]D)$.*

Now that we can consider $Q(C)$ a subring of the center of $Q(F[G])$ the final step is to show that it is the whole center. We already know that the center of $Q(F[H])$ is contained in $Q(F[\Delta])$, by Theorem 7.

LEMMA 12. *If $d \in C$, $\text{Ann}_{F[G]}(d) = F[G]e$ for some central idempotent e of $F[G]$.*

Proof. Let $B = \text{Ann}_{F[G]}(d)$, $A = \text{Ann}_{F[G]}(B)$. A is an annihilator ideal of $F[G]$ and $d \in A$ so a theorem of M. Smith [7, Corollary 5.6] [5, Theorem 25.4] says that there is a central idempotent e of $F[G]$ such that $e \in A$ and $de = d$. Then $B = \text{Ann}_{F[G]}(d) = F[G](1 - e)$.

LEMMA 13. *Let D be a two-sided ideal of $F[\Delta]$ which is dense as a left ideal in $F[\Delta]$ and invariant under conjugation by elements of G . Then $D \cap C$ is dense in C .*

Proof. Since C is commutative, to show that $D \cap C$ is dense in C it suffices to show that $\text{Ann}_c(D \cap C) = 0$. Let $c \in C$, $c \neq 0$. Then $\text{Ann}_{F[G]}(c) = F[G]e$ for some central idempotent e of $F[G]$ by Lemma 12. Since D is dense, $D(1 - e) \neq 0$ and by Lemma 9, $D(1 - e) \cap C$ contains a nonzero d . Now $dc \neq 0$ since otherwise we would have $de = d$, $d(1 - e) = d$, whence $d = 0$.

Now consider a central element of $Q(F[G])$. By Lemma 6, it is represented by a map $f: D \rightarrow F[G]$ where D is a two-sided ideal of $F[G]$, $\theta(D) \subseteq D$, and $f(\theta(D)) \subseteq F[\Delta]$. By Theorem 7 and the remarks preceding, it lies in $Q(F[\Delta])$ where it is represented by $f|_{\theta(D)}: \theta(D) \rightarrow F[\Delta]$. $\theta(D)$ satisfies the hypothesis of Lemma 13, so $\theta(D) \cap C$ is dense in C . f maps $D \cap C$ into C since $f(g^{-1}dg) = g^{-1}f(d)g$ for all $d \in D$. Thus $f|_{(\theta(D) \cap C)}: \theta(D) \cap C \rightarrow C$ represents an element of $Q(C)$ and we have shown

THEOREM 14. *Let $F[G]$ be a semiprime group ring with center C . Then $Q(C)$ is the center of $Q(F[G])$.*

ACKNOWLEDGMENT. I am indebted to D. S. Passman for many improvements in the presentation.

REFERENCES

1. W. Burgess, *Rings of quotients of group rings*, Canad. J. Math., **21** (1969), 865-875.
2. I. N. Herstein and L. Small, *Rings of quotients of group algebras*, J. Algebra, **19** (1971), 153-155.
3. J. Lambek, *Lectures on Rings and Modules*, Blaisdell, Waltham, Massachusetts, 1966.
4. B. H. Neumann, *Groups with finite classes of conjugate elements*, Proc. London Math. Soc., (3) **1** (1951), 178-187.
5. D. S. Passman, *Infinite Group Rings*, Dekker, New York, 1971.
6. ———, *On the ring of quotients of a group ring*, Proc. Amer. Math. Soc., **33** (1972), 221-225.
7. M. Smith, *Group algebras*, J. Algebra, **18** (1971), 477-499.
8. P. F. Smith, *Localization in group rings*, Proc. London Math. Soc., (3) **22** (1971), 69-90.
9. Y. Utumi, *On quotient rings*, Osaka Math. J., **8** (1956), 1-18.

Received June 13, 1973. Research supported by NRC (Canada) grant A7171.

CARLETON UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 53, No. 1

March, 1974

Martin Bartelt, <i>Strongly unique best approximates to a function on a set, and a finite subset thereof</i>	1
S. J. Bernau, <i>Theorems of Korovkin type for L_p-spaces</i>	11
S. J. Bernau and Howard E. Lacey, <i>The range of a contractive projection on an L_p-space</i>	21
Marilyn Breen, <i>Decomposition theorems for 3-convex subsets of the plane</i>	43
Ronald Elroy Bruck, Jr., <i>A common fixed point theorem for a commuting family of nonexpansive mappings</i>	59
Aiden A. Bruen and J. C. Fisher, <i>Blocking sets and complete k-arcs</i>	73
R. Creighton Buck, <i>Approximation properties of vector valued functions</i>	85
Mary Rodriguez Embry and Marvin Rosenblum, <i>Spectra, tensor products, and linear operator equations</i>	95
Edward William Formanek, <i>Maximal quotient rings of group rings</i>	109
Barry J. Gardner, <i>Some aspects of T-nilpotence</i>	117
Juan A. Gatica and William A. Kirk, <i>A fixed point theorem for k-set-contractions defined in a cone</i>	131
Kenneth R. Goodearl, <i>Localization and splitting in hereditary noetherian prime rings</i>	137
James Victor Herod, <i>Generators for evolution systems with quasi continuous trajectories</i>	153
C. V. Hinkle, <i>The extended centralizer of an S-set</i>	163
I. Martin (Irving) Isaacs, <i>Lifting Brauer characters of p-solvable groups</i>	171
Bruce R. Johnson, <i>Generalized Lerch zeta function</i>	189
Erwin Kleinfeld, <i>A generalization of $(-1, 1)$ rings</i>	195
Horst Leptin, <i>On symmetry of some Banach algebras</i>	203
Paul Weldon Lewis, <i>Strongly bounded operators</i>	207
Arthur Larry Lieberman, <i>Spectral distribution of the sum of self-adjoint operators</i>	211
I. J. Maddox and Michael A. L. Willey, <i>Continuous operators on paranormed spaces and matrix transformations</i>	217
James Dolan Reid, <i>On rings on groups</i>	229
Richard Miles Schori and James Edward West, <i>Hyperspaces of graphs are Hilbert cubes</i>	239
William H. Specht, <i>A factorization theorem for p-constrained groups</i>	253
Robert L Thele, <i>Iterative techniques for approximation of fixed points of certain nonlinear mappings in Banach spaces</i>	259
Tim Eden Traynor, <i>An elementary proof of the lifting theorem</i>	267
Charles Irvin Vinsonhaler and William Jennings Wickless, <i>Completely decomposable groups which admit only nilpotent multiplications</i>	273
Raymond O'Neil Wells, Jr, <i>Comparison of de Rham and Dolbeault cohomology for proper surjective mappings</i>	281
David Lee Wright, <i>The non-minimality of induced central representations</i>	301
Bertram Yood, <i>Commutativity properties in Banach $*$-algebras</i>	307