

Pacific Journal of Mathematics

**COMPLETELY DECOMPOSABLE GROUPS WHICH ADMIT
ONLY NILPOTENT MULTIPLICATIONS**

CHARLES IRVIN VINSONHALER AND WILLIAM JENNINGS WICKLESS

COMPLETELY DECOMPOSABLE GROUPS WHICH ADMIT ONLY NILPOTENT MULTIPLICATIONS

C. VINSONHALER AND W. J. WICKLESS

A triangle of size n is a collection $\{A_u\}$ of $n(n+1)/2$ (not necessarily distinct) rank one torsion-free abelian groups indexed by all integer sequences of the form $u = i, i+1, \dots, i+j$ with $1 \leq i \leq i+j \leq n$, satisfying $T(A_u) + T(A_s) \leq T(A_{us})$ for all consecutive sequences u, s . Here $T(A_v)$ denotes the type of the rank one torsion-free abelian group A_v . If $A = \bigoplus_{i \in I} A_i$ is a direct sum of rank one torsion-free abelian groups A_i , let $\Delta(A) = \sup \{n \mid \exists \text{ a triangle of size } n \text{ of groups chosen, possibly with repetitions, from } \{A_i \mid i \in I\}\}$, $\Delta'(A) = \sup \{n \mid \exists \text{ a triangle of size } n \text{ of groups chosen without repetition from } \{A_i \mid i \in I\}\}$. An abelian group $(G, +)$ is radical iff whenever $(R, +, \cdot)$ is a ring with $(R, +) \cong (G, +)$ there exists a positive integer n with $R^n = (0)$.

THEOREM. *Let $A = \bigoplus_{i \in I} A_i$ be such that $\{T(A_i) \mid i \in I\}$ is an ordered set and $\Delta(A) < \infty$. Then A is radical.*

THEOREM. *Let $A = \bigoplus_{i \in I} A_i$ be such that $\Delta'(A) = \infty$. Then A is not radical.*

THEOREM. *Let $A = \bigoplus_{i \in I} A_i$, $B = \bigoplus_{j \in J} B_j$ be such that $\Delta(A) < \infty$, $\Delta(B) < \infty$. Then if $\{T(A_i) \mid i \in I\} \cup \{T(B_j) \mid j \in J\}$ is an ordered set $A \oplus B$ is radical. A bound is given for the index of nilpotency of any multiplication on $A \oplus B$.*

1. Preliminaries. Several authors ([2], [3], [4], [5]) have studied the class of abelian groups $(A, +)$ which admit only a trivial ring structure; i.e., if $(R, +, \cdot)$ is a ring with $(R, +) \cong (A, +)$, then $R^2 = (0)$. These are called nil groups.

In [6] a larger class was introduced—abelian groups which admit only nilpotent multiplications. More precisely:

DEFINITION 1.1. An abelian group $(A, +)$ is a radical group iff whenever $(R, +, \cdot)$ is a ring with $(R, +) \cong (A, +)$, we have $R^n = 0$ for some positive integer n .

In [6], using the techniques of [1], the class of finite rank torsion free radical groups was studied, and it was shown that this class is closed under finite direct sums.

In this paper we study completely decomposable radical groups. We work toward the goals of characterizing such groups and of obtaining information on finite direct sums of such groups.

Throughout, the word “group” means torsion-free abelian group.

We let $A = \bigoplus_{i \in I} A_i$, $B = \bigoplus_{j \in J} B_j$ be arbitrary completely decompo-

$\{x_1, \dots, x_n\} \subseteq A$. Then a triangle of size n can be formed by choosing (possibly with repetitions) groups from $\{A_i \mid i \in I\}$.

Proof. Let $x_1 * x_2 * \dots * x_n \neq 0$ and choose, for each i , some $0 \neq a_i \in A_i$. Write each x_j as a finite rational combination $x_j = \sum_{\text{finite}} \alpha_{j_i} a_i$. It follows immediately that $a_{i_1} * a_{i_2} * \dots * a_{i_n} \neq 0$ for some (not necessarily distinct) indices $\{i, \dots, i_n\} \subseteq I$. Reindex, setting $A_1 = A_{i_1}$, $A_2 = A_{i_2}$, \dots , $A_n = A_{i_n}$. The groups A_1, \dots, A_n form the top row of the triangle.

For $1 \leq i < i + j \leq n$ write $a_i * \dots * a_{i+j} = \sum \alpha_s a_s$, and let $A_{i \dots i+j}$ by any A_s such that $T(A_s) = T_A(a_i * \dots * a_{i+j})$. Such A_s always exists since $\{T(A_i) \mid i \in I\}$ is an ordered set. It is easy to check that the types of the rank one groups thus chosen satisfy the additive requirements of Definition 2.1.

The requirement that $\{T(A_i) \mid i \in I\}$ be an ordered set cannot be deleted in the above theorem. One can construct a completely decomposable group $A = \bigoplus_{i=1}^{11} A_i$ such that no triangle of size 4 can be formed by choosing groups from $\{A_i \mid 1 \leq i \leq 11\}$ but such that $A^4 \neq (0)$ under an appropriate multiplication. (The numbers 11 and 4 are the minimum rank and index for such an example.)

THEOREM 2.2. *Let $\{A_i \mid i \in I\}$ be a set of rank one groups such that a triangle of size n can be formed by choosing $n(n + 1)/2$ distinct groups from the set. Then there is a multiplication on $A = \bigoplus_{i \in I} A_i$ with a nonzero product of length n .*

Proof. Reindex the groups chosen from $\{A_i \mid i \in I\}$ as in Definition 2.1. For each index, u , choose $0 \neq a_u \in A_u$ such that for consecutive indices $r = i \dots i + j$, $s = i + j + 1 \dots i + j + k$, $i + j + k \leq n$, $h(a_r) + h(a_s) \leq h(a_{i \dots i+j+k}) = h(a_{rs})$. Such a choice is possible because of the type relationships in the triangle. Now define $a_r \cdot a_s = a_{rs}$ for all consecutive indices r, s . If r and s are not of the above form, define $a_r \cdot a_s = 0$. If A_i is not a group in the triangle, let $A_i A = A A_i = (0)$. These products define a unique associative ring structure on A . Moreover, in this ring $a_1 \cdot a_2 \cdot \dots \cdot a_n \neq 0$.

Note that the above method (or simple variations of it) cannot be used to obtain multiplication on $\bigoplus A_i$ from triangles formed by groups arbitrarily chosen from $\{A_i \mid i \in I\}$. If the same group is allowed in repeated positions, it seems impossible to overcome the consistency problems arising in the definitions of the products $a_r \cdot a_s$. However, we can obtain a nonassociative multiplication in this case.

The results of this section can be used to obtain information on completely decomposable radical groups and on direct sums of such

groups. First we prove theorem on the existence of subtriangles of a given triangle.

3. Subtriangles.

DEFINITION 3.1. Let $\{A_u\}$ be a triangle of size n , indexed as in Definition 2.1. For $1 \leq m \leq n$, a subtriangle of size m is an array of $m(m + 1)/2$ (not necessarily distinct) groups chosen from $\{A_u\}$ such that the resulting array, after reindexing, satisfies the requirements of Definition 2.1.


EXAMPLE. The array $A_1 \quad A_{34} \quad A_5$ is a subtriangle of any

$$\begin{array}{cc} A_{1234} & A_{345} \\ & A_{123456} \end{array}$$

triangle of size $n \geq 6$.

THEOREM 3.1. Let $A = \{A_i\}$, $B = \{B_i\}$ and let n be any positive integer. If m is any integer such that $m \geq f(n) \equiv \max \{3n - 1, n(n + 1)/2\}$, then any triangle of size m^m of groups from $A \cup B$ contains a subtriangle of size n of groups from A alone or from B alone.

Proof. Use induction on n . The result is easy for $n = 1$. Now consider a triangle of size m^m of groups from $A \cup B$, when $m \geq f(n)$, (n an integer not less than 2). As in Diagram 1, divide the first m^{m-1} rows into m disjoint subtriangles of size m^{m-1} . Since $m^{m-1} \geq$



$(m - 1)^{m-1}$ and $m - 1 \geq f(n - 1) = \max \{3(n - 1) - 1, (n - 1)n/2\}$, by the induction hypothesis, each of these disjoint triangle of size m^{m-1} contains a subtriangle of size $n - 1$ of groups from A alone or from B alone.

Case I. Both triangles of size $n - 1$ of groups from A and triangles of size $n - 1$ of groups from B appear as subtriangles of the triangles of size m^{m-1} . In this case, there occurs a subtriangle of size $n - 1$ of groups from one collection, say A , such that either to the right or to the left of this subtriangle there are n disjoint triangles of size m^{m-1} , each of which contains subtriangles of size $n - 1$ of groups from B . This is due to the fact that in any arrangement of $3n - 1$ objects which are of two types, one of the types must have n objects of the other type either to the right or to the left.

Without loss of generality, assume that the left-most triangle (1) contains a subtriangle of size $n - 1$ of groups from A , and that there are n triangles to the right which contain subtriangles of groups from B . The technique is to expand one of these subtriangles to a triangle of size n .

Let $G_{i_k, i_{k+1}, \dots, i_{k+j_k}}$ be the rightmost group in the k^{th} row of the subtriangle of groups from A contained in triangle 1, $1 \leq k \leq n - 1$ (see Diagram 1). If triangle r_k , $1 \leq k \leq n$, contains the k^{th} subtriangle to the right of triangle 1 of groups from B , consider, for all $2 \leq k \leq n$, the sequence of groups

$$(*)_k \quad G_{i_k, i_{k+1}, \dots, (r_k, -1)^{m^{m-1}+1}} G_{i_k, i_{k+1}, \dots, (r_{k-1})^{m-1}+2}, \dots, G_{i_k, i_{k+1}, \dots, r_k m^{m-1}} .$$


(Each of these sequences is represented by a diagonal in Diagram 1.)

Claim: Either $(*)_k$ contains a group from A or the subtriangle of groups from B in triangle r_k can be expanded to a triangle of size n of groups from B . Suppose $(*)_k$ contains no group from A . Let B^l be any group from B in triangle r_{k-1} . Now let B^l , $2 \leq l \leq n$, be the group in the sequence $(*)_k$ whose index ends in the same number as the index of the left-most group in the $l-1$ row of the subtriangle of groups from B in triangle r_k . By adjoining each B^l as the left-most group in the l^{th} row of the triangle, the subtriangle of groups from B in triangle r_k is expanded to a triangle of size n of groups from B . In this case we are done, so assume now that each sequence $(*)_k$, $2 \leq k \leq n$, contains a group from A -call it A^k . Let A^1 be any group from A in triangle r_1 (if none exists, we're done). Now by adjoining A^k as the right-most group in the k^{th} row of the A -subtriangle of triangle 1 we obtain an A -subtriangle of size n .

Case II. There exist only A -subtriangles of size $n - 1$ in the triangles 1, 2, \dots , m . If this is the case, then *any* subtriangle of

size m^{m-1} contained in the first m^{m-1} rows most contain only A -subtriangles of size $n - 1$. If there were a B -subtriangle of size $n - 1$, the techniques of Case I could be applied to construct an A - or B -subtriangle of size n .

Now consider any diagonal sequence of the form $(+): G_{i, \dots, i+j}, G_{i, \dots, i+j+1}, \dots, G_{i, \dots, i+j+m^{m-1}-1}$ where $2 \leq i \leq m^m - m^{m-1} - n - 1, n \leq j \leq m^m - m^{m-1} - i$. These $(+)$ sequences are illustrated in Diagram 2. If all of the groups in $(+)$ are in A then consider the A -subtriangle of size $n - 1$ contained in the triangle of size m^{m-1} whose top row is $G_{i+j}, G_{i+j+1}, \dots, G_{i+j+m^{m-1}-1}$. (Refer to Diagram 2.) Again using the techniques of Case I, this A -subtriangle can be expanded to a triangle of size n using the sequence $(+)$. Therefore, without loss of generality, assume that every sequence $(+)$ contains a group from B . This assumption can be used to build a B -subtriangle of size n .



Choose from triangle 1 any group B_1 from B . Say $B_1 = G_{i, \dots, i+j}$ where $2 \leq i \leq m^{m-1}$ and $0 \leq j \leq m^{m-1} - i$. If such a B_1 does not exist, we are done, for triangle 1 is then an A -subtriangle of size $m^{m-1} \geq n$. Similarly choose from triangle 2 a group $B_2 = G_{i', \dots, i'+j'}$ from B , where $m^{m-1} < i' \leq 2m^{m-1}, 0 \leq j' \leq 2m^{m-1} - i'$. Now choose a group B_{12} from B from the sequence $G_{i, i+1, \dots, i'+j'}, \dots, G_{i, i+1, \dots, i'+j'+m^{m-1}-1}$. (See Diagram 2.) This is a $(+)$ sequence unless $i' + j' - i < n$. In this case replace $i' + j'$ with $i + n$ in the above sequence. Now let B_3 be any group from B found in triangle 4. The group B_{23} is obtained from the $(+)$ sequence which begins where the diagonals from B_2 and B_3 intersect (see Diagram 2). The group B_{123} is obtained

from the $(+)$ sequence which begins where the diagonals from B_{12} and B_{23} intersect. The group B_i is then chosen from triangle 7 and the process continued. Since each new group requires a new $(+)$ sequence, it is not difficult to see that $n(n+1)/2$ $(+)$ sequences are needed in this process. Hence the requirement $m \geq n(n+1)/2$. Thus a B -triangle of size n can be constructed and the proof is complete.

REMARK. The number $f(n) = \max \{3n - 1, n(n+1)/2\}$ is probably much too large as is the triangle size m^m used in the proof. These in fact could be lowered slightly, but are retained in the interest of a less complicated proof.

4. Applications. In this section, the results of the previous sections are consolidated to give some information on when completely decomposable groups are radical. We begin with

DEFINITION 4.1. Let $A = \bigoplus_{i \in I} A_i$ be any completely decomposable group, and assume $T(A_i)$ is nil for all $i \in I$. Let $\Delta(A) = \sup \{n \mid \exists \text{ a triangle of size } n \text{ chosen, possibly with repetitions, from } \{A_i \mid i \in I\}\}$. Let $\Delta'(A) = \sup \{n \mid \exists \text{ a triangle of size } n \text{ chosen without repetition from } \{A_i \mid i \in I\}\}$.

THEOREM 4.1. Let $A = \bigoplus_{i \in I} A_i$ be as above. If $\{T(A_i) \mid i \in I\}$ is an ordered set and $\Delta(A)$ is finite, then A is radical.

Proof. If A is not radical, then for any $n \geq 0$, there exists a multiplication on A which yields a nonzero product of length greater than n . By Theorem 2.1, $\Delta(A) > n$. This contradicts $\Delta(A)$ finite.

A sort of converse to Theorem 4.1 is provided by

THEOREM 4.2. If $A = \bigoplus_{i \in I} A_i$ is completely decomposable with $\Delta'(A) = \infty$ then A is not radical.

Proof. If $\Delta'(A)$ is infinite, we show there exists a multiplication on A which produces nonzero products of arbitrary length. Let n be any positive integer. Since $\Delta'(A)$ is infinite, there exists a triangle T_1 of size $m > n$ consisting of distinct groups from $\{A_i\}$. Now let $k = f(m+1)$ and form a triangle of size $l \geq k^k$ consisting of distinct groups from $\{A_i\}$. By applying Theorem 3.1 to the disjoint collections of groups T_1 and $\{A_i\} \setminus T_1$, we obtain a subtriangle T_2 of size $m+1$ consisting of distinct groups from $\{A_i\} \setminus T_1$. Continuing this process we obtain a disjoint sequence of collections of groups, T_1, T_2, T_3, \dots , such that distinct groups in T_j may be used to form a triangle of

size at least $m + j - 1$. Thus by Theorem 2.1 a ring may be defined on the direct sum of the groups in T_j which contains a nonzero product of length $m + j - 1$. The direct sum of these rings is then non-nilpotent and is a group direct summand of A . Therefore A cannot be radical.

The final result concerns the direct sum of two completely decomposable radical groups.

THEOREM 4.3. *Let $A = \bigoplus_{i \in I} A_i$, $B = \bigoplus_{j \in J} B_j$ be completely decomposable radical groups such that $\Delta(A) = n$ and $\Delta(B) = m$ are finite. Then if $\{T(A_i) \mid i \in I\} \cup \{T(B_j) \mid j \in J\}$ is an ordered set, $A \oplus B$ is radical. Moreover, for any multiplication on $A \oplus B$, $(A \oplus B)^{k^k} = 0$ for $k = \max\{f(n + 1), f(m + 1)\}$.*

Proof. If $A \oplus B$ is not radical, then $\Delta(A \oplus B)$ is infinite by Theorem 4.1. But Theorem 3.1 then implies that either $\Delta(A)$ or $\Delta(B)$ is infinite, a contradiction. In fact, if $(A \oplus B)^{k^k} \neq (0)$ for some multiplication on $A \oplus B$, then $\Delta(A \oplus B) \geq k^k$. But by Theorem 3.1, this implies that $\Delta(A) \geq n + 1$ or $\Delta(B) \geq m + 1$, a contradiction.

REFERENCES

1. R. A. Beaumont and R. S. Pierce, *Torsion free rings*, Illinois J. Math., **5**, 61-98.
2. L. Fuchs, *Abelian Groups*, Pergamon Press, New York, 1960.
3. R. Ree and R. J. Wisner, *A note on torsion free nil groups*, Amer. Math. Soc. Proc., **7**, 6-8.
4. T. Szele, *Gruppen theoretische beziehungen bei gewissen ringkonstruktionen*, Math. Z., **54**, 168-180.
5. ———, *Zur theorie der zeroringe*, Math. Ann., **121**, 242-246.
6. W. J. Wickless, *Abelian groups which admit only nilpotent multiplications*, Pacific J. Math., **40**, 251-259.

Received February 5, 1973 and in revised form January 30, 1974.

UNIVERSITY OF CONNECTICUT

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 53, No. 1

March, 1974

Martin Bartelt, <i>Strongly unique best approximates to a function on a set, and a finite subset thereof</i>	1
S. J. Bernau, <i>Theorems of Korovkin type for L_p-spaces</i>	11
S. J. Bernau and Howard E. Lacey, <i>The range of a contractive projection on an L_p-space</i>	21
Marilyn Breen, <i>Decomposition theorems for 3-convex subsets of the plane</i>	43
Ronald Elroy Bruck, Jr., <i>A common fixed point theorem for a commuting family of nonexpansive mappings</i>	59
Aiden A. Bruen and J. C. Fisher, <i>Blocking sets and complete k-arcs</i>	73
R. Creighton Buck, <i>Approximation properties of vector valued functions</i>	85
Mary Rodriguez Embry and Marvin Rosenblum, <i>Spectra, tensor products, and linear operator equations</i>	95
Edward William Formanek, <i>Maximal quotient rings of group rings</i>	109
Barry J. Gardner, <i>Some aspects of T-nilpotence</i>	117
Juan A. Gatica and William A. Kirk, <i>A fixed point theorem for k-set-contractions defined in a cone</i>	131
Kenneth R. Goodearl, <i>Localization and splitting in hereditary noetherian prime rings</i>	137
James Victor Herod, <i>Generators for evolution systems with quasi continuous trajectories</i>	153
C. V. Hinkle, <i>The extended centralizer of an S-set</i>	163
I. Martin (Irving) Isaacs, <i>Lifting Brauer characters of p-solvable groups</i>	171
Bruce R. Johnson, <i>Generalized Lerch zeta function</i>	189
Erwin Kleinfeld, <i>A generalization of $(-1, 1)$ rings</i>	195
Horst Leptin, <i>On symmetry of some Banach algebras</i>	203
Paul Weldon Lewis, <i>Strongly bounded operators</i>	207
Arthur Larry Lieberman, <i>Spectral distribution of the sum of self-adjoint operators</i>	211
I. J. Maddox and Michael A. L. Willey, <i>Continuous operators on paranormed spaces and matrix transformations</i>	217
James Dolan Reid, <i>On rings on groups</i>	229
Richard Miles Schori and James Edward West, <i>Hyperspaces of graphs are Hilbert cubes</i>	239
William H. Specht, <i>A factorization theorem for p-constrained groups</i>	253
Robert L Thele, <i>Iterative techniques for approximation of fixed points of certain nonlinear mappings in Banach spaces</i>	259
Tim Eden Traynor, <i>An elementary proof of the lifting theorem</i>	267
Charles Irvin Vinsonhaler and William Jennings Wickless, <i>Completely decomposable groups which admit only nilpotent multiplications</i>	273
Raymond O'Neil Wells, Jr, <i>Comparison of de Rham and Dolbeault cohomology for proper surjective mappings</i>	281
David Lee Wright, <i>The non-minimality of induced central representations</i>	301
Bertram Yood, <i>Commutativity properties in Banach $*$-algebras</i>	307