

Pacific Journal of Mathematics

THE NORM OF A CERTAIN DERIVATION

CHARLES ALAN MCCARTHY

THE NORM OF A CERTAIN DERIVATION

CHARLES A. MCCARTHY

J. C. Stampfli has asked whether the norm of the derivation $\mathfrak{D}_T: A \rightarrow TA - AT$ as a mapping of the subalgebra \mathfrak{A} of $\mathfrak{B}(H)$ into $\mathfrak{B}(H)$ is given by $\inf\{2\|T - A'\|: A' \in \mathfrak{A}'\}$. That this need not be the case is shown through an example in 4×4 matrices.

H is a Hilbert space. $\mathfrak{B}(H)$ is the algebra of all bounded linear operators on H . \mathfrak{A} is a subalgebra of $\mathfrak{B}(H)$ and \mathfrak{A}' is the commutant of \mathfrak{A} .

In [6], J. C. Stampfli proved that the norm of \mathfrak{D}_T as a mapping of $\mathfrak{B}(H)$ into itself is precisely $2 \inf_{\lambda} \|T - \lambda\|$. Thus the question about $\|\mathfrak{D}_T\|$ as a mapping from \mathfrak{A} to $\mathfrak{B}(H)$ naturally arises. In addition, Kadison, Lance, and Ringrose [2, Theorem 3.1] show that if $T = T^*$ and \mathfrak{D}_T maps \mathfrak{A} into itself, then $\|\mathfrak{D}_T\| = \inf\{2\|T - A'\|: A' \in \mathfrak{A}'\}$. Our example will have T self-adjoint, which shows that their hypothesis $\mathfrak{D}_T(\mathfrak{A}) \subset \mathfrak{A}$ is not inessential.

For our example, we take H to be complex four-dimensional Hilbert spaces; elements of H are to be thought of as column 4-vectors, and elements of $\mathfrak{B}(H)$ as 4×4 matrices. We take \mathfrak{A} to be the subalgebra of diagonal matrices, so $\mathfrak{A}' = \mathfrak{A}$.

For T we take the Hermitian matrix

$$T = \frac{1}{12} \begin{pmatrix} 1 & & -4 & \frac{1}{\sqrt{14}}(-5 + 6i\sqrt{3}) & \frac{1}{\sqrt{14}}(-5 - 6i\sqrt{3}) \\ & -4 & & 4 & -2\sqrt{14} \\ \frac{1}{\sqrt{14}}(-5 - 6i\sqrt{3}) & -2\sqrt{14} & \frac{7}{2} & & \frac{1}{14}(-95 + 12i\sqrt{3}) \\ \frac{1}{\sqrt{14}}(-5 + 6i\sqrt{3}) & -2\sqrt{14} & \frac{1}{14}(-95 - 12i\sqrt{3}) & \frac{7}{2} & \end{pmatrix}$$

T is of the form $P - Q$ where P and Q are self-adjoint projections. The range of P is two-dimensional and is spanned by the orthogonal unit vectors

$$p^{(1)} = \frac{1}{2\sqrt{3}} \left(1, -1 + i\sqrt{3}, \frac{1}{\sqrt{14}}(4 - 5i\sqrt{3}), \frac{1}{\sqrt{14}}(-2 + i\sqrt{3}) \right),$$

$$p^{(2)} = \frac{1}{2\sqrt{3}} \left(1, -1 - i\sqrt{3}, \frac{1}{\sqrt{14}}(-2 - i\sqrt{3}), \frac{1}{\sqrt{14}}(4 + 5i\sqrt{3}) \right);$$

the range of Q is one-dimensional and is spanned by the unit vector

$$q = \frac{1}{2\sqrt{3}}\left(1, 2, \sqrt{\frac{7}{2}}, \sqrt{\frac{7}{2}}\right).$$

First we show that $\|\mathfrak{D}_T\| = \sup\{\|TX - XT\|: X \in \mathfrak{A}, \|X\| = 1\} < 2$. As the unit sphere of \mathfrak{A} is the convex hull of the unitary matrices in \mathfrak{A} , it suffices to consider $\|TX - XT\|$ only for diagonal unitary matrices X . As T has norm 1 and any X has norm 1, $\|TX - XT\| \leq 2$. Suppose then that there were an X for which $\|TX - XT\| = 2$ (since the set of X under consideration is compact, the supremum defining \mathfrak{D}_T is attained). Then there must be a unit vector $u \in H$ for which $\|(TX - XT)u\| = 2$, and since TX and XT are both of norm 1, we must have $\|TXu\| = 1 = \|XTu\|$; and since the norm of H is strictly convex, we must have $TXu = -XTu$. Further, since $\|Tu\| = 1$, we must have $u = Pu + Qu$. The next two relations are consequences of $TXu = -XTu$; start in the middle and work towards either end.

$$\begin{aligned} PXPu + PXQu &= PX(P + Q)u = PXu \\ &= P(P - Q)Xu = PTXu = -PXTu \\ &= -PXPu + PXQu, \end{aligned}$$

so $PXPu = 0$;

$$\begin{aligned} -QXPu - QXQu &= -QXu \\ &= QTXu = -QXTu \\ &= -QXPu + QXQu; \end{aligned}$$

so $QXQu = 0$.

Next we observe that XPu is in the range of Q and XQu is in the range of P ; for if one of these were not the case, we should have the strict inequality below:

$$\begin{aligned} 1 &= \|u\|^2 = \|Qu\|^2 + \|Pu\|^2 = \|XQu\|^2 + \|XPu\|^2 \\ &> \|PXQu\|^2 + \|QXPu\|^2 \\ &= \|PX(P + Q)u\|^2 + \|QX(P + Q)u\|^2 \\ &= \|PXu\|^2 + \|QXu\|^2 = \|TXu\|^2. \end{aligned}$$

But if $\|\mathfrak{D}_T\|$ is to be 2, we cannot allow $\|TXu\| < 1$.

Since $\|XPu\|^2 + \|XQu\|^2 = 1$, not both of XPu and XQu may be zero. Observe that operation on a vector by the diagonal unitary X does not change the absolute value of any component. If $XPu \neq 0$, then XPu is in the range of Q and the conclusion we draw is that there must be a nonzero vector in the range of p with moduli of components the same as that of q . If $XPu = 0$, then $XQu \neq 0$ and XQu is in the range of P ; we draw the same conclusion.

To finally reach the desired contradiction to the assumption $\|\mathfrak{D}_T\| = 2$, we need only show that no vector in the range of p has components of the same modulus as q . Indeed, if there were, such a vector must be of the form $p = e^{i\beta}(\cos \theta p^{(1)} + e^{i\Phi} \sin \theta p^{(2)})$ for some real β, θ, Φ . Equating the squares of the moduli of the first two components yields

$$\begin{aligned} 1 &= |\cos \theta + e^{i\Phi} \sin \theta|^2 = 1 + 2 \cos \theta \sin \theta \cos \Phi, \\ 4 &= |\cos \theta(-1 + i\sqrt{3}) + e^{i\Phi} \sin \theta(-1 - i\sqrt{3})|^2 \\ &= 4 + 8 \cos \theta \sin \theta \cos\left(\Phi + \frac{2\pi}{3}\right). \end{aligned}$$

Thus $\cos \theta \sin \theta = 0$ and p must be a multiple of $p^{(1)}$ or $p^{(2)}$; but neither of these has the moduli of their last two components the same as q .

Having demonstrated that $\|\mathfrak{D}_T\| < 2$, we show now that $\|T - A'\| \geq 1$ for every $A' \in \mathfrak{U}$. As $\|T\| = 1$, this is equivalent to showing that $\|T - D\| \geq 1$ for every diagonal matrix D . Suppose, then, that there were a diagonal matrix D with diagonal entries d_1, d_2, d_3, d_4 —for which $\|T - D\| < 1$. We may assume D real, since $\|T - \operatorname{Re} D\| = \|\operatorname{Re}(T - D)\| \leq \|T - D\| < 1$, where by $\operatorname{Re} A$ we mean $1/2(A + A^*)$.

Let p be any unit vector in the range of p , and q as before. Consider the inner product

$$((T - D)(\cos \theta p + e^{i\Phi} \sin \theta q), (\cos \theta p - e^{i\Phi} \sin \theta q)).$$

This is equal to

$$1 - (D(\cos \theta p + e^{i\Phi} \sin \theta q), (\cos \theta p - e^{i\Phi} \sin \theta q)),$$

but has absolute value less than 1. Hence

$$\operatorname{Re}(D(\cos \theta p + e^{i\Phi} \sin \theta q), (\cos \theta p - e^{i\Phi} \sin \theta q)) > 0,$$

for any choice of p, θ , and Φ . The choices $\theta = 0, p = p^{(1)}$, and $p = p^{(2)}$ give

$$\begin{aligned} \left(d_1 + 4d_2 + \frac{13}{2}d_3 + \frac{1}{2}d_4\right) &> 0, \\ \left(d_1 + 4d_2 + \frac{1}{2}d_3 + \frac{13}{2}d_4\right) &> 0, \end{aligned}$$

and hence

$$\left(d_1 + 4d_2 + \frac{7}{2}d_3 + \frac{7}{2}d_4\right) > 0.$$

But the choice $\theta = \pi/2$ gives

$$-\left(d_1 + 4d_2 + \frac{7}{2}d_3 + \frac{7}{2}d_4\right) > 0.$$

This incompatibility is a contradiction to $\|T - D\| < 1$.

The reader will observe the similarity with Example 5.5 of [3]. In spirit, we have the logarithmic analogue of the problem of conditioning matrices. One can ascertain conditions that $\|T - A'\| \geq \|T\|$ for all $A' \in \mathfrak{A}'$ by consideration of the norms $\|T\|_p = [\text{trace}(T^*T)^{p/2}]^{1/p}$ as $p \rightarrow \infty$, as in [4, Lemma 4.7, Theorem 4.8] or, more generally, [5, § 6]. For T self-adjoint, the relevant condition to have $\|T - A'\| \geq \|T\|$ for all diagonal A' is that both numbers $-\|T\|$ and $\|T\|$ are eigenvalues of T and that the spectral projections associated with these eigenvalues have proportional diagonals. Conditions involving suprema of norms over the group of diagonal unitaries are related to the moduli of components of certain vectors; see [4, Theorem 5.4], as well as [1, 2]. Finally, we note that for the analogous problem of conditioning matrices, examples such as we have constructed are not available in 3×3 matrices, nor with 4×4 real matrices.

REFERENCES

1. G. E. Forsythe and E. G. Straus, *On best conditioned matrices*, Proc. Amer. Math. Soc., **6** (1955), 340-345.
2. G. H. Golub and J. M. Varah, *On a characterization of the best l_2 scaling of a matrix*, To appear in SIAM J. Numer. Anal.
3. R. V. Kadison, E. C. Lance, and J. R. Ringrose, *Derivations and automorphisms of operator algebras*, II. J. Funct. Anal., **1** (1967), 204-221.
4. C. Mc Carthy and G. Strang, *Optimal conditioning of matrices*, SIAM J. Numer. Anal., **10** (1973), 370-388.
5. C. Mc Carthy, *Optimal conditioning of operators on Hilbert space*, Functional Analysis (Proc. symposium, Monterey, California, 1969). Academic Press, (1970), 107-125.
6. J. G. Stampfli, *The norm of a derivation*, Pacific J. Math., **33** (1970), 131-141.

Received May 2, 1973. Supported in part by NSF GP28933. I should like to thank the Department of Mathematics, University of Newcastle upon Tyne, for their hospitality and encouragement in this work.

UNIVERSITY OF MINNESOTA

AND

CHALMERS TEKNISKA HÖGSKOLA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific of Journal Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 53, No. 2

April, 1974

Kenneth Abernethy, <i>On characterizing certain classes of first countable spaces by open mappings</i>	319
Ross A. Beaumont and Donald Lawver, <i>Strongly semisimple abelian groups</i>	327
Gerald A. Beer, <i>The index of convexity and parallel bodies</i>	337
Victor P. Camillo and Kent Ralph Fuller, <i>On Loewy length of rings</i>	347
Stephen LaVern Campbell, <i>Linear operators for which T^*T and TT^* commute. II</i>	355
Charles Kam-Tai Chui and Philip Wesley Smith, <i>Characterization of a function by certain infinite series it generates</i>	363
Allan L. Edelson, <i>Conjugations on stably almost complex manifolds</i>	373
Patrick John Fleury, <i>Hollow modules and local endomorphism rings</i>	379
Jack Tilden Goodykoontz, Jr., <i>Connectedness in kleinen and local connectedness in 2^X and $C(X)$</i>	387
Robert Edward Jamison, II, <i>Functional representation of algebraic intervals</i>	399
Athanassios G. Kartsatos, <i>Nonzero solutions to boundary value problems for nonlinear systems</i>	425
Soon-Kyu Kim, Dennis McGavran and Jingyal Pak, <i>Torus group actions on simply connected manifolds</i>	435
David Anthony Klarner and R. Rado, <i>Arithmetic properties of certain recursively defined sets</i>	445
Ray Alden Kunze, <i>On the Frobenius reciprocity theorem for square-integrable representations</i>	465
John Lagnese, <i>Existence, uniqueness and limiting behavior of solutions of a class of differential equations in Banach space</i>	473
Teck Cheong Lim, <i>A fixed point theorem for families on nonexpansive mappings</i> ...	487
Lewis Lum, <i>A quasi order characterization of smooth continua</i>	495
Andy R. Magid, <i>Principal homogeneous spaces and Galois extensions</i>	501
Charles Alan McCarthy, <i>The norm of a certain derivation</i>	515
Louise Elizabeth Moser, <i>On the impossibility of obtaining $S^2 \times S^1$ by elementary surgery along a knot</i>	519
Gordon L. Nipp, <i>Quaternion orders associated with ternary lattices</i>	525
Anthony G. O'Farrell, <i>Equiconvergence of derivations</i>	539
Dorte Olesen, <i>Derivations of AW^*-algebras are inner</i>	555
Dorte Olesen and Gert Kjærgaard Pedersen, <i>Derivations of C^*-algebras have semi-continuous generators</i>	563
Duane O'Neill, <i>On conjugation cobordism</i>	573
Chull Park and S. R. Paranjape, <i>Probabilities of Wiener paths crossing differentiable curves</i>	579
Edward Ralph Rozema, <i>Almost Chebyshev subspaces of $L^1(\mu; E)$</i>	585
Lesley Millman Sibner and Robert Jules Sibner, <i>A note on the Atiyah-Bott fixed point formula</i>	605
Betty Salzberg Stark, <i>Irreducible subgroups of orthogonal groups generated by groups of root type 1</i>	611
N. Stavrakas, <i>A note on starshaped sets, (k)-extreme points and the half ray property</i>	627
Carl E. Swenson, <i>Direct sum subset decompositions of Z</i>	629
Stephen Tefteller, <i>A two-point boundary problem for nonhomogeneous second order differential equations</i>	635
Robert S. Wilson, <i>Representations of finite rings</i>	643