

Pacific Journal of Mathematics

DIRECT SUM SUBSET DECOMPOSITIONS OF Z

CARL E. SWENSON

DIRECT SUM SUBSET DECOMPOSITIONS OF Z

CARL SWENSON

Let Z be the set of integers. In this paper it is shown that there is no effective characterization of all direct sum subset decompositions of Z i.e., where $A+B=Z$ and the sums are distinct. Further the result is generalized to include decompositions of a product of sets where Z is a set in the product, and to cases where the number of subsets in the decomposition is greater than two.

The question of characterizing all direct sum subset decompositions for Z , the infinite cyclic group, seems first to have been raised explicitly by de Bruijn [1]. It was mentioned again by de Bruijn [2] in 1956, and Long [5] in 1967. The notation $A \oplus B$ will denote $A + B$ where the sums are distinct. Without loss of generality we will assume 0 is a member of each summand.

For the semi-group Z^+ there is a particularly nice characterization of all direct sum decompositions. The result, which was implicit from the work of de Bruijn [2], was first explicitly by Long [5]. It is the following:

THEOREM 1. *Let $|A| = |B| = \infty$. $A \oplus B = Z^+$ if and only if there exists an infinite sequence of integers $\{m_i\}_{i \geq 1}$ with $m_i \geq 2$ for all i , such that A and B are the sets of all finite sums of the form*

$$\begin{aligned} a &= \sum x_{2i} M_{2i} \\ b &= \sum x_{2i+1} M_{2i+1} \end{aligned}$$

respectively, where $0 \leq x_i < m_{i+1}$ for $i \geq 0$ where $M_0 = 1$ and $M_i = \prod_{j=1}^i m_j$ for $i \geq 1$.

In case $|A| < \infty$ or $|B| < \infty$, a similar characterization holds with the change that the sequence $\{m_i\}$ will be of finite length r and the only restriction on x_r is that it be nonnegative.

A distinguishing characteristic of decompositions obtained as in Theorem 1 is that either A or B has the property that each of its elements is a multiple of some integer $m \geq 2$ and it has been conjectured that this property would necessarily hold for any decomposition of Z . The following theorem shows that this is not the case and that the decomposing sets A and B can be quite arbitrary. It follows that there is no real possibility of effectively characterizing A and B . We do obtain a rather weak characterization in Theorem 3.

Throughout this paper, unless otherwise noted, all maximums and minimums will be taken with respect to the following order $0 < 1 < -1 < 2 < -2 < 3 < \dots$.

THEOREM 2. *Suppose that A_1 and B_1 are finite, that $0 \in A_1 \cap B_1$, and that $A_1 + B_1 = A_1 \oplus B_1$, then there exist sets A and B , both infinite, such that $A_1 \subset A$, $B_1 \subset B$, and $A \oplus B = Z$.*

Proof. We first let

$$n_1 = \min(Z \sim (A_1 \oplus B_1))$$

and

$$m_1 = |\max(A_1 \cup B_1 \cup \{n_1\})|$$

where the min and max are taken with respect to the previously mentioned order. We now construct A and B by an inductive procedure. Let

$$A_2 = A_1 \cup \{n_1 + 5m_1\}$$

and

$$B_2 = B_1 \cup \{-5m_1\}$$

then

$$A_2 + B_2 = (A_1 \oplus B_1) \cup (\{n_1 + 5m_1\} + B_1) \cup (\{-5m_1\} + A_1) \cup \{n_1\}.$$

We now claim that $A_2 + B_2 = A_2 \oplus B_2$. Of course, this is immediate if the sets in the above union are mutually disjoint. The fact that they are disjoint is assured by the following inequalities which derive from the definitions of n_1 and m_1 and the fact $m_1 > 0$.

$$\begin{aligned} -5m_1 + a &\leq -4m_1 < -2m_1 \leq a' + b \\ n_1 + 5m_1 + b &\geq 3m_1 > 2m_1 \geq a + b' \\ n_1 + 5m_1 + b &> 0 > (-5m_1) + a \\ n_1 + 5m_1 + b &\geq 3m_1 > |n_1| \\ |-5m_1 + a| &\geq 4m_1 > |n_1| \end{aligned}$$

for all $a, a' \in A_1$ and $b, b' \in B_1$.

Note that $n_1 \in A_2 \oplus B_2$ so that we have enlarged the interval about the origin in which all integers are represented. Also it is clear the process can be repeated to obtain sets A_i and B_i for $i \geq 2$ which are supersets of A_{i-1} and B_{i-1} and such that $A_i + B_i = A_i \oplus B_i$. Setting $A = \bigcup_{i=1}^{\infty} A_i$ and $B = \bigcup_{i=1}^{\infty} B_i$ we have desired decomposition $A \oplus B = Z$, since any given n is an element of $A_{2^{n+1}} \oplus B_{2^{n+1}}$. This completes the proof.

Theorem 2 shows that any two finite sets A and B with $A \subset Z$, $B \subset Z$ and $A + B = A \oplus B$ can be extended to two infinite sets which are a direct sum decomposition of Z . This shows that it is certainly not necessarily the case that every element of A or B has a multiple of some integer $m \geq 2$. It also shows that no condition can be placed on the size or location of the two elements which sum to a given n . Thus, a best possible type of characterization of decomposition of Z will be of the nature of Theorem 3. Let $A(k) = \{a \in A \mid |a| \leq |k|\}$.

THEOREM 3. $A \oplus B = Z$ if and only if for each $n \in Z$ there exists $k \in Z$ such that

$$A(k) \oplus B(k) \supset Z(n).$$

Proof. Suppose first that $A \oplus B = Z$ and let $n \in Z$. For each $i \in Z(n)$, set

$$k_i = \max\{a_i, b_i \mid a_i + b_i = i, a_i \in A, b_i \in B\}.$$

Also, set $k = \max\{k_i \mid i \in Z(n)\}$. Then, since $i \in A(k_i) + B(k_i)$, $A(k_i) \subset A(k)$ and $B(k_i) \subset B(k)$ for all $i \in Z(n)$, it follows that

$$Z(n) \subset A(k) + B(k) = A(k) \oplus B(k).$$

The last equality follows from the fact that $A(k) \subset A$, $B(k) \subset B$ and $A + B = A \oplus B$.

Conversely, suppose that for each $n \in Z$ there exists k_n such that

$$A(k_n) \oplus B(k_n) \supset Z(n).$$

If we set $A = \bigcup_{n=1}^{\infty} A(k_n)$ and $B = \bigcup_{n=1}^{\infty} B(k_n)$, then clearly $n \in A + B$ for any $n \in Z$; i.e., $Z \subset A + B$. Since $A + B \subset Z$ is trivially true, it follows that $Z = A + B$. If $A + B \neq A \oplus B$, then there exist positive integers i and j and an integer n such that

$$n = a + b = a_i + b_j$$

with $a \in A(k_n)$, $b \in B(k_n)$, $a_i \in A(k_i)$, $b_j \in B(k_j)$, $a \neq a_i$, and $b \neq b_j$. But then, if $k' = \max\{k_i, k_j, k_n\}$, it is clear that

$$A(k_p) + B(k_q) \subset A(k') + B(k')$$

for p and $q \in \{i, j, n\}$, and this implies that n has two representations in $A(k') + B(k')$ in violation to the fact

$$A(k') + B(k') = A(k') \oplus B(k').$$

This completes the proof.

We now consider the remaining case for $A \oplus B = Z$ when one

of A and B is finite. We assume without loss of generality that $|A| < \infty$.

THEOREM 4. *Let $|A| < \infty$, then $A \oplus B = Z$ if and only if there exists an n such that $B = nZ \oplus C$ where $C \subset \{0, 1, \dots, n-1\}$ and $A \oplus C$ is a complete residue system modulo n .*

Proof. Since $A \oplus C$ is a complete residue system modulo n , it is clear that

$$Z = (A \oplus C) \oplus nZ = A \oplus (C \oplus nZ) = A \oplus B.$$

Conversely suppose $A \oplus B = Z$ with A of finite order. Under these conditions, Hajós [3] proved that B is periodic, i.e., there exists an $n \neq 0$ such that $n + B = B$. Since $0 \in B$ we have $nZ \subset B$. Letting $C = B \cap \{0, 1, \dots, n-1\}$ we have $B = nZ \oplus C$. Since $Z = A \oplus B = (A \oplus C) + nZ$, it is clear that $A \oplus C$ must be a complete residue system modulo n .

2. Generalizations. Consider subsets $D_i \subset Z$ with $0 \in D_i$ for $1 \leq i \leq n$, then a generalization of Theorem 2 is obtained by replacing Z with $Z \times D_1 \times \dots \times D_k$. The method of proof is similar except that the order is replaced by $(x_0, \dots, x_k) > (y_0, y_1, \dots, y_k)$ if $\sum |x_i| > \sum |y_i|$ or, in case $\sum |x_i| = \sum |y_i|$, $(x_0, \dots, x_k) > (y_0, \dots, y_k)$ if $x_i > y_i$ for the least i such that $x_i \neq y_i$. The least element in the ordering for which there is no representation by $A_1 \oplus B_1$ is n_1 . The m_1 is the maximum over any entry in any element of A_1, B_1 or $\{n_1\}$. In particular, this shows that no strong characterization such as that exhibited for $Z^+ \times Z^+$ by Hanson [4] and Niven [6] can exist for $Z \times Z$ or $Z \times Z^+$.

The preceding theorems all generalize from the case of two summands to the case of any number of summands. Further, since the construction of Theorem 2 is by a single element at a time, the summands can be created with any given order.

REFERENCES

1. N. G. de Bruijn, *On bases for the set of integers*, Publ. Math. Debrecen, **1** (1950), 232-242.
2. ———, *On number systems*, Nieuw Arch. Wisk., **4** (1956), 15-17.
3. G. Hajós, *Sur la factorisation des groupes abéliens*, Casopis Pest. Mat. Fys., **74** (1950), 157-162.
4. R. T. Hanson, *Complementing pairs of subsets in the plane*, Duke Math. J., **36** (1969), 441-449.
5. C. T. Long, *Addition theorems for sets of integers*, Pacific J. Math., **22** (1967), 107-112.

6. I. Niven, *A characterization of complementing sets of pairs of integers*, Duke Math. J., **38** (1971), 193-203.

Received March 8, 1973. The results constitute a part of the author's doctoral dissertation written under the supervision of Professor C. T. Long.

CLAFLIN COLLEGE
ORANGEBURG, SC.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. Index to Vol. **39**. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific of Journal Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Kenneth Abernethy, <i>On characterizing certain classes of first countable spaces by open mappings</i>	319
Ross A. Beaumont and Donald Lawver, <i>Strongly semisimple abelian groups</i>	327
Gerald A. Beer, <i>The index of convexity and parallel bodies</i>	337
Victor P. Camillo and Kent Ralph Fuller, <i>On Loewy length of rings</i>	347
Stephen LaVern Campbell, <i>Linear operators for which T^*T and TT^* commute.</i> <i>II</i>	355
Charles Kam-Tai Chui and Philip Wesley Smith, <i>Characterization of a function by certain infinite series it generates</i>	363
Allan L. Edelson, <i>Conjugations on stably almost complex manifolds</i>	373
Patrick John Fleury, <i>Hollow modules and local endomorphism rings</i>	379
Jack Tilden Goodykoontz, Jr., <i>Connectedness in kleinen and local connectedness in 2^X and $C(X)$</i>	387
Robert Edward Jamison, II, <i>Functional representation of algebraic intervals</i>	399
Athanasios G. Kartsatos, <i>Nonzero solutions to boundary value problems for nonlinear systems</i>	425
Soon-Kyu Kim, Dennis McGavran and Jingyal Pak, <i>Torus group actions on simply connected manifolds</i>	435
David Anthony Klarner and R. Rado, <i>Arithmetic properties of certain recursively defined sets</i>	445
Ray Alden Kunze, <i>On the Frobenius reciprocity theorem for square-integrable representations</i>	465
John Lagnese, <i>Existence, uniqueness and limiting behavior of solutions of a class of differential equations in Banach space</i>	473
Teck Cheong Lim, <i>A fixed point theorem for families on nonexpansive mappings</i> ..	487
Lewis Lum, <i>A quasi order characterization of smooth continua</i>	495
Andy R. Magid, <i>Principal homogeneous spaces and Galois extensions</i>	501
Charles Alan McCarthy, <i>The norm of a certain derivation</i>	515
Louise Elizabeth Moser, <i>On the impossibility of obtaining $S^2 \times S^1$ by elementary surgery along a knot</i>	519
Gordon L. Nipp, <i>Quaternion orders associated with ternary lattices</i>	525
Anthony G. O'Farrell, <i>Equiconvergence of derivations</i>	539
Dorte Olesen, <i>Derivations of AW^*-algebras are inner</i>	555
Dorte Olesen and Gert Kjærsgaard Pedersen, <i>Derivations of C^*-algebras have semi-continuous generators</i>	563
Duane O'Neill, <i>On conjugation cobordism</i>	573
Chull Park and S. R. Paranjape, <i>Probabilities of Wiener paths crossing differentiable curves</i>	579
Edward Ralph Rozema, <i>Almost Chebyshev subspaces of $L^1(\mu; E)$</i>	585
Lesley Millman Sibner and Robert Jules Sibner, <i>A note on the Atiyah-Bott fixed point formula</i>	605
Betty Salzberg Stark, <i>Irreducible subgroups of orthogonal groups generated by groups of root type 1</i>	611
N. Stavrakas, <i>A note on starshaped sets, (k)-extreme points and the half ray property</i>	627
Carl E. Swenson, <i>Direct sum subset decompositions of Z</i>	629
Stephen Tefteller, <i>A two-point boundary problem for nonhomogeneous second order differential equations</i>	635
Robert S. Wilson, <i>Representations of finite rings</i>	643