

Pacific Journal of Mathematics

**CERTAIN REPRESENTATIONS OF INFINITE GROUP
ALGEBRAS**

INDRANAND SINHA

CERTAIN REPRESENTATIONS OF INFINITE GROUP ALGEBRAS

I. SINHA

For any group G , let ρ be an irreducible representation of the group algebra $\mathfrak{F}G$ over a field \mathfrak{F} . Then by Schur's lemma, the center \mathcal{A} of its commuting ring, is a field containing \mathfrak{F} . If ρ is finite-dimensional over \mathcal{A} , then it is called finite and if it is finite-dimensional over \mathfrak{F} itself, then it is called strongly finite. In this paper, certain conditions are given for finiteness of ρ . Also it is shown that for some types of groups, finiteness of ρ is related to the existence of abelian subgroups of finite index in certain quotient of the group. Conditions under which finiteness and strongly finiteness are equivalent, are given. Finally, consequences of ρ being faithful on G , or being faithful on $\mathfrak{F}G$, are studied.

Study of finiteness of irreducible representations was initiated by Kaplansky in [3], and later carried to a great extent by Passman, Issacs, and others: {see [5] and relevant references therein}. Finiteness and strong finiteness were studied in [6]. Using a slight modification of the technique of [4] to suit our nonsemisimple case, we get Theorem 1 which includes the results of [4] and gives us Theorem 2 whose corollaries contains the result of [3].

We further recall the well-known result that for a finite group G , if the kernel of an irreducible representation ρ contains the commutator subgroup G' , then the representation is 1-dimensional over \mathcal{A} . As corollary to our Theorem 3, we prove that in general, if G' is contained in the kernel of ρ , then ρ is finite whether G is finite or not.

2. Finiteness of representation. In this section we study conditions under which a given irreducible representation is finite, and also the conditions under which all irreducible representations are finite. We need the following:

DEFINITIONS. 1. Let ρ be a representation of $\mathfrak{F}G$. Then $G_\rho = \{g \in G \mid \rho(g) = 1\}$, and $\text{Kern } \rho = \text{kernel } \rho = \{x \in \mathfrak{F}G \mid \rho(x) = 0\}$.

Thus ρ is G -faithful if $G_\rho = 1$, while ρ is $\mathfrak{F}G$ -faithful if $\text{Kern } \rho = 0$.

2. Let $\mathfrak{B} \leq \text{Aut } G$. For $S \leq G$, we shall write $\mathfrak{A}_\mathfrak{B}(S)$ for the left-ideal $\{\sum x_i(\mathcal{S}_i^{\beta_i} - 1) \mid x_i \in \mathfrak{F}G, \mathcal{S}_i \in S, \beta_i \in \mathfrak{B}\}$. {For a general study of such ideals we may refer to [6] and [8].} We write $\mathfrak{A}(S)$, if $\mathfrak{B} = \{\text{identity}\}$.

3. We also define the \mathfrak{B} -kernel of ρ in $H \leq G$ to be

$$\{h \in H \mid \rho(h^\beta) = 1, \forall \beta \in \mathfrak{B}\},$$

and set $K_n^{\mathfrak{B}}(H) = \bigcap \{\mathfrak{B}\text{-kernels of } \rho \text{ in } H\}$, where the intersection runs through all irreducible representations ρ of G for which $\dim_{\Delta} \rho > n^2$, where Δ is the center of the commuting ring of ρ . If no such ρ exists, then we put $K_n^{\mathfrak{B}}(H) = G$.

4. Let S_{2n} be the symmetric group of degree $2n$. Then for an algebra A , the sums

$$\sigma_n = \sum_{\sigma \in S_{2n}} (\text{sgn } \sigma) x_{\sigma(1)} \cdot x_{\sigma(2)} \cdots x_{\sigma(2n)}, x_{\rho} \in A,$$

are called the Standard Monomial Sums (of parity n).

5. Define $\Sigma_n(G)$ to be the \mathfrak{F} -space spanned by all the σ_n in $\mathfrak{F}G$. {We shall frequently write Σ_n wherever the group in question is clear from context.}

This Σ_n plays a significant role in determining the degrees of irreducible representations.

Specifically we have:

PROPOSITION 1. *Let ρ be an irreducible representation of $\mathfrak{F}G$. Then $\dim_{\Delta} \rho \leq n^2$ if and only if $\Sigma_n \subseteq \text{Kern } \rho$.*

Proof. If $\dim_{\Delta} \rho \leq n^2$ then $\mathfrak{F}G/\text{Kern } \rho$ is a primitive algebra of matrices of $\dim n$ over Δ . By Theorem 1 of [1], for any $\sigma_n \in \mathfrak{F}G$, $\rho(\sigma_n) = 0$, whence $\sigma_n \in \text{Kern } \rho$ so that $\Sigma_n \subseteq \text{Kern } \rho$.

Conversely, suppose $\Sigma_n \subseteq \text{Kern } \rho$. Then $\mathfrak{F}G/\text{Kern } \rho$ is a primitive algebra satisfying $\sigma_n = 0$ for every σ_n in $\mathfrak{F}G/\text{Kern } \rho$. Then by Theorem 1 of [2], p. 226, $\mathfrak{F}G/\text{Kern } \rho$ is a central simple algebra of $\dim \leq n^2$. Hence $\dim_{\Delta} \rho \leq n^2$.

Using this result we obtain:

THEOREM 1. *Let $S \leq H \leq G$. Then $S \subseteq K_n^{\mathfrak{B}}(H)$ if and only if $\mathfrak{A}_{\mathfrak{B}}(S) \cdot \Sigma_n \subseteq \text{Rad } \mathfrak{F}G$.*

Proof. We observe that $\mathfrak{A}_{\mathfrak{B}}(S) = \{\sum x_i (s_i^{\beta_i} - 1) \mid x_i \in \mathfrak{F}G, s_i \in S, \beta_i \in \mathfrak{B}\}$. Thus, to show that $\mathfrak{A}_{\mathfrak{B}}(S) \cdot \Sigma_n \subseteq \text{Rad } \mathfrak{F}G$, it suffices to show that $(s^\beta - 1) \cdot \Sigma_n \subseteq \text{Rad } \mathfrak{F}G$, for $\forall \beta \in \mathfrak{B}, \forall s \in S$. Now let $h \in K_n^{\mathfrak{B}}(H)$ and ρ be an irreducible representation of G . If $\dim_{\Delta} \rho > n^2$, then $\rho(h^\beta) = 1$ or $\rho(h^\beta - 1) = 0, \forall \beta \in \mathfrak{B}$, by the very definition of $K_n^{\mathfrak{B}}(H)$. On the other hand, if $\dim_{\Delta} \rho \leq n^2$, then by Proposition 1, $\rho(\Sigma_n) = 0$.

Thus, in both cases, $\rho[(h^\beta - 1) \cdot \sum_n] = 0$. Since ρ is arbitrary, so $(h^\beta - 1) \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G$. Hence $S \subseteq K_n^{\mathfrak{B}}(H)$ implies $\mathfrak{U}_{\mathfrak{B}}(S) \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G$.

Conversely, suppose $\mathfrak{U}_{\mathfrak{B}}(S) \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G$. Then, in particular,

$$(s^\beta - 1) \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G, \forall \beta \in \mathfrak{B}, s \in S.$$

We define the left-idealiser [7], of \sum_n into $\text{Rad } \mathfrak{F}G$, by $L_{\text{Rad}(\sum_n)\mathfrak{F}G} = L(\sum_n) = \{x \in \mathfrak{F}G \mid x \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G\}$. This is clearly a left-ideal. Also $[L(\sum_n) \cdot g] \cdot \sum_n = L(\sum_n)[g \cdot \sum_n \cdot g^{-1}] \cdot g = [L(\sum_n) \cdot \sum_n] \cdot g \subseteq \text{Rad } \mathfrak{F}G \cdot g = \text{Rad } \mathfrak{F}G$, for $\forall g \in G$. Hence $L(\sum_n)$ is a two-sided ideal of $\mathfrak{F}G$.

Now let ρ be an irreducible representation of G , afforded by the $\mathfrak{F}G$ -module \mathfrak{M} . Since $L(\sum_n)$ is a two-sided ideal, so $\text{Ann } L(\sum_n) = \{m \in \mathfrak{M} \mid L(\sum_n)m = 0\}$ is an $\mathfrak{F}G$ -submodule of \mathfrak{M} . Thus either $\text{Ann } L(\sum_n) = 0$ or \mathfrak{M} . Now assume that $\dim_{\Delta} \rho > n^2$. Again, by Proposition 1, $\rho(\sum_n) \neq 0$ so that $\sum_n \cdot \mathfrak{M} \neq 0$, whence $\sum_n \not\subseteq \text{Rad } \mathfrak{F}G$. But $L(\sum_n) \cdot [\sum_n \cdot \mathfrak{M}] = [L(\sum_n) \cdot \sum_n] \cdot \mathfrak{M} = 0$, since $L(\sum_n) \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G$. Thus $\text{Ann } L(\sum_n) = \mathfrak{M}$. Then, as by hypothesis $s^\beta - 1 \in L(\sum_n)$, so $(s^\beta - 1) \cdot \mathfrak{M} = 0$; or $\rho(s^\beta - 1) = 0$. As ρ was arbitrary with $\dim_{\Delta} \rho > n^2$; so $s \in K_n^{\mathfrak{B}}(H)$.

Letting $\mathfrak{F} = \text{complex-field}$, we have $\text{Rad } \mathfrak{F}G = 0$. Taking $\mathfrak{B} = \{1\}$ in this case, we obtain the result of Passman [4]:

COROLLARY. $g \in K_n(G)$ if and only if $(g - 1) \cdot \sum_n = 0$.
We also deduce:

THEOREM 2. Let $S \leq G$ and $\mathfrak{B} \leq \text{Aug } G$ such that $S^{\mathfrak{B}} = G$. Then $\mathfrak{U}_{\mathfrak{B}}(S) \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G$ if and only if $\dim_{\Delta} \rho \leq n^2$ for every irreducible representation ρ of G . {Of course, Δ depends on ρ .}

Proof. By Theorem 1, $\mathfrak{U}_{\mathfrak{B}}(S) \cdot \sum_n \subseteq \text{Rad } \mathfrak{F}G$ if and only if $S \subseteq K_n^{\mathfrak{B}}(G)$: $\{G = H\}$.

The latter condition is equivalent to the statement that for every irreducible representation ρ with $\dim_{\Delta} \rho > n^2$, we have $\rho(s^\beta) = 1$, $\forall \beta \in \mathfrak{B}, s \in S$. Since $S^{\mathfrak{B}} = G$, so we deduce that $\rho = 1$.

COROLLARY 1. If $\sum_n \subseteq \text{Rad } \mathfrak{F}G$ for some n , then every irreducible representation of $\mathfrak{F}G$ is finite.

COROLLARY 2. [3]. If in $\mathfrak{F}G$, $\sum_n = 0$ for some n , then every irreducible representation of G is finite.

Next recall that if $|G| < \infty$ then $G' \subseteq G_\rho$ for any irreducible

representation ρ , if and only if ρ is of dim. 1. A generalization of sort, is obtained in the corollary to:

THEOREM 3. *Let ρ be an irreducible representation of $\mathfrak{F}G$.*

(a) *If either (i) $\sum_n(G/G_\rho) = 0$ for some n , or (ii) $\exists A \leq G \ni \cdot$. $G_\rho \leq A$, $|G:A| < \infty$ and A/G_ρ is abelian, then ρ is finite.*

(b) *(Conversely) If ρ is finite and $\mathfrak{F}G$ satisfies either of the following conditions:*

(i) *G/G_ρ is periodic and $\mathfrak{F}(G/G_\rho)$ is nonmodular;*

(ii) *G/G_ρ is periodic with a finite p -Sylow subgroup for Char. $\mathfrak{F} = p \neq 0$;*

(iii) *G/G_ρ satisfies minimum-condition on subgroups; then $\exists A \leq G \ni \cdot$. $G_\rho \leq A$, $|G:A| < \infty$ and A/G_ρ is abelian.*

Proof. (a) Suppose (i) holds. In the notation of [6], $G_\rho = \mathfrak{X}^{-1}(\text{Kern } \rho)$ where for any ideal I of $\mathfrak{F}G$, $\mathfrak{X}^{-1}(I) = \{g \in G \mid g - 1 \in I\}$, and hence $\mathfrak{X}(G_\rho)$ is a sub-ideal in $\text{Kern } \rho$. Since $\mathfrak{X}(G_\rho)$ is the kernel of the linear extension of the canonical map $G \rightarrow G/G_\rho$, so $\mathfrak{F}G/\mathfrak{X}(G_\rho) \cong \mathfrak{F}(G/G_\rho)$. Therefore, $\sum_n(G/G_\rho) = 0$ implies that the standard monomialsum, in $\mathfrak{F}G/\mathfrak{X}(G_\rho)$, all vanish. Now $\mathfrak{F}G/\text{Kern } \rho \cong \mathfrak{F}G/\mathfrak{X}(G_\rho)/\text{Kern } \rho/\mathfrak{X}(G_\rho)$; therefore, the same holds for $\mathfrak{F}G/\text{Kern } \rho$. In particular, $\sum_n(G) \subseteq \text{Kern } \rho$. Then, by Proposition 1, ρ is finite. Next let (ii) hold. Then $|G/G_\rho: A/G_\rho| = n < \infty$, and A/G_ρ is abelian. Therefore, by the result of Kaplansky mentioned before, or by Theorems. 5.1, 8.1 of [5], all the irreducible representations of G/G_ρ are finite.

Now if ρ is afforded by the $\mathfrak{F}G$ -module \mathfrak{M} , then putting $\bar{\rho}(\bar{g}) \cdot m = \rho(g) \cdot m$, for $\bar{g} \in G/G_\rho$, and observing that $G_\rho = \{g \in G \mid \rho(g) = 1\}$, we get a representation $\bar{\rho}$ of G/G_ρ , such that $\bar{\rho}$ is irreducible and the commuting ring of $\bar{\rho}$ in $\text{Hon}_{\mathfrak{F}}(\mathfrak{M}, \mathfrak{M})$, is the same as that of ρ .

Thus the finiteness of $\bar{\rho}$ implies the finiteness of ρ .

(b) $G/G_\rho \cong S \leq GL(n, A)$ and any such S satisfying either of the conditions (i), (ii) or (iii), is abelian by finite: {see [9], Corollaries 9.4, 9.7, 9.8, and 9.23}. We then get our A , by taking the complete inverse-image of the abelian part of G/G_ρ .

Since the group-algebra of an abelian group always satisfies $\sum_n = 0$, so we obtain:

COROLLARY. *If $G' \cong G_\rho$, then ρ is finite.*

3. Strong finiteness of representations. In this section we give a result which shows the equivalence of finiteness and strong-finiteness in certain conditions.

THEOREM 4. *Under either of the following conditions, an irreducible representation ρ of G is finite if and only if it is strongly finite:*

- (i) G is finitely generated;
- (ii) ρ is absolutely irreducible;
- (iii) $\exists H \trianglelefteq G \ni |G:H| < \infty$ and ρ_H has a strongly finite constituent.

Proof. (i) This is the content of Lemma 7 of [6].

(ii) Let the absolutely irreducible finite representation ρ , be afforded by the $\mathfrak{F}G$ -module \mathfrak{M} . Since $\Delta \cong \text{Hom}_{\mathfrak{F}}(\mathfrak{M}, \mathfrak{M})$, $\mathfrak{F} \cong \Delta$, so we can make $\Delta \otimes_{\mathfrak{F}} \mathfrak{M}$ into a ΔG -module by letting $g \cdot (d \otimes m) = d \otimes \rho(g)m$.

Define $\psi: \Delta \otimes_{\mathfrak{F}} \mathfrak{M} \rightarrow \mathfrak{M}$ by $\psi(d \otimes m) = dm$. Since ρ and d commute, so

$$\psi(g \cdot (d \otimes m)) = \psi(d \otimes \rho(g)m) = d(\rho(g)m) = \rho(g)(dm).$$

Thus ψ is a ΔG -homomorphism. Since \mathfrak{M} is absolutely irreducible, so $\Delta \otimes_{\mathfrak{F}} \mathfrak{M}$ is irreducible. So ψ is an isomorphism. Then $\dim_{\Delta}(\Delta \otimes_{\mathfrak{F}} \mathfrak{M}) = \dim_{\Delta} \mathfrak{M} < \infty$. Thus ρ is also finite-dimensional over \mathfrak{F} .

(iii) By Clifford's theorem, $\rho_H = \bigoplus \sum_{i=1}^{[G:H]} \rho_i$, where ρ_i are all conjugate irreducible-representations of H . Hence, if one of them is finite-dimensional over \mathfrak{F} , then so are all; and hence ρ .

4. Faithful representation. Finally, let ρ be a representation (not necessarily irreducible) of the group algebra $\mathfrak{F}G$. For any left-ideal I we shall write ρ^I for the representation afforded by the module $I \cdot \mathfrak{M}$, where ρ is afforded by \mathfrak{M} . We shall let $\mathfrak{A} = \mathfrak{A}(G)$ denote the augmentation-ideal of $\mathfrak{F}G$ and $J = [\mathfrak{F}G, \mathfrak{F}G]$. Let $\text{Char } \mathfrak{F} = p \neq 0$.

We then investigate the consequences of ρ being faithful as a representation of G and as a representation of $\mathfrak{F}G$ respectively. Recalling that if $H \leq G$, then $\mathfrak{A}(H)$ is the left-ideal in $\mathfrak{F}G$ generated by $\{h - 1 \mid h \in H\}$: [6], we have the following:

THEOREM 5. (a) *If ρ is faithful on G , then $A = \mathfrak{A}^{-1}(\text{Kern } \rho^{\mathfrak{A}})$ is an elementary abelian normal p -subgroup which is central if $\mathfrak{A} \cong \text{Kern } \rho^{\mathfrak{A}}$.*

(b) *If ρ is faithful on $\mathfrak{F}G$, then*

- (i) $A = 1$ unless $|G| = 2$, $p = 2$ in which case $A = G$;
- (ii) $B = \mathfrak{A}^{-1}(\text{Kern } \rho^J) = 1$ unless $p = 2$ and $B = G$ is abelian, or $p = 2$, G is nonabelian and B is central.

Proof. (a) Since $\text{Kern } \rho^{\mathfrak{A}} \leq \mathfrak{F}G$ so $A \leq G$.

Let \mathfrak{M} afford ρ so that $\mathfrak{X}\cdot\mathfrak{M}$ affords $\rho^{\mathfrak{a}}$. Hence $g \in A$ if and only if $(g - 1) \cdot \sum_{\substack{x \in G \\ \lambda_x \in \mathfrak{F}}} \lambda_x(x - 1)m = 0$ for each $m \in \mathfrak{M}$. Since $p \neq 0$, so $(g^p - 1)m = (g - 1)^p m = (g - 1)[(g - 1)^{p-1}m] = 0$ as $(g - 1)^{p-1}m \in \mathfrak{X}\cdot\mathfrak{M}$. Thus $g^p m = m$ and faithfulness implies that $g^p = 1$. Further, if $h \in A$ then $(h - 1)m$ and $(g - 1)m$ are both in $\mathfrak{X}\cdot\mathfrak{M}$. Then,

$$(g - 1)(h - 1)m = 0 = (h - 1)(g - 1)m$$

or

$$g^{-1}h^{-1}gh m = m .$$

Again faithfulness gives that $gh = hg$; i.e., A is abelian.

If $\mathfrak{X} \subseteq \text{Kern } \rho^{\mathfrak{a}}$, then $g \in G, h \in A$ implies

$$\begin{aligned} (gh - 1)m &= [(g - 1)(h - 1) + (g - 1) + (h - 1)]m \\ &= (g - 1)m + (h - 1)m = (hg - 1)m . \end{aligned}$$

Thus $gh m = hg m$ and faithfulness gives that $A \subseteq Z(G)$.

(b) (i) Now let ρ be $\mathfrak{F}G$ -faithful. Then the Kern $\rho^{\mathfrak{a}} = \text{Ann } \mathfrak{X}$ in $\mathfrak{F}G$. It is well-known that this annihilator is 0 unless $|G| < \infty$ and $\text{Ann } \mathfrak{X} = \mathfrak{F} \cdot (\sum_{g \in G} g)$. Now if $g_i \in A$, then $g_i - 1 = a \in \text{Kern } \rho^{\mathfrak{a}}$ so that $g_i - 1 = k \cdot \sum g, k \in \mathfrak{F}$. Linear independence of the group elements, gives us that $g_i = 1$, and $k = 0$, or $|G| = 2, g_1 = 1, i = 2, k = 1$, and $+1 = -1$ in \mathfrak{F} .

(ii) Again by faithfulness $\text{Kern } \rho^{\mathfrak{J}} = \text{Ann } J$. So $g \in B$ implies that $(g - 1)(hk - kh) = 0, \forall h, k \in G$, i.e., $ghk - gkh - hk + kh = 0$. If $\text{Char } \mathfrak{F} \neq 2$, then we must either have $ghk = gkh$ in which case $hk = kh$, or $ghk = hk$ in which case $g = 1$.

In case $\text{Char } \mathfrak{F} = 2$ and g is noncentral then choose $k \in G$ such that $gk \neq kg$. Put $h = g^{-1}$. Then the above identity gives,

$$k - gkg^{-1} - g^{-1}k + kg^{-1} = 0 .$$

Since $gkg^{-1} \neq k$, so either $k = g^{-1}k$ or $k = kg^{-1}$, both leading to $g = 1$, a contradiction. Thus in this case $g \in Z(G)$.

REFERENCES

1. S. A. Amitsur and J. Levitzki, *Minimal identities for algebras*, Proc. Amer. Math. Soc., **1** (1950), 449-463.
2. N. Jacobson, *Structure of Rings*, Amer. Math. Soc. Coll. Publ., **28** 1964, Rhode Island, Providence.
3. I. Kaplansky, *Groups with representations of bounded degree*, Canad. Math. J., **1** (1949), 575-580.
4. D. S. Passman, *Character kernels in discrete groups*, Proc. Amer. Math. Soc., **17** (1966), 487-492.
5. ———, *Infinite Group Rings*, Marcel-Dekker Inc., N. Y., 1971.

6. I. Sinha, *On the augmentation maps of the subgroups of a group*, Math. Zeitsch., **94** (1966), 193-206.
7. I. Sinha and J. Srivastava, *Ideals in commutative domains*, Math. Mag., **44** (3), (1971), 124-127.
8. J. Srivastava, Ph. D. Thesis, I. I. T., Kanpur, India.
9. B. A. F. Wehrfritz, *Infinite Linear Groups*, Queen Mary College Math. Notes, 1969.

Received April 18, 1973 and in revised form October 16, 1973.

MICHIGAN STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 54, No. 1

May, 1974

Ralph K Amayo, <i>Engel Lie rings with chain conditions</i>	1
Bernd Anger and Jörn Lembcke, <i>Hahn-Banach type theorems for hypolinear functionals on preordered topological vector spaces</i>	13
Gregory Frank Bachelis and Samuel Ebenstein, <i>On $\Lambda(p)$ sets</i>	35
Harvey Isaac Blau, <i>Indecomposable modules for direct products of finite groups</i>	39
Larry Eugene Bobisud and James Calvert, <i>Singular perturbation of a time-dependent Cauchy problem in a Hilbert space</i>	45
Walter D. Burgess and Robert Raphael, <i>Abian's order relation and orthogonal completions for reduced rings</i>	55
James Diederich, <i>Representation of superharmonic functions mean continuous at the boundary of the unit ball</i>	65
Aad Dijkema and Hendrik S. V. de Snoo, <i>Self-adjoint extensions of symmetric subspaces</i>	71
Gustave Adam Efroymsen, <i>A Nullstellensatz for Nash rings</i>	101
John D. Elwin and Donald R. Short, <i>Branched immersions onto compact orientable surfaces</i>	113
John Douglas Faires, <i>Comparison of the states of closed linear transformations</i>	123
Joe Wayne Fisher and Robert L. Snider, <i>On the von Neumann regularity of rings with regular prime factor rings</i>	135
Franklin Takashi Iha, <i>A unified approach to boundary value problems on compact intervals</i>	145
Palaniappan L. Kannappan and Che Tat Ng, <i>On functional equations connected with directed divergence, inaccuracy and generalized directed divergence</i>	157
Samir A. Khabbaz and Elias Hanna Toubassi, <i>The module structure of $\text{Ext}(F, T)$ over the endomorphism ring of T</i>	169
Garo K. Kiremidjian, <i>On deformations of complex compact manifolds with boundary</i>	177
Dimitri Koutroufiotis, <i>Mappings by parallel normals preserving principal directions</i>	191
W. K. Nicholson, <i>Semiperfect rings with abelian adjoint group</i>	201
Norman R. Reilly, <i>Extension of congruences and homomorphisms to translational hulls</i>	209
Sadahiro Saeki, <i>Symmetric maximal ideals in $M(G)$</i>	229
Brian Kirkwood Schmidt, <i>On the homotopy invariance of certain functors</i>	245
H. J. Shyr and T. M. Viswanathan, <i>On the radicals of lattice-ordered rings</i>	257
Indranand Sinha, <i>Certain representations of infinite group algebras</i>	261
David Smallen, <i>The group of self-equivalences of certain complexes</i>	269
Kalathoor Varadarajan, <i>On a certain problem of realization in homotopy theory</i>	277
James Edward West, <i>Sums of Hilbert cube factors</i>	293
Chi Song Wong, <i>Fixed points and characterizations of certain maps</i>	305