

# Pacific Journal of Mathematics

**CLOSURE THEOREMS FOR AFFINE TRANSFORMATION  
GROUPS**

RICHARD P. GOSSELIN

## CLOSURE THEOREMS FOR AFFINE TRANSFORMATION GROUPS

R. P. GOSSELIN

Let  $\mathcal{H}$  be a closed subgroup of the group of linear transformations of  $R_n$  onto itself. Let  $hx$  denote the image of the point  $x$  under the transformation  $h$ , and let  $\mathcal{G}$  be the transpose group of  $\mathcal{H}$ : i.e. its elements are associated with matrices which are the transposes of those in  $\mathcal{H}$ . For  $f$  in  $L^2(R_n)$ , let  $Cl\{f; \mathcal{H} \times R_n\}$  denote the closure in the  $L^2$  norm of the linear span of functions of the form  $f(hx + t)$  where  $h$  is in  $\mathcal{H}$ , and  $t$  is in  $R_n$ . Since this space is translation-invariant, it is of the form  $L^2(\mathcal{S})$ : i.e. the set of  $L^2$  functions  $r(x)$  such that the nonzero set of  $\hat{r}$ , the Fourier transform of  $r$ , is, except for a set of measure zero, included in  $\mathcal{S}$ . In the first theorem a precise description of  $\mathcal{S}$  is given, and in the second, a function is constructed in a natural way whose translates alone generate the given space.

$\mathcal{S}$  is roughly the orbit of  $N(f)$ , the nonzero set of  $\hat{f}$ , under the group  $\mathcal{G}$ :

$$\bigcup_{g \in \mathcal{G}} g(N(f)) = \mathcal{G}(N(f)).$$

However a difficulty arises in that  $N(f)$  is determined only to within a set of measure zero, and  $\mathcal{G}$  may transform sets of measure zero into nonmeasurable sets. For example, when the rotation group of the plane acts on a nonmeasurable linear set (of the  $x$ -axis, say), a nonmeasurable planar set results. Hence some care is required in defining  $\mathcal{S}$ . Let  $E_f$  denote the set of points of density (one) of  $N(f)$ . Since the exceptional set of  $N(f)$  has measure zero, every point of  $E_f$  has density one with respect to  $E_f$ . The orbit of the set  $E_f$  under the group  $\mathcal{G}$  will be used as  $\mathcal{S}$ , and as part of our first theorem, it will be shown that  $\mathcal{S}$  is measurable.

The fact that closed translation-invariant subspaces of  $L^2$  are of the form  $L^2(\mathcal{S})$  is due to L. Schwartz [3]. The characterization of  $Cl\{f; \mathcal{H} \times R_n\}$  in Theorem 1 reduces to the familiar Wiener theorem when  $\mathcal{H}$  consists only of the identity and has been proved by S. R. Harasymiv for  $L^p$  spaces and for general distribution spaces [1, 2] when  $\mathcal{H}$  is the diagonal group. The second theorem involves the construction of a function  $p$ , arising naturally from  $f$  by an integration over  $\mathcal{G}$ , such that the translates of  $p$  generate the same space: i.e. such that  $Cl\{p; R_n\} = Cl\{f; \mathcal{H} \times R_n\}$ .

## 2. The closure theorem.

**THEOREM 1.** *Let  $\mathcal{H}$  be a closed group of linear transformations of  $R_n$  onto itself with transpose group  $\mathcal{S}$ . For  $f$  in  $L^2(R_n)$ , let  $E_f$  be the set of points of density of  $N(f)$ . Then*

- (i)  $S = \mathcal{S}(E_f)$  is measurable, and
- (ii)  $Cl\{f; \mathcal{H} \times R_n\} = L^2(\hat{S})$ .

For  $x$  in  $\mathcal{S}(E_f)$ , let  $g = g(x)$  be any element of  $\mathcal{S}$  such that  $x$  is in  $g(E_f)$ . For such a point, there exists  $\delta = \delta(g) > 0$  such that, if  $B_r(x)$  is the ball of radius  $r$  about  $x$  as center, then

$$m\{B_r(x) \cap g(E_f)\} \geq 2\delta m\{B_r(x)\}$$

for sufficiently small  $r$ . Here  $m$  denotes Lebesgue measure. Hence  $B_r(x) \cap g(E_f)$  contains a closed set  $A_r(x)$  such that

$$m\{A_r(x)\} \geq \delta m\{B_r(x)\}.$$

It may be assumed that  $x$  itself belongs to  $A_r(x)$ . The sets  $A_r(x)$ ,  $r \leq r(x)$ ,  $x$  in  $\mathcal{S}(E_f)$ , constitute a family of closed sets covering  $\mathcal{S}(E_f)$  in the sense of Vitali. Thus there exists a countable subfamily covering  $\mathcal{S}(E_f)$  except for a set of measure zero. But since each set  $A_r(x)$  is contained in  $\mathcal{S}(E_f)$ , then  $\mathcal{S}(E_f)$  is measurable. This completes the proof of part (i).

For reference below, let  $A_r^{(n)}(x)$ ,  $n = 1, 2, \dots$  denote the countable subcover, and let  $\{g_n\}$  denote the corresponding sequence in  $\mathcal{S}$ . Thus the difference set

$$(1) \quad \mathcal{S}(E_f) \sim \bigcup_n g_n(E_f)$$

is of measure zero.

Now the proof of part (ii) of the theorem follows standard lines. In particular, if  $k$  is orthogonal to the space  $Cl\{f; \mathcal{H} \times R_n\}$ , then  $\hat{k}(x)\hat{f}(g_n x) = 0$  for every  $g_n$  of the sequence in (1). Hence  $\hat{k}$  must vanish for almost every point of  $\mathcal{S}(E_f) = S$ .

**3. An equivalent space of translates.** Since  $L^2(\hat{S})$  is translation-invariant, it is generated by the translates of any function  $p$  such that the sets  $N(p)$  and  $S$  differ by a set of measure zero. Such a function arises naturally from an integration with respect to Haar measure on  $\mathcal{S}$ . Consider  $\hat{f}(gx) = \hat{f}(g, x)$  as a function on the product space  $\mathcal{S} \times R$  where the measure associated with  $\mathcal{S}$  is  $\mu$ , the right invariant Haar measure, and the measure

associated with  $R_n$  is  $m$ , Lebesgue measure. It is shown below that  $\hat{f}(gx)$  is measurable on this product space. Now let  $\beta$  be a strictly positive measurable function of  $\mathcal{S}$  such that  $|g^{-1}|^{1/2} \beta(g)$  is in  $L^1(\mathcal{S})$ . Here  $|g^{-1}|$  denotes the absolute value of the determinant associated with  $g^{-1}$ . Assuming the measurability of the function  $\hat{f}(gx)$ , let us consider the function

$$(2) \quad \hat{p}(x) = \int_{\mathcal{S}} |\hat{f}(gx)| \beta(g) d\mu(g) .$$

In view of Minkowski's inequality,  $\hat{p}$  is in  $L^2(R_n)$ , and its inverse Fourier transform,  $p$ , is the one we wish to consider in our second theorem.

**THEOREM 2.** *Let  $f$  be in  $L^2(R_n)$  and  $S$  the orbit under  $\mathcal{S}$  of the set of points of density of  $N(f)$ . Then*

- (i)  $\hat{f}(gx)$  is measurable on  $\mathcal{S} \times R_n$ , and
- (ii) the function  $p$ , with Fourier transform  $\hat{p}$  defined by (2), is in  $L^2(R_n)$ , and  $Cl\{p; R_n\} = L^2(\hat{S})$ .

Let  $\mathcal{O}$  be an open set in the complex plane. Then  $(\hat{f})^{-1}(\mathcal{O})$  is measurable in  $R_n$  and can be expressed as  $B \cup C$ , where  $B$  is a Borel set and  $C$  has measure zero. Let  $\alpha$  be the map from  $\mathcal{S} \times R_n$  defined as  $\alpha(g, x) = gx$ . We have

$$\alpha^{-1}\{(\hat{f})^{-1}(\mathcal{O})\} = \{\alpha^{-1}(B)\} \cup \{\alpha^{-1}(C)\} .$$

Since  $\alpha$  is continuous,  $\alpha^{-1}(B)$  is a Borel set in  $\mathcal{S} \times R_n$ . It is thus enough to prove that  $\alpha^{-1}(C)$  is of measure zero: i.e. that  $\alpha^{-1}$  is absolutely continuous as a set function. Let  $K$  be a compact set in  $\mathcal{S}$ , and let  $D$  be an open set containing  $C$  such that  $m(D) < \epsilon = \epsilon(K)$ . A routine calculation shows that

$$(\mu \times m)\{\alpha^{-1}(D) \cap (K \times R_n)\} \leq \epsilon \int_K |g^{-1}| d\mu(g) .$$

Thus the outer measure of  $\alpha^{-1}(C) \cap (K \times R_n)$  is zero. Since  $K$  is an arbitrary compact set of  $\mathcal{S}$ ,  $\alpha^{-1}(C)$  is measurable and of measure zero. This completes the proof of (i).

It has already been noted that  $\hat{p}$ , and hence  $p$ , are in  $L^2(R_n)$ . To complete the proof of part (ii), it is enough to show that  $N(p)$ , the nonzero set of  $\hat{p}$ , and  $S$  differ by a set of measure zero. Let  $D$  be the set of points  $x$  such that, for any neighbourhood of  $x$ ,  $\mathcal{N}(x)$ ,

$$\int_{\mathcal{N}(x)} \hat{p}(y) dy > 0 .$$

The sets  $D$  and  $N(p)$  differ by a set of measure zero so that it is enough to show the same for  $D$  and  $S$ .

Our first step is to show that  $D$  is invariant with respect to  $\mathcal{G}$  : i.e.

$$(3) \quad gD \subset D \text{ for all } g \text{ of } \mathcal{G} .$$

Since  $\mathcal{G}$  is a group, this is equivalent to saying that  $gD = D$  for all  $g$  of  $\mathcal{G}$  . It is clear that  $S$  is invariant in this same sense. Since the complement of an invariant set is also invariant, and since the intersection of two invariant sets is invariant, it will follow from (3) that both difference sets,  $S \sim D$  and  $D \sim S$ , are invariant. The proof is then completed by integration of  $\hat{p}$  over these sets.

Let  $x$  be in  $D$ ,  $g_0$  in  $\mathcal{G}$  , and  $\epsilon > 0$ . Since  $\mu$  is right invariant Haar measure, it follows that

$$\int_{|g_0x-y|<\epsilon} \hat{p}(y) dm(y) = |g_0| \int_{|g_0(x-z)|<\epsilon} dm(z) \int_{\mathcal{G}} |\hat{f}(gz)| \beta(gg_0^{-1}) d\mu(g) .$$

By the continuity of  $g_0$ , there is a  $\delta > 0$  such that if  $|x - z| < \delta$ , then  $|g_0(x - z)| < \epsilon$  so that the above integral exceeds

$$(4) \quad |g_0| \int_{|x-z|<\delta} dm(z) \int_{\mathcal{G}} |\hat{f}(gz)| \beta(gg_0^{-1}) d\mu(g) .$$

Since  $\beta(gg_0^{-1}) > 0$  for all  $g$  of  $\mathcal{G}$  , and since

$$\int_{|x-z|<\delta} dm(z) \int_{\mathcal{G}} |\hat{f}(gz)| \beta(g) d\mu(g) > 0$$

the integral of (4) must be positive. Thus  $g_0x$  is also in  $D$ , and so  $D$  is invariant in the sense of (3).

Let  $A = D \sim S$ , which, in view of (3), is invariant. Let  $q$  be a strictly positive function of  $L^2(R_n)$ . Then

$$\int_A \hat{p}(x) q(x) dm(x) = \int_{\mathcal{G}} \beta(g) d\mu(g) \int_A |\hat{f}(gx)| q(x) dm(x) .$$

If this integral is positive, then for some  $g$  of  $\mathcal{G}$  ,  $gA \cap E_f$  has positive measure. Since  $gA = A$ , and since  $E_f \subset S$ , then  $A \cap S$  has positive measure. Since the latter is impossible, the integral is zero. Since  $q(x) > 0$  for all  $x$ , and since  $p(x) > 0$  for almost every  $x$  of  $D$ , this implies that  $A = D \sim S$  has zero measure.

Let  $B = S \sim D$ . Then

$$0 = \int_B |\hat{p}(x)| dm(x) = \int_{\mathcal{G}} \beta(g) |g^{-1}| d\mu(g) \int_{gB} |\hat{f}(x)| dm(x) .$$

Since  $\beta$  is a strictly positive function, then for almost every  $g$  of  $\mathcal{G}$ ,

$$(5) \quad \int_{gB} |\hat{f}(x)| dm(x) = 0.$$

Since  $B$  is invariant, (5) is true for all  $g$  of  $\mathcal{G}$ , with  $gB = B$ . If  $m(B) > 0$ , then for some  $g_n$  of the sequence in (1),  $m(B \cap g_n E_f) > 0$ . By the invariance of  $B$ ,  $m(B \cap E_f) > 0$ . But this contradicts (5) with  $g$  taken as the identity.

#### REFERENCES

1. S. R. Harasymiv, *A note of dilations*, Pacific J. Math., **21** (1967), 493-501.
2. ———, *On approximation by dilations of distributions*, Pacific J. Math., **28** (1969), 363-374.
3. L. Schwartz, *Théorie général des fonctions moyene-périodiques*, Ann. Math., **48** (1947), 857-929.

Received February 21, 1973. The work on this paper was supported by National Science Foundation Grant GP 9053.

UNIVERSITY OF CONNECTICUT



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)

University of California  
Los Angeles, California, 90024

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington  
Seattle, Washington 98105

D. GILLBARG AND J. MILGRAM

Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INST. OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NAVAL WEAPONS CENTER

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate if possible, may be sent to any one of the five editors. Please classify according to the scheme of Math. Rev. Index to Vol. 39. All other communications to the editors should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California 90708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright ©1974 by Pacific Journal of Mathematics  
Manufactured and first issued in the U.S.A.



# Pacific Journal of Mathematics

Vol. 54, No. 2

June, 1974

John Edward Coury, <i>Walsh series with coefficients tending monotonically to zero</i> .....	1
Patrick Michael Fitzpatrick and Walter Volodymyr Petryshyn, <i>Fixed point theorems for multivalued noncompact acyclic mappings</i> .....	17
Irving Leonard Glicksberg, <i>More on Phragmén-Lindelöf for function algebras</i> .....	25
Adilson Goncalves, <i>Structural constants. II</i> .....	39
Richard P. Gosselin, <i>Closure theorems for affine transformation groups</i> ....	53
Ralph Peter Grimaldi, <i>Baer and UT-modules over domains</i> .....	59
Edward Grossman, <i>On the prime ideal divisors of <math>(a^n - b^n)</math></i> .....	73
A. Hedayat and Ester Seiden, <i>On the theory and application of sum composition of Latin squares and orthogonal Latin squares</i> .....	85
Gerald L. Itzkowitz, <i>Continuous measures, Baire category, and uniform continuity in topological groups</i> .....	115
Francis Masat, <i>Right simple congruences on a semigroup</i> .....	127
Robert Harvey Oehmke, <i>Right congruences and semisimplicity for Rees matrix semigroups</i> .....	143
Qazi Ibadur Rahman and Jan Stankiewicz, <i>Differential inequalities and local valency</i> .....	165
William John Reed, <i>Random points in a simplex</i> .....	183
Mohan S. Shrikhande, <i>Strongly regular graphs and group divisible designs</i> .....	199
Zahava Shmuely, <i>The structure of Galois connections</i> .....	209
Robert C. Shock, <i>Dual generalizations of the Artinian and Noetherian conditions</i> .....	227
Arne Stray, <i>Approximation and interpolation for some spaces of analytic functions in the unit disc</i> .....	237
Eldon Jon Vought, <i>Monotone decompositions into trees of Hausdorff continua irreducible about a finite subset</i> .....	253
James Wirth, <i>The mapping cylinder axiom for WCHP fibrations</i> .....	263
Gordon S. Woodward, <i>Invariant means and ergodic sets in Fourier analysis</i> .....	281