Pacific Journal of Mathematics

THE BACK-AND-FORTH ISOMORPHISM CONSTRUCTION

DALE WAYNE MYERS

Vol. 55, No. 2

October 1974

THE BACK-AND-FORTH ISOMORPHISM CONSTRUCTION

DALE MYERS

Frequently in model theory and occasionally elsewhere, a back-and-forth construction is used to show that any two countable structures satisfying a given relation are isomorphic. Such a construction is used to show that any two countable dense linear orders without end points are isomorphic (Cantor), that any two countable reduced *p*-primary abelian groups with the same Ulm invariants are isomorphic (see Kaplansky, *Infinite Abelian Groups*), and that any two countable elementarily equivalent saturated structures are isomorphic (Morley and Vaught). The back-and-forth arguments using these constructions can often be reduced to an application of the following result: If *R* is a symmetric relation between countable structures such that (1) \mathfrak{ARB} implies \mathfrak{A} and \mathfrak{B} satisfy the same atomic sentences and (2) \mathfrak{ARB} and $a \in \mathfrak{A}$ implies there is a $b \in \mathfrak{B}$ such that (\mathfrak{A}, a) $R(\mathfrak{B}, b$), then \mathfrak{ARB} implies $\mathfrak{A} \cong \mathfrak{B}$.

Loosely, the second condition requires that related structures have enough related expansions by constants. We prove a similar result in which the second condition requires, loosely, that related structures have enough similar decompositions into related components. The prototype of our result is a theorem of Vaught's on Boolean algebras mentioned in the last section. In order to suitably formalize "decomposition" we use category theory.

In a subsequent paper, our two theorems will be used to solve the problem of determining when two free locally-finite cylindric algebras are isomorphic. In particular, the answer to both questions in problem 2.8 of [4, p. 463] is no. The answer to the first is yes if $|\Delta\xi| = \kappa$ and $|\Delta'\xi| = \kappa$ are replaced by $|\Delta\xi| \ge \kappa$ and $|\Delta'\xi| \ge \kappa$ respectively.

Constructors and decomposable relations. We shall use [7] as our notational reference. A directed category is a small category generated by a preorder for which every pair of elements has an upper bound. Let \underline{C} be a category whose arrows are monics. Let \underline{C}^* be the category whose objects are functors from directed categories to \underline{C} and for which an arrow from an object $A: \alpha \to \underline{C}$ to an object $B: \beta \to \underline{C}$ is an ordered pair $\langle f, \tau \rangle$ such that $f: \alpha \to \beta$ is a functor and $\tau: A \to$ $f \circ B$ is a natural transformation. The components of $\langle f, \tau \rangle: A \to B$ are those of τ , i.e., the maps τ_i for i in the domain of A. If $\langle f$, $\tau
angle$: $A \to B$ and $\langle g, \sigma \rangle$: $A \to D$ and *i* is in the domain of *A*, then τ_i and σ_i are corresponding components. By identifying an object *a* of \underline{C} with the functor $1 \to \underline{C}$ whose unique object value is *a*, we can consider \underline{C} to be a subcategory of \underline{C}^* . For an object *A* of \underline{C}^* , an object $\bigcup A$ of \underline{C} is a colimit of *A* iff there is an arrow $A \to \bigcup A$ which is universal from *A* to \underline{C} . Such an arrow is called a *universal* cone. An object *c* in \underline{C} is compact iff for every universal cone $A \to$ $\bigcup A$, every $c \to \bigcup A$ factors through $A \to \bigcup A$; an arrow is compact iff its domain is. (See the last section for examples.) A category is compactly generated iff for any two arrows $a \to_1 b$ and $a \to_2 b$, $a \to_1 b \neq$ $a \to_2 b$ implies $c \to a \to_1 b \neq c \to a \to_2 b$ for some compact arrow $c \to a$. An object in \underline{C} is countable iff it is a colimit $\bigcup C$ of some *C* in \underline{C}^*

A partial isomorphism between objects a and b of \underline{C} is a pair of arrows $a \leftarrow d \rightarrow b$ in \underline{C} (recall that arrows of \underline{C} are monics). Two objects a and b are partially isomorphic, written $a \cong_p b$, iff there is a nonempty set I of partial isomorphisms between a and b such that for every $a \leftarrow d \rightarrow b$ in I, every compact arrow to a or b factors through an extension in I of $a \leftarrow d \rightarrow b$, i.e.,



is completable with $a \leftarrow d' \rightarrow b$ in *I*. If *I* is as above, we write $I: a \cong_p b$. This definition is due to Karp [6]; see [1] for details. Two objects are partially isomorphic iff player II has a winning strategy in an appropriate Ehrenfeucht-Fraïssé game.

For any functor $\varphi: \underline{C} \times \cdots \times \underline{C} \to \underline{C}$ and arrows $a_1 \to b_1, \cdots, a_n \to b_n$, let $\varphi(a_1, \cdots, a_n) \to \varphi(b_1, \cdots, b_n)$ be the arrow $\varphi(a_1 \to b_1, \cdots, a_n \to b_n)$. A functor $\varphi: \underline{C} \times \cdots \times \underline{C} \to \underline{C}$ is a constructor iff for any objects $a_1, \cdots, a_n \in \underline{C}$, every compact arrow $c \to \varphi(a_1, \cdots, a_n)$ factors through $\varphi(c_1, \cdots, c_n) \to \varphi(a_1, \cdots, a_n)$ for some compact arrows $c_1 \to a_1, \cdots, c_n \to a_n$. Zero-ary functors $1 \to \underline{C}$ are also regarded as constructors. Evidently the identity functor on \underline{C} is a constructor and a composition of constructors is a constructor. For any set φ of constructors let $\overline{\phi}$ be the smallest set including φ and the identity functor on \underline{C} and closed under composition.

It is not hard to verify that if every object of \underline{C} is a directed colimit of compacts, then φ is a constructor iff φ preserves directed colimits.

Suppose each component of \underline{C} has an initial object. Let Φ be a set of constructors on \underline{C} . For any $\varphi \in \Phi$, any objects $a_1, \dots, a_n \in \underline{C}$ whose components have initial objects p_1, \dots, p_n respectively, let $\varphi(p_1, \dots, p_n) \to \varphi(a_1, \dots, a_n)$ be $\varphi(p_1 \to a_1, \dots, p_n \to a_n)$ where $p_i \to a_i$ is the unique arrow from p_i to a_i . A symmetric binary relation R on the objects of \underline{C} is Φ -decomposable iff for any objects a and b in \underline{C} (1) aRb implies a and b are in the same component of \underline{C} and (2) aRband $c \to a$ compact implies there is a φ in $\overline{\Phi}, a_1, \dots, a_n, b_1, \dots, b_n$ in \underline{C} , and isomorphisms $a \cong \varphi(a_1, \dots, a_n)$ and $\varphi(b_1, \dots, b_n) \cong b$ such that a_1Rb_1, \dots, a_nRb_n , and $c \to a$ factors through $\varphi(p_1, \dots, p_n) \to \varphi(a_1, \dots, a_n)$ a where p_i is an initial object of the common component of a_i and b_i . (See the last section for examples.)

The back-and-forth theorems. The following lemma is related to the final functor theorem [7, p. 213].

LEMMA 1. For any compactly generated category <u>C</u> of monics, any A in <u>C</u>^{*} with universal cone $A \rightarrow \bigcup A$, and any $B \rightarrow \bigcup A$ in <u>C</u>^{*}, if every component of $A \rightarrow \bigcup A$ factors through a component of $B \rightarrow \bigcup A$, then $\bigcup A$ is a colimit of B.

Proof. Suppose \underline{C} , $A \to \bigcup A$, and $B \to \bigcup A$ are as hypothesized. We claim that $B \to \bigcup A$ is a universal cone. Given $B \to d$, let $A \to d$ be the unique arrow such that for any $A \to \bigcup A$ component $a \to \bigcup A$, the corresponding $A \to d$ component is $a \to d = a \to b \to d$ where (*) $b \to d$ is a $B \to d$ component and $a \to b$ is an arrow such that $a \to b \to$ $\bigcup A = a \to \bigcup A$ where $b \to \bigcup A$ is the $B \to \bigcup A$ component corresponding to $b \to d$. To show that $a \to d$ is independent of the choice of b, let $a \to b_1 \to d$ and $a \to b_2 \to d$ satisfy (*). Thus, in the diagram below with $c \to a$, $c \to b_1$, and $\bigcup A \to d$ omitted, the triangles H commute by hypothesis. Since B has a directed domain, there are arrows $b_1 \to b_3 \leftarrow b_2$ and components $b_3 \to \bigcup A$ and $b_3 \to d$ such that the triangles N commute by naturality.

Hence $a \to b_1 \to b_3 \to \bigcup A = a \to b_2 \to b_3 \to \bigcup A$ and, since $b_3 \to \bigcup A$ is monic, $a \to b_1 \to b_3 = a \to b_2 \to b_3$. Hence $a \to b_1 \to d = a \to b_2 \to d$. Finally, these components "add up" to an arrow (cone) $A \to d$ since if A assigns $a \to a'$ to an arrow of its domain, if $a \to \bigcup A$ and $a' \to \bigcup A$ are the associated $A \to \bigcup A$ components and if $a \to d = a \to b \to d$ and $a' \to d = a' \to b' \to d$ are the corresponding arrows satisfying (*); then, letting $a \to b' = a \to a' \to b'$ and noting that $a \to b' \to d$ satisfies (*), we have $a \to a' \to d = a \to a' \to b' \to d = a \to b' \to d = a \to d$, the last equality due to the independence of $a \to d$ from the choice of b. By universality of $A \to \bigcup A$, there is a $\bigcup A \to d$ such that $A \to d$



 $\bigcup A \to d = A \to d$. We show that $B \to \bigcup A \to d = B \to d$. Since <u>C</u> is compactly generated, it suffices to show that $c \rightarrow b \rightarrow \bigcup A \rightarrow d = c \rightarrow d$ $b \rightarrow d$ for every pair $b \rightarrow \bigcup A$ and $b \rightarrow d$ of corresponding components of $B \to \bigcup A$ and $B \to d$ and every compact arrow $c \to b$. Suppose $b_1 \rightarrow \bigcup A$ and $b_1 \rightarrow d$ are corresponding components of $B \rightarrow \bigcup A$ and $B \rightarrow d$ and $c \rightarrow b_1$ is compact. By compactness $c \rightarrow b_1 \rightarrow \bigcup A = c \rightarrow d$ $a \to \bigcup A$ for some $A \to \bigcup A$ component $a \to \bigcup A$. Let $a \to d$ be the $A \rightarrow d$ component corresponding to $a \rightarrow \bigcup A$. Then by definition of $A \rightarrow d, a \rightarrow d = a \rightarrow b_2 \rightarrow d$ for some $B \rightarrow d$ component $b_2 \rightarrow d$ and some $a \rightarrow b_2$ such that $a \rightarrow b_2 \rightarrow \bigcup A = a \rightarrow \bigcup A$. Thus in the diagram above, this time with $a \rightarrow b_1$ omitted, the triangle H commutes by hypothesis and the triangle C by choice of $\bigcup A \rightarrow d$. Since B has a directed domain, there are arrows $b_1 \rightarrow b_3 \leftarrow b_2$ and components $b_i \rightarrow \bigcup A$ and $b_i \rightarrow d$ such that the triangles N commute by naturality. Now $c \rightarrow d$ $b_1 \rightarrow b_3 \rightarrow \bigcup A = c \rightarrow b_1 \rightarrow \bigcup A = c \rightarrow a \rightarrow \bigcup A = c \rightarrow a \rightarrow b_2 \rightarrow b_3 \rightarrow \bigcup A$ and $b_3 \rightarrow \bigcup A$ is monic so $c \rightarrow b_1 \rightarrow b_3 = c \rightarrow a \rightarrow b_2 \rightarrow b_3$ and the quintilateral Q commutes. Using the commutativity of the triangles H, N, C, and ab_2d and the quadrilateral $cb_1 \bigcup Aa, c \rightarrow b_1 \rightarrow \bigcup A \rightarrow d = c \rightarrow d$ $a \rightarrow \bigcup A \rightarrow d = c \rightarrow a \rightarrow d = c \rightarrow a \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_3 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow a \rightarrow b_2 \rightarrow d = c \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow b_1 \rightarrow b_2 \rightarrow d = c \rightarrow b_2$ $c \rightarrow b_1 \rightarrow b_3 \rightarrow d = c \rightarrow b_1 \rightarrow d$. Hence $B \rightarrow \bigcup A \rightarrow d = B \rightarrow d$.

To see that $\bigcup A \to d$ is the unique such arrow, use the same diagram, this time with c, b_1 , and b_3 omitted and $\bigcup A \to d$ replaced by $\bigcup A \to d$, to show that if $B \to \bigcup A \to d = B \to d$, then $A \to \bigcup A \to d = A \to d$ and hence, by the universality of $A \to \bigcup A, \bigcup A \to d' = \bigcup A \to d$.

THEOREM 2. (Countable and \cong_{p} implies \cong .) If a and b are countable objects in a compactly generated category of monics, then $a \cong_{p} b$ implies $a \cong b$.

Proof. Let $A \to a$ and $B \to b$ be universal cones from functors with countable domains and compact values; let $a_1 \to a, a_3 \to a, a_5 \to a, \cdots$ and $b_2 \to b, b_4 \to b, \cdots$ be enumerations of their (compact) components; and let $I: a \cong_p b$. Let $a \leftarrow d_0 \to b$ be any element of I. Given arrows $a \leftarrow d_k \to b$ in I and $d_{k-1} \to d_k, k = 1, \cdots, i-1$, such that the solid part of the diagram below commutes, let $a \leftarrow d_i \to b$ in I be an extension of $a \leftarrow d_{i-1} \to b$ via some $d_{i-1} \to d_i$ such that $a_i \to a$, if i is odd, or $b_i \to b$, if i is even, factors through $a \leftarrow d_i \to b$.



Let D in \underline{C}^* be the functor from the directed category $\boldsymbol{\omega}$ for which $D(i) = d_i$ and $D(i \to i + 1) = d_i \to d_{i+1}$, i.e., $D = "d_0 \to d_1 \to d_2 \cdots$ ", and let $D \to a$ be the arrow of \underline{C}^* whose *i*th component is $d_i \to a$. By construction, every component of $A \to a$ factors through a component of $D \to a$ and so, by Lemma 1, a is a colimit of D. Likewise b is a colimit of D and hence $a \cong b$.

THEOREM 3. (Decomposably related implies \cong_p .) If every component of the category <u>C</u> has an initial object, if Φ is a set of constructors on <u>C</u>, and if R is a Φ -decomposable relation, then aRb implies $a \cong_p b$.

Proof. Suppose \emptyset and R are as hypothesized and suppose aRb. Let I be the set of partial isomorphisms from a to b of the form $a \cong \varphi(a_1, \dots, a_n) \leftarrow \varphi(p_1, \dots, p_n) \rightarrow \varphi(b_1, \dots, b_n) \cong b$, where φ is in $\overline{\phi}$, a_iRb_i, p_i is an initial object of the common component of a_i and b_i , $a \cong \varphi(a_1, \dots, a_n)$ and $\varphi(b_1, \dots, b_n) \cong b$ are isomorphisms, and for d_i in the component of p_i and $p_i \rightarrow d_i$ the unique arrows, $\varphi(p_1, \dots, p_n) \rightarrow \varphi(d_1, \dots, d_n)$ is $\varphi(p_1 \rightarrow d_1, \dots, p_n \rightarrow d_n)$.

If p is an initial object of the common component of a and b, then, since the identity is in $\overline{\Phi}, a \leftarrow p \rightarrow b$ is in I and hence I is nonempty. If $a \cong \varphi(a_1, \dots, a_n) \leftarrow \varphi(p_1, \dots, p_n) \rightarrow \varphi(b_1, \dots, b_n) \cong b$ is in I and $c \rightarrow a$ compact, then, since φ is a constructor, the compact arrow $c \rightarrow \varphi(a_1, \dots, a_n) = c \rightarrow a \cong \varphi(a_1, \dots, a_n)$ factors through $\varphi(c_1,$

DALE MYERS

 $\cdots, c_n) \to \varphi(a_1, \cdots, a_n) \text{ for some compacts } c_i \to a_i. \text{ Since } R \text{ is } \Phi \text{-} \text{decomposable, each } c_i \to a_i \text{ factors through } \varphi(p_{i1}, \cdots, p_{i_{p_i}}) \to \varphi_i(a_{i1}, \cdots, a_{in_i}) \cong a_i \text{ for some } \varphi_i \text{ in } \overline{\Phi}, \text{ some } a_{ij} \text{ and } b_{ij} \text{ with } a_{ij}Rb_{ij} \text{ and some } a_i \cong \varphi_i(a_{i1}, \cdots, a_{in_i}) \text{ and } \varphi_i(b_{i1}, \cdots, b_{in_i}) \cong b_i. \text{ Hence } c \to a \text{ factors through } a \cong \varphi(a_1, \cdots, a_n) \cong \varphi(\varphi_1(a_{11}, \cdots, a_{1n_1}), \cdots, \varphi_n(a_{n1}, \cdots, a_{nn_n})) \leftarrow \varphi(\varphi_1(p_{11}, \cdots, p_{1n_1}), \cdots, \varphi_n(p_{n1}, \cdots, p_{nn_n})) \to \varphi(\varphi_1(b_{11}, \cdots, b_{1n_1}), \cdots, \varphi_n(b_{n1}, \cdots, b_{nn_n})) \cong \varphi(a_1, \cdots, a_n) \cong \varphi(p_1, \cdots, p_n) \to \varphi(b_1, \cdots, b_n) \cong b \text{ via } \varphi(p_1, \cdots, p_n) \to \varphi(\varphi_1(p_{11}, \cdots, p_{1n_1}), \cdots, \varphi_n(p_{n1}, \cdots, p_{nn_n})). \text{ Hence } I: a \cong_p b.$

Suppose for the remainder of the section that \underline{C} has an initial object p. Then condition (1) of the Φ -decomposability definition is trivial.

DEFINITION. With respect to a given set Φ of constructors on \underline{C} , an object b is a factor of an object a iff $a \cong \varphi(a_1, \dots, a_n, b)$ for some a_1, \dots, a_n and some $\varphi \in \overline{\Phi}$. \underline{C} is Φ -decomposable iff every compact arrow $c \to a$ factors through $\varphi(p, \dots, p) \to \varphi(a_1, \dots, a_n) \cong a$ for some a_1, \dots, a_n , some $\varphi \in \overline{\Phi}$, and some $\varphi(a_1, \dots, a_n) \cong a$.

NOTE. <u>C</u> is Φ -decomposable iff the equality relation on <u>C</u> is Φ -decomposable.

DEFINITION. For any set Φ of constructors and any object a, the Φ -factor algebra of a is the partial algebra consisting of isomorphism types of factors of a under the operations of Φ plus a constant for the isomorphism type of a. (See the next section for examples.)

COROLLARY 4. If \underline{C} is Φ -decomposable and objects a and b have isomorphic factor algebras, then $a \cong_{p} b$.

Proof. Let R be the relation on \underline{C} such that aRb iff a and b have isomorphic Φ -factor algebras. Suppose aRb and $c \to a$ is a compact arrow. Then by decomposability of \underline{C} , $c \to a$ factors through $\varphi(p, \dots, p) \to \varphi(a_1, \dots, a_n) \cong a$ for some $\varphi \in \overline{\Phi}$ and some $\varphi(a_1, \dots, a_n) \cong a$. Let f be an isomorphism from the Φ -factor algebra of a to that of b, let \overline{a}_i be the isomorphism type of a_i , and let b_i be of the type $f(\overline{a}_i)$. Then $b \cong \varphi(b_1, \dots, b_n)$ and a_iRb_i . Hence R is Φ -decomposable and, by Theorem 3, aRb implies $a \cong_p b$.

EXAMPLES. In the following we give examples of constructors, decomposable relations, and factor algebras in the categories of sets, linear orders, Boolean algebras, bordered surfaces, and structures of a first order language. Sets. Let Set be the category of sets and 1-1 functions. The empty set is the initial object and the finite and countable sets are the compact and countable objects respectively. Let φ : Set \rightarrow Set be the functor such that $\varphi(X) = X \cup \{X\}$ for any set X and $\varphi(f) = f \cup \{\langle \text{domain } f, \text{ range } f \rangle\}$ for any function f. Then φ is a constructor. If R is a relation on sets such that XRY iff X and Y have the same finite cardinality or are both infinite, then R is $\{\varphi\}$ -decomposable. The $\{\varphi\}$ -factor algebra of the set $\{0, 1, \dots, n-1\}$ is $\langle\{0, 1, \dots, n\}, S, n\rangle$ where S(i) is i+1 if i < n and undefined if i = n. The $\{\varphi\}$ -factor algebra of $\omega = \{0, 1, \dots\}$ is (isomorphic to) $\langle \omega \cup \{\infty\}, S, \infty \rangle$ where $S(\alpha) = \alpha + 1$ if $\alpha \in \omega$ and ∞ if $\alpha = \infty$.

Linear orders. Let Lin be the category of linear orders and 1-1 order preserving functions. The empty order is the initial object and the finite and countable orders are the compact and countable objects respectively. Let φ : Lin \times Lin \rightarrow Lin be a functor such that $\varphi(K, L)$ is an order consisting of an initial segment of type K followed by a new point followed by a final segment of type L and for functions f and g, $\varphi(f, g)$ is, essentially, f on the first segment, the new point goes to the new point, and g on the final segment. Then φ is a constructor. If R is the relation such that KRL iff K and L are both dense linear orders without end points, then R is $\{\varphi\}$ decomposable. If R is any symmetric relation on linear orders such that (1) KRL implies K is empty iff L is and (2) KRL and $a \in k$ implies there is a $b \in L$ such that $\langle \{x \in K : x <_{\kappa} a\}, \langle_{\kappa} \rangle R \langle \{x \in L : x <_{\iota} b\},$ $<_L$ and $\langle \{x \in K : x >_K a\}, <_K \rangle R \langle \{x \in L : x >_L b\}, <_L \rangle$, then R is $\{\varphi\}$ decomposable. The $\{\varphi\}$ -factor algebra of the empty order is $\{\{1\}, +, \}$ 1) where 1 + 1 is undefined. The $\{\varphi\}$ -factor algebra of the rationals with the usual order is $\langle \{\infty\}, +, \infty \rangle$ where $\infty + \infty = \infty$.

Boolean algebras. Let **Bool** be the category of Boolean algebras with two or more elements and monomorphisms. The two-element Boolean algebra is the initial object and the finite and countable algebras are the compact and countable objects respectively. Let φ : **Bool** \times **Bool** \rightarrow **Bool** be the cartesian product functor. Then φ is a constructor. If R is the relation such that ARB iff A and B are atomless Boolean algebras, then R is $\{\varphi\}$ -decomposable. If R is any symmetric relation on Boolean algebras such that (1) ARB implies Ais the two-element algebra iff B is and (2) ARB and $a \in A$ implies there is a $b \in B$ such that A[a]RB[b] and A[-a]RB[-b] where A[a]is the ideal (considered as a Boolean algebra) generated by a, then R is $\{\varphi\}$ -decomposable. Relations of this type were first studied by Vaught [13] who showed that any two countable Boolean algebras related by such a relation were isomorphic. Decomposability is

DALE MYERS

simply a categorical generalization of conditions (1) and (2). The $\{\mathcal{P}\}$ -factor algebras of the two-element algebra and the atomless algebra are the partial algebras $\langle \{1\}, +, 1 \rangle$ and $\langle \{\infty\}, +, \infty \rangle$ respectively defined in the previous example. For primitive Boolean algebras, $\{\mathcal{P}\}$ -factor algebras are nothing more than algebraic versions of the structure diagrams of Hanf [3] and Pierce [10] which have played an important role in classifying such Boolean algebras and determining isomorphism types of Lindenbaum-Tarski algebras [3, 9, 12].

Bordered surfaces. Let Sur be the category whose objects are bordered orientable surfaces (orientable two-dimensional manifolds whose boundary is a disjoint union of simple closed curves) each with a designated homeomorphism from the closed unit disc onto a subspace of the surface's interior and whose arrows are homeomorphisms from one surface onto a subsurface of a second such that the designated map of the first composed with the arrow is the designated map of the second and such that a boundary curve of the first is either carried onto a boundary curve of the second or into the interior of the second. The closed disc with the identity map as the designated map is the initial object, the compact bordered surfaces are the compact objects (this is false if the boundary curve condition on arrows is dropped), and the separable bordered surfaces are the countable objects. Let C be a cylinder — finite length, closed, and open at both ends — with a designated (map from the) disc in its interior. Let φ : Sur \times Sur \rightarrow Sur be a functor such that for any objects S and T, $\varphi(S, T)$ is a surface obtained by cutting out the interiors of the designated disc of S and T and gluing their perimeters to the open ends of C (C becomes a tube joining the two surfaces) and letting the designated disc of C be the designated disc of the result; and such that for any arrows f and g, $\varphi(f, g)$ is, essentially, f on the first surface, g on the second, and the identity on C. Let σ and τ be the 0-ary functors whose unique values are the sphere and torus respectively with designated discs. Then φ , σ , and τ are constructors. If R is the relation between surfaces such that SRT iff S and T have the same number of boundary components, the same genus, and homeomorphic ideal boundaries (see [11]), then R is $\{\varphi, \sigma, \tau\}$ -decomposable. The $\{\varphi, \sigma, \tau\}$ factor algebra of the surface of a solid semi-infinite rod is $\langle \{s, t, 1\}, +, \rangle$ s, t, 1 where s + s = s, s + t = t + s = t, s + 1 = 1 + s = 1, and all other sums are undefined. The $\{\varphi, \sigma, \tau\}$ -factor algebra of the surface of an infinite binary tree with solid branches of nonzero width is $\langle \{s, t, \infty\}, +, s, t, \infty \rangle$ where $s + s = s, s + t = t + s = t, s + \infty = \infty + s = s = t$ ∞ , $\infty + \infty = \infty$, and all other sums are undefined.

Structures. Given a first-order language with at least one constant

symbol, let Str be the category whose objects are finite expansions of structures of the language via constants, i.e., structures of the form $(\mathfrak{A}, a_1, \cdots, a_n)$ where \mathfrak{A} is a structure of the language and a_1, \cdots, a_n $a_n \in \mathfrak{A}$, and whose arrows are isomorphisms from one structure to a substructure of a second, i.e., monomorphisms which preserve quantifier-free formulas. For any object B of Str the initial object of its component is Prime (\mathfrak{B}) , the substructure of \mathfrak{B} generated by its Finitely generated structures and countably generated constants. structures are the compact and countable objects respectively. Let $\varphi: \mathbf{Str} \to \mathbf{Str}$ be the functor such that for any structure \mathfrak{A} of the language $\varphi(\mathfrak{A}) = \mathfrak{A}$ and for any $a_1, \dots, a_n \in \mathfrak{A}, \varphi((\mathfrak{A}, a_1, \dots, a_n)) = (\mathfrak{A}, \mathfrak{A})$ a_1, \dots, a_{n-1}). Then φ is a constructor. If R is the relation such that \mathfrak{ARB} iff \mathfrak{A} and \mathfrak{B} are elementarily equivalent ω -saturated structures, then R is $\{\varphi\}$ -decomposable. If R is a symmetric relation such that (1) \mathfrak{ARB} implies Prime $\mathfrak{A} \cong$ Prime \mathfrak{B} and (2) \mathfrak{ARB} and $a \in \mathfrak{A}$ implies there is a $b \in \mathfrak{B}$ such that $(\mathfrak{A}, a)R(\mathfrak{B}, b)$, then R is $\{\varphi\}$ -decomposable.

References

1. Jon Barwise, *Back and forth through infinitary logic*, Studies in Model Theory (M. D. Morley, Editor), The Mathematical Association of America, 1973.

 Georg Cantor, Beiträge zur Begründung der transfiniten Mengenlehre, Math. Ann., 96 (1895), 481-512.

 William Hanf, Primitive Boolean algebras, Proceedings of the Symposium in Honor of Alfred Tarski, (Berkeley, 1971), Amer. Math. Soc., Providence, R. I., 25 (1974), 75-90.
 Henkin, Monk, and Tarski, Cylindric Algebras, North-Holland Publishing Company, Amsterdam, 1971.

5. Irving Kaplansky, Infinite Abelian Groups, Michigan Press, Ann Arbor, 1969.

6. Carol Karp, Finite Quantifier Equivalence, The Theory of Models, North-Holland Publishing Company, Amsterdam, 1965.

7. Saunders Mac Lane, Categories for the Working Mathematician, Springer-Verlag, New York, 1971.

8. M. Morley and R. L. Vaught, *Homogeneous universal models*, Math. Scand., **11** (1962), 37-57.

9. Dale Myers, The Boolean algebras of abelian groups and well-orders, J. Symbolic Logic, **39** (1974), 452-458.

10. R. S. Pierce, Compact zero-dimensional metric spaces of finite type, Memoirs of the Amer. Math Soc., no. 130 (1972).

11. Ian Richards, On the classification of noncompact surfaces. Trans. Amer. Math. Soc., 106 (1963), 259-269.

12. Roger Simons, The Boolean algebra of sentences of the theory of a function, Ph. D. thesis, Berkeley, 1972.

13. R. L. Vaught, Topics in the theory of arithmetical classes and Boolean algebras, Ph. D. thesis, Berkeley, 1954.

Received June 11, 1974.

UNIVERSITY OF HAWAII

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor) University of California Los Angeles, California 90024 J. DUGUNDJI

Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

ASSOCIATE EDITORS

E.F. BECKENBACH

R. A. BEAUMONT

University of Washington

Seattle, Washington 98105

B. H. NEUMANN

F. WOLF K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

AMERICAN MATHEMATICAL SOCIETY NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

> Copyright © 1973 by Pacific Journal of Mathematics Manufactured and first issued in Japan

Pacific Journal of Mathematics Vol. 55, No. 2 October, 1974

Walter Allegretto, On the equivalence of two types of oscillation for elliptic operators	319
Edward Arthur Bertram, A density theorem on the number of conjugacy classes in	
finite groups	329
Arne Brøndsted, On a lemma of Bishop and Phelps	335
Jacob Burbea, Total positivity and reproducing kernels	343
Ed Dubinsky, Linear Pincherle sequences	361
Benny Dan Evans, Cyclic amalgamations of residually finite groups	371
Barry J. Gardner and Patrick Noble Stewart, A going down theorem for certain	201
In the Light Cross and Themas William Tusker, Quatients of semplate engrha	301
Jonathan Light Gross and Thomas witham Tucker, Quotients of complete graphs:	201
Seu Demen Henremin Course of matrices active on distribution and	391
Sav Roman Harasymiv, Groups of matrices acting on distribution spaces	403
linearly ordered spaces	/10
Chung Wu Ho. Deforming p. 1. homeomorphisms on a convex polygonal	419
2-disk	427
Richard Farl Hodel Metrizability of topological spaces	441
Wilfried Imrich and Mark F. Watkins. On araphical regular representations of	771
cvclic extensions of groups	461
Jozef Krasinkiewicz. Remark on mappings not raising dimension of curves	479
Melven Robert Krom, Infinite games and special Baire space extensions	483
S. Leela. Stability of measure differential equations	489
M. H. Lim, Linear transformations on symmetric spaces	499
Teng-Sun Liu Arnoud C M van Rooij and Ju-Kwei Wang <i>On</i> some group algebra	
modules related to Wiener's algebra M_1	507
Dale Wayne Myers. The back-and-forth isomorphism construction	521
Donovan Harold Van Osdol. Extensions of sheaves of commutative algebras by	
nontrivial kernels	531
Alan Rahilly, Generalized Hall planes of even order	543
Joylyn Newberry Reed. On completeness and semicompleteness of first countable	
spaces	553
Alan Schwartz, Generalized convolutions and positive definite functions associated	
with general orthogonal series	565
Thomas Jerome Scott, <i>Monotonic permutations of chains</i>	583
Eivind Stensholt, An application of Steinberg's construction of twisted groups	595
Yasuji Takeuchi, On strongly radicial extensions	619
William P. Ziemer, Some remarks on harmonic measure in space	629
John Grant, Corrections to: "Automorphisms definable by formulas"	639
Peter Michael Rosenthal, Corrections to: "On an inversion for the general	
Mehler-Fock transform pair"	640
Carl Clifton Faith, Corrections to: "When are proper cyclics injective"	640