# Pacific Journal of Mathematics

# ABELIAN GROUPS, A, SUCH THAT Hom(A, --)PRESERVES DIRECT SUMS OF COPIES OF A

DAVID MARION ARNOLD AND CHARLES ESTEP MURLEY

Vol. 56, No. 1

November 1975

## ABELIAN GROUPS, A, SUCH THAT HOM (A,-)PRESERVES DIRECT SUMS OF COPIES OF A

D. M. ARNOLD AND C. E. MURLEY

An *R*-module, *A*, is *self-small* if Hom(A,-) preserves direct sums of copies of *A*. Various conditions on the endomorphism ring of a module which guarantee that it is self-small are studied. Various results are proved about subgroups of direct sums or direct products of copies of a self-small abelian group *A*, which generalize results previously known when *A* is torsion free of rank one.

**0.** Introduction. An *R*-module, *A*, is *self-small* if  $\text{Hom}_R(A, -)$  preserves direct sums of copies of *A*. Homological arguments show that if *A* is a self-small *R*-module with *R* a commutative ring with 1, then the category of direct summands of direct sums of copies of *A* is equivalent to the category of projective right  $\text{End}_R(A)$ -modules  $(\text{End}_R(A))$  is the *R*-endomorphism ring of *A*). Consequently, direct sum decompositions of direct sums of copies of *A* may be interpreted in terms of direct sum decompositions of free  $\text{End}_R(A)$ -modules.

An *R*-module, *A*, is self-small in the following cases: (a) *A* is *small* (i.e., Hom<sub>*R*</sub>(*A*,-) preserves arbitrary direct sums of *R*-modules); (b)  $A = \prod_{i \in I} A_i$ , where each  $A_i$  is a self-small *R*-module and Hom<sub>*R*</sub>( $A_i, A_j$ ) = 0 if  $i \neq j$ ; (c) End<sub>*R*</sub>(*A*) is countable.

If the finite topology on  $\operatorname{End}_R(A)$  is discrete, then A is selfsmall. In certain cases, the converse is true.

COROLLARY I. Suppose that A is a countably generated Rmodule. Then A is self-small iff the finite topology on  $End_{R}(A)$  is discrete. If R is countable, then A is self-small iff  $End_{R}(A)$  is countable.

A left ideal, I, of  $\operatorname{End}_R(A)$  is an annihilator ideal if  $I = \{f \in \operatorname{End}_R(A): f(x) = 0 \text{ for all } x \in A \text{ with } Ix = 0\}.$ 

PROPOSITION II. Suppose that A is an R-module and that  $End_R(A)$  has the minimum condition on left annihilator ideals. Then the finite topology on  $End_R(A)$  is discrete and A is the finite direct sum of indecomposable R-modules.

The remainder of the paper is devoted to self-small abelian groups (although many of the arguments are valid in a more general setting). Self-small torsion abelian groups are finite. Section 3 and examples in §5 demonstrate that self-small torsion free abelian groups are both profuse and diverse. Self-small mixed abelian groups with finite torsion free rank are characterized by Proposition 3.6. Generalizations of homogeneous separable and completely decomposable torsion free abelian groups are considered in §4. We demonstrate that many of the classical properties of these groups may be viewed as consequences of the fact that endomorphism rings of rank 1 torsion free abelian groups are principal ideal domains.

Let G and A be abelian groups. Define G to be A-free if G is isomorphic to a direct sum of copies of A; A-projective if G is a summand of an A-free group; and locally A-projective [locally A-free] if every finite subset of G is contained in an A-projective [A-free] summand of G. An R-module, M, is locally projective [locally free] if every finite subset of M is contained in an R-projective [R-free] summand of M. Note that if A is torsion free of rank 1, then the class of locally A-projective [A-projective] groups coincides with the class of homogeneous separable [completely decomposable] groups (with type = type of A). Define  $S_A(G)$  to be the subgroup of G generated by  $\{f(A)|f \in Hom(A, G)\}$ .

THEOREM III. Suppose that A is an abelian group and that End(A) is discrete in the finite topology. Then the category of locally A-projective abelian groups is equivalent to the category of locally projective right End(A)-modules.

Results of Chase [3] and Theorem III suffice to prove:

COROLLARY IV. Let A be a torsion free abelian group such that End(A) is a principal ideal domain and A/im f is torsion for all  $0 \neq f \in End(A)$ .

(a) If B is a pure subgroup of an A-free group, G, such that  $S_A(B) = B$  and if End(B) is discrete in the finite topology, then B is an A-free summand of G.

(b) A group, G, is locally A-free iff  $S_A(G) = G$  and G is isomorphic to a pure subgroup of  $S_A(\prod_{i \in I} A_i)$ , where  $A_i \cong A$  for all  $i \in I$ .

(c) If B is a pure subgroup of a locally A-free group, G, and if  $S_A(B) = B$ , then B is locally A-free. Moreover, if End(B) is discrete in the finite topology, then B is an A-free summand of G.

(d) Countable locally A-free groups are A-free.

Note that if A is torsion free of rank 1 and type  $\tau$ , then End(A) is a principal ideal domain and  $S_A(G) = G(\tau)$ . Thus Corollary IV includes the classical properties of homogeneous separable groups as a special case (see Fuchs [7]).

Fundamental references, for this paper, are Fuchs [6] and [7].

**1. Self-small modules.** Let  $\sum_{i \in I} \bigoplus A_i$  be a direct sum of copies of an *R*-module *A*. There is a natural monomorphism  $e_I: \sum_{i \in I} \bigoplus \operatorname{Hom}_R(A, A_i) \to \operatorname{Hom}_R(A, \sum_{i \in I} \bigoplus A_i)$  induced by projection maps. Consequently, *A* is self-small iff for every countable index set *I* 

and every *R*-homomorphism  $\phi: A \to \sum_{i \in I} \bigoplus A_i$ , there is a finite subset, *I'*, of *I* with  $\phi(A) \subseteq \sum_{i \in I'} \bigoplus A_i$ . Every finitely generated *R*-module is self-small. On the other hand, if *A* is an infinite direct sum of *R*-modules, then *A* is not self-small.

Observe that endomorphic images (in particular, direct summands) of self-small *R*-modules are self-small.

Let X and Y be nonempty subsets of A and  $\operatorname{End}_R(A)$ , respectively. Define  $X^* = \{f \in \operatorname{End}_R(A) | f(X) = 0\}$  and  $Y^* = \{x \in A | f(x) = 0 \text{ for all } f \in Y\}$ . A left ideal, I, of  $\operatorname{End}_R(A)$  is an annihilator ideal if  $I = I^{**}$ . An R-submodule, B, of A is a kernel submodule if  $B = B^{**}$ . One can easily verify that  $X^*$  is a left ideal of  $\operatorname{End}_R(A)$ ;  $Y^* = \bigcap_{f \in Y} \ker f$  is a submodule of A;  $X \subseteq X^{**}$ ;  $Y \subseteq Y^{**}$ ; a left ideal, I, of  $\operatorname{End}_R(A)$  is an annihilator ideal iff  $I = X^*$  for some  $X \subseteq A$ ; and a submodule, B, of A is a kernel submodule iff  $B = Y^*$  for some  $Y \subseteq \operatorname{End}_R(A)$ .

LEMMA 1.1. There is a 1-1 order inverting correspondence between the kernel submodules of an R-module, A, and the left annihilator ideals of  $End_R(A)$ .

*Proof.* The correspondence is given by  $B \rightarrow B^*$  and  $I \rightarrow I^*$ .

**PROPOSITION 1.1.** The following statements are equivalent for an R-module, A:

(a) A is not self-small;

(b) there is a chain  $A_1 \subseteq \cdots \subseteq A_n \subseteq \cdots$  of (proper) submodules of A such that  $A = \bigcup_{n=1}^{\infty} A_n$  and  $A_n^* \neq 0$  for all n;

(c) there is a chain  $A_1 \subseteq \cdots \subseteq A_n \subseteq \cdots$  of proper kernel submodules of A such that  $A = \bigcup_{n=1}^{\infty} A_n$ ;

(d) there is a chain  $I_1 \supseteq \cdots \supseteq I_n \supseteq \cdots$  of nonzero left annihilator ideals of  $\operatorname{End}_R(A)$  such that  $A = \bigcup_{n=1}^{\infty} I_n^*$ . In this case,  $\bigcap_{n=1}^{\infty} I_n = 0$ ;

(e) there is an infinite subset, S, of  $\operatorname{End}_{\mathbb{R}}(A)$  such that  $S \setminus (X^* \cap S)$  is finite for all finite subsets, X, of A.

**Proof.** (a)  $\Rightarrow$  (b) Since A is not self-small, there is an R-homomorphism  $\phi: A \to \sum_{i=1}^{\infty} \bigoplus B_i$  such that  $\prod_i \phi \neq 0$  for all *i*, where  $\prod_i : \Sigma \bigoplus B_i \to B_j = A$  is the projection map for all *j*. For  $n \ge 1$  let  $A_n = \{x \in A \mid \prod_i \phi(x) = 0 \text{ for } i > n\}$  so that  $A_1 \subseteq \cdots \subseteq A_n \subseteq \cdots$  is a chain of proper submodules of A with  $A = \bigcup_{n=1}^{\infty} A_n$ . Clearly,  $A_n^* \neq 0$  for all n.

(b)  $\Rightarrow$  (c) Replace each  $A_n$  by  $A_n^{**}$ , thereby obtaining an ascending chain of kernel submodules with  $A = \bigcup_{n=1}^{\infty} (A_n^{**})$  (since  $A_n \subseteq A_n^{**}$ ). Each  $A_n^{**}$  is proper, for if  $A_n^{**} = A$ , then  $A_n^* = A^* = 0$ , an impossibility by (b).

(c)  $\Rightarrow$  (d) A consequence of Lemma 1.1.

(d)  $\Rightarrow$  (e) For each *n*, choose  $f_n \in I_n \setminus I_{n+1}$  if  $I_n \neq I_{n+1}$  and let  $f_n = 0$  if  $I_n = I_{n+1}$ . Then  $S = \{f_n | n = 1, 2, \dots\}$  is an infinite subset of End<sub>R</sub>(A)

(since  $\bigcap_{n=1}^{\infty} I_n = 0$  and each  $I_n$  is nonzero). If X is a finite subset of A, then  $X \subseteq I_n^*$  for some n (since  $A = \bigcup_{n=1}^{\infty} I_n^*$ ). Consequently,  $I_m = I_m^{**} \subseteq X^*$  for all  $m \ge n$  so that  $S \setminus (X^* \cap S) \subseteq \{f_1, \dots, f_{n-1}\}$ , a finite set.

(e)  $\Rightarrow$  (a) Define an *R*-homomorphism  $\phi: A \to \Sigma_{f \in S} \bigoplus A_f$  by  $\phi(a) = \Sigma_{f \in S} f(a)$  where  $f(a) \in A_f = A$ . The hypotheses guarantee that  $\phi$  is well defined. Since *S* is infinite,  $\phi(A) \not\subseteq \Sigma_{f \in S'} \bigoplus A_f$  for all finite subsets, *S'*, of *S*. Thus *A* is not self-small.

**REMARK.** In the language of Szele [13], (e) may be restated as: End<sub>R</sub>(A) has an infinite 0-system (equivalently, End<sub>R</sub>(A) has an infinite summable system.

COROLLARY 1.3. Let  $\{B_i\}_{i \in I}$  be a family of self-small R-modules with  $\operatorname{Hom}_{\mathbb{R}}(B_i, B_j) = 0$  for  $i \neq j$ . Then  $A = \prod_{i \in I} B_i$  is self-small.

**Proof.** Assume that A is not self-small and choose a chain  $A_1 \subseteq \cdots \subseteq A_n \subseteq \cdots$  of proper kernel submodules of A with  $A = \bigcup_{n=1}^{\infty} A_n$ . For each  $i \in I$ , there is a least integer n(i) with  $B_i \subseteq A_{n(i)}$  (otherwise,  $B_i$  is not self-small by Proposition 1.2.b). If  $\{n(i)|i \in I\}$  is bounded, say by m, then for any  $0 \neq f \in A_m^*$   $(A_m^* \neq 0$  since  $A_m = A_m^{**} \neq A$ ) it follows that f(A) = 0, a contradiction. If  $\{n(i)|i \in I\}$  is not bounded, then there is some  $a = (b_i) \in A$  with  $a \notin A_m$  for all m, a contradiction (since  $A = \bigcup_{n=1}^{\infty} A_n$ ).

COROLLARY 1.4. If A is an R-module such that  $End_R(A)$  is countable, then A is self-small.

**Proof.** Assume that A is not self-small. By Proposition 1.2.d there is a chain  $I_1 \supset \cdots \supset I_n \supset \cdots$  of nonzero left annihilator ideals of  $\operatorname{End}_R(A)$  such that  $A = \bigcup_{n=1}^{\infty} I_n^*$  (choose an appropriate subchain to guarantee that  $I_n \neq I_{n+1}$  for all n). For each n, choose  $g_n \in I_n \setminus I_{n+1}$ ; let  $S = \{g_n\}$ ; and for each subset, L, of  $Z^+$  (the positive integers) define  $g_n^L = g_n$  if  $n \in L$  and  $g_n^L = 0$  otherwise. Define  $g^L \in \operatorname{End}_R(A)$  by  $g^L(a) = \Sigma g_i^L(a)$ , a well defined homomorphism (since for each  $a \in A$ ,  $g_i^L(a) = 0$  for almost all i).

We prove that  $g^{L} = g^{K}$  iff K = L; in which case,  $\operatorname{End}_{R}(A)$  is uncountable, a contradiction. Assume that  $K \neq L$  and choose a least integer *n* with  $g_{n}^{L} \neq g_{n}^{K}$ , say  $g_{n}^{L} = g_{n}$  and  $g_{n}^{K} = 0$ . Since  $g_{n} \notin I_{n+1}$ , there is some  $a \in I_{n+1}^{*}$  such that  $a \notin \ker g_{n}$ . Thus  $g^{L}(a) - g^{K}(a) = g_{n}(a) \neq 0$ and  $g^{K} \neq g^{L}$ .

2. The Finite Topology on  $\operatorname{End}_R(A)$ . The finite topology on  $\operatorname{End}_R(A)$  is defined by letting  $\{X^*|X \text{ is a finite subset of } A\}$  be a basis of open neighborhoods of 0. It is known that  $\operatorname{End}_R(A)$  is a complete Hausdorff topological ring in the finite topology (see Fuchs [7], p. 221, for the case that R = Z; the general argument is similar). The finite topology on  $\operatorname{End}_{R}(A)$  is discrete (i.e.,  $X^{*} = 0$  for some finite subset of A) iff  $B^{*} = 0$  for some finitely generated sub-module, B, of A.

COROLLARY 2.1. If the finite topology on  $\operatorname{End}_{\mathbb{R}}(A)$  is discrete, then A is self-small.

*Proof.* If A is not self-small, then Proposition 1.2.b leads to a contradiction.

**PROPOSITION 2.2.** Suppose that A is an R-module such that the finite topology on  $\operatorname{End}_{R}(A)$  is either first countable or locally compact. Then A is self-small iff the finite topology on  $\operatorname{End}_{R}(A)$  is discrete.

*Proof.* ( $\Leftarrow$ ) Corollary 2.1.

(⇒) Assume that  $\operatorname{End}_R(A)$  is first countable. Then there is a set  $\{X_n | n = 1, 2, \cdots\}$  of finite subsets of A such that  $X_1^* \supseteq X_2^* \supseteq \cdots$ ;  $\bigcap_{n=1}^{\infty} X_n^* = 0$ ; and if V is a neighborhood of 0, then  $X_n^* \subseteq V$  for some n. If  $x \in A$ , then  $X_n^* \subseteq \{x\}^*$  for some n, so that  $x \in X_n^{**}$ . Consequently,  $A = \bigcup_{n=1}^{\infty} (X_n^*)^*$ . By Proposition 1.2.d. and the preceding remarks,  $X_n^* = 0$  for some n, i.e.,  $\operatorname{End}_R(A)$  is discrete.

Assume that  $\operatorname{End}_R(A)$  is locally compact. Then there is a compact ideal neighborhood, *I*, of 0. Since  $\operatorname{End}_R(A)$  is Hausdorff, *I* is both closed and complete. But  $I \setminus (X^* \cap I)$  is discrete (since  $X^*$  is open) and compact, thus finite, for all finite subsets, *X*, of *A*. Since *A* is self-small, *I* must be finite (Proposition 1.2.e.). But *I* is a neighborhood of 0 and  $\operatorname{End}_R(A)$  is Hausdorff, so  $X^* = 0$  for some finite subset, *X*, of *A*.

COROLLARY 2.3. Suppose that A is a countably generated R-module.

- (a) A is self-small iff the finite topology on  $End_R(A)$  is discrete.
- (b) If R is countable, then A is self-small iff  $End_R(A)$  is countable.

**Proof.** (a) Let S be a countable set of generators for A. Then  $\{X^*|X \text{ is a finite subset of } S\}$  is a countable neighborhood basis of 0; i.e., End<sub>R</sub>(A) is first countable. Now apply Proposition 2.2.

(b) In view of Corollary 1.4, it suffices to prove that if R is countable and A is self-small, then  $\operatorname{End}_R(A)$  is countable. By (a),  $\operatorname{End}_R(A)$  is discrete, i.e.,  $X^* = 0$  for some finite subset  $X = \{x_1, \dots, x_n\}$  of A. Define  $\phi: \operatorname{End}_R(A) \to \sum_{i=1}^n \bigoplus A_i$  by  $\phi(f) = f(x_1) + \dots + f(x_n)$ , where  $f(x_i) \in A_i = A$ . Then  $\phi$  is a monomorphism and  $\operatorname{End}_R(A)$  is countable (since A is countable).

**PROPOSITION 2.4.** Assume that  $\text{End}_{R}(A)$  has the minimum condition on left annihilator ideals.

(a) The finite topology on  $End_{R}(A)$  is discrete.

(b) If B is a summand of A, then  $End_{R}(B)$  has the minimum condition on left annihilator ideals.

(c) A is a finite direct sum of indecomposable R-modules.

**Proof.** (a) Let  $S = \{X^* | X \text{ is a finite nonzero subset of } A \}$  be a collection of left annihilator ideals of  $\operatorname{End}_R(A)$ . If  $X^* = 0$  for some X, then  $\operatorname{End}_R(A)$  is discrete. Otherwise, S has a minimal nonzero element, say  $Y^*$ . If X is a finite subset of A, then  $(Y \cup X)^* \subseteq Y^*$ , so by the minimality of  $Y^*$ ,  $Y^* \cap X^* = (Y \cup X)^* = Y^*$ . Thus  $Y^* \subseteq \cap \{X^* | X^* \in S\} = 0$ , a contradiction.

(b) Write  $A = B \bigoplus C$  and identify  $\operatorname{End}_R(B)$ , in the usual way, with a subring of  $\operatorname{End}_R(A)$  so that  $\operatorname{End}_R(B) \subseteq C^* \subseteq \operatorname{End}_R(A)$ . For *I*, a left annihilator ideal of  $\operatorname{End}_R(B)$ , let  $e(I) = (I^* \bigoplus C)^* \subseteq$  $\operatorname{End}_R(A)$ . Note that  $e(\operatorname{End}_R(B)) = C^*$  and that *e* is an order preserving map from the annihilator ideals of  $\operatorname{End}_R(B)$  to the annihilator ideals of  $\operatorname{End}_R(A)$ . Therefore, it is enough to show that *e* is monic. Let *J* be an annihilator ideal of  $\operatorname{End}_R(B)$  and let  $\phi \in I \setminus J$ . Then  $\phi \in e(I)$ , but  $\phi(J^* \oplus C) = \phi(J^*) \neq 0$ , i.e.,  $\phi \notin e(J)$ . Consequently, e(I) = e(J)iff I = J.

(c) First of all, A must have a nonzero indecomposable summand. Otherwise, there is a chain  $A_1 \subset \cdots \subset A_n \subset \cdots$  of submodules of A such that each  $A_n$  is a proper summand of A and  $A_{n+1}$ . Thus each  $A_n^*$  is a nonzero annihilator ideal and  $\{A_n^* | n = 1, 2, \cdots\}$  has no minimal element, a contradiction.

Write  $A = A_1 \bigoplus B$ , where  $A_1$  is a nonzero indecomposable summand of A. Using (b) and the fact that A is self-small, one sees that (c) is true.

### 3. Self-small Abelian groups.

**PROPOSITION 3.1.** Every self-small torsion group is finite.

**Proof.** Let A be a self-small torsion group and for each integer n regard n! as an element of End(A) (i.e., multiplication by n factorial). Define  $A_n = \{n!\}^* \subseteq A$  so that  $0 = A_1 \subseteq A_2 \subseteq \cdots \subseteq A_n \subseteq \cdots$ is chain of subgroups of A. Each  $A_n$  is bounded, hence a direct sum of cyclic groups. Consequently,  $A_n = A$  for some n and A is finite. Otherwise, each  $A_n$  is proper,  $A_n^* \neq 0$ , and A is not self-small (Proposition 1.2.b), a contradiction.

If A is a torsion free abelian group such that End(A) is countable, then A is self-small (Corollary 1.4). Examples of such groups include all torsion free abelian groups of finite rank, all of the groups constructed by Corner [4], and all rigid groups (see Fuchs [7], p. 124). Other examples, not requiring the countability of End(A), are given by: PROPOSITION 3.2. Let A be a reduced torsion free abelian group. (a) If End(A) is a Dedekind domain, then every nonzero endomorphism of A is monic.

(b) If every nonzero endomorphism of A is monic, then End(A) has the minimum condition on annihilator ideals.

(c) If B is a torsion free abelian group and if  $B \bigotimes_z A$  is self-small, then A is self-small.

**Proof.** (a) Let R = End(A) and regard A as a left R-module. Since R has no nontrivial idempotents, A is indecomposable as an abelian group and consequently as an R-module. Suppose that A is not a torsion free R-module. Since R is Dedekind and A is reduced, then A is R-isomorphic to R/I, where I is a nonzero ideal of R (see Kaplansky [8]). There is some  $0 \neq r \in I$  so that  $Rr \subseteq I$  and r(A) = 0, a contradiction. Thus A is a torsion free R-module and (a) is proved.

(c) Assume that A is not self-small. By Proposition 1.2.c. there is a chain  $A_1 \subseteq \cdots \subseteq A_n \subseteq \cdots$  of proper kernel subgroups of A there is a chain  $A_1 \subseteq \cdots \subseteq A_n \subseteq \cdots$  of proper kernel subgroups of A such that  $A = \bigcup_{n=1}^{\infty} A_n$ . Now kernel subgroups are pure  $(A_n = A_n^{**} = \bigcap_{f \in A_n^*} (\ker f))$  and B is torsion free so each  $B \bigotimes_Z A_n$  is proper in  $B \bigotimes_Z A$ ;  $B \bigotimes_Z A_1 \subseteq \cdots \subseteq B \bigotimes_Z A_n \subseteq \cdots$ ;  $\bigcup_{n=1}^{\infty} (B \bigotimes_Z A_n) = B \bigotimes_Z A$  and  $(B \bigotimes_Z A_n)^* \neq 0$  for all n (choose  $f_n \in \text{End}(A)$  with  $f_n(A_n) = 0$  and note that  $(1 \bigotimes f_n) (B \bigotimes A_n) = 0$ ). By Proposition 1.2.b.,  $B \bigotimes_Z A$  is not selfsmall, a contradiction.

For a prime, p, let  $Z_p$  be the localization of Z at a prime p (i.e., the subring of Q consisting of elements with denominator prime to p).

COROLLARY 3.3. Let A be a reduced torsion free abelian group. (a) If  $Z_p \bigotimes_Z A$  is a self-small  $Z_p$ -module for some prime, p, then A is self-small.

(b) If there is a prime p such that the cardinality of A/pA is finite and if  $\bigcap_{n=1}^{\infty} p^n A = 0$ , then A is self-small.

*Proof.* (a) is a consequence of Proposition 3.2.

(b) Choose  $X = \{x_1, \dots, x_n\}$ , where  $\{x_1 + pA, \dots, x_n + pA\}$  is a basis for A/pA (as a Z/pZ vector space). Since  $\bigcap_{n=1}^{\infty} p^n A = 0$ , one can easily see that  $X^* = 0$ , i.e., End(A) is discrete.

**REMARK.** Examples of Corollary 3.3.b include p-pure subgroups of finite direct sums of copies of the p-adic integers.

Conditions guaranteeing that a torsion free abelian group A is self-small may be recast in terms of the *quasi-endomorphism ring of A*, denoted by  $\mathscr{C}(A)$ . Walker [14] observed that  $\mathscr{C}(A)$  may be regarded as  $Q \otimes_Z \operatorname{End}(A)$  (also see Reid [11]). If  $f \in \mathscr{C}(A)$ , then  $f = 1/n \otimes g$  for some  $g \in \operatorname{End}(A)$  and  $n \in Z$ . Define ker  $f = \{x \in A | g(x) = 0\}$ ; if  $X \subseteq$ A, let  $X_* = \{f \in \mathscr{C}(A) | X \subseteq \ker f\}$ , a left ideal in  $\mathscr{C}(A)$ ; and if J is a left ideal of  $\mathscr{C}(A)$ , let  $J_* = \bigcap_{f \in J} \ker f$ . Call J an annihilator ideal of  $\mathscr{C}(A)$  if  $J = J_{**}$ .

**PROPOSITION 3.4.** If A is a torsion free abelian group, then there is a 1-1 order preserving correspondence between left annihilator ideals in End(A) and left annihilator ideals in  $\mathscr{E}(A)$ .

**Proof.** The correspondence is given by  $I \rightarrow QI$  and  $J \rightarrow J \cap$ End(A), where End(A) is canonically embedded in  $\mathscr{C}(A)$  (via  $f \rightarrow 1 \otimes f$ ). Note that left annihilator ideals of End(A) are pure subgroups of Hom(A, A). The remainder of the proof is straightforward and is left to the reader.

COROLLARY 3.5. Let A be a torsion free abelian group.

(a) End(A) has minimum condition on left annihilator ideals iff  $\mathscr{E}(A)$  has minimum condition on left annihilator ideals.

(b) End(A) is discrete iff  $\mathscr{E}(A)$  is discrete.

(c) A is not self-small iff there is a chain  $J_1 \supset \cdots \supset J_n \supset \cdots$  of nonzero annihilator ideals of  $\mathscr{C}(A)$  with  $A = \bigcup_{n=1}^{\infty} (J_n)_*$ .

REMARK. If  $\mathscr{E}(A)$  has the minimum condition on left ideals, then A is self-small. This class of groups has been considered by Reid [11] and others.

The situation is even more complicated for mixed abelian groups, A. Let tA be the torsion subgroup of A and, for p a prime, let  $(tA)_p$  be the p-component of tA. If A is a torsion free abelian group of finite rank, then the R-type of A is the quasi-isomorphism class of A/F, where F is a free subgroup of A with A/F torsion (e.g., see Richman [12]).

**PROPOSITION 3.6.** Suppose that A is a mixed abelian group and that A/tA has finite rank. Then A is self-small iff (a) for all primes, p,  $(tA)_p$  is finite or zero; and (b) the R-type of A/tA is p-divisible for all primes p with  $(tA)_p \neq 0$ .

*Proof.* ( $\Rightarrow$ ) Let *p* be a prime with  $(tA)_p \neq 0$ . Since *A* is self-small and A/pA is a direct sum of cyclic groups of order *p*, it follows that A/pA is finite (otherwise map each summand of A/pA into  $(tA)_p \subset A$ ). But  $(tA)_p/p(tA)_p$  is a summand of A/pA, so  $(tA)_p$  is bounded. Thus  $(tA)_p$  is a bounded pure subgroup of *A* hence a direct summand of *A*. Since summands of self-small groups are self-small and since self-small torsion groups are finite,  $(tA)_p$  is finite.

Since A is self-small, every endomorphic image of A is selfsmall. Let B = A/tA and let F be a free subgroup of B with B/Ftorsion. Now B/F is a homomorphic image of A, so  $(B/F)_p$  must be divisible for all but a finite number of primes with  $(tA)p \neq 0$  (or else A has an infinite torsion endomorphic image, a con-tradiction). Consequently,  $B/F = G \bigoplus T$  where T is finite and G is p-divisible for all primes p with  $(tA)_p \neq 0$ . This proves (b).

 $(\Leftarrow)$  We prove that End(A) is countable and apply Corollary 1.4. Let F be a free subgroup of A such that A/F is torsion. By (b),  $A/F + tA \simeq (A/tA)/((F + tA)/tA) = G \oplus T$ , where T is finite and G is *p*-divisible for almost all primes *p* with  $(tA)_p \neq 0$ . Moreover. Hom (A/F, tA) is countable. In view of the exact sequence  $0 \rightarrow \text{Hom}(A/F, tA) \rightarrow \text{Hom}(A, tA) \rightarrow \text{Hom}(F, tA)$  and the fact that F is finitely generated and tA is countable, one sees that Hom(A, tA) is countable. Furthermore, there exact sequences are  $0 \rightarrow \text{Hom}(A, tA) \rightarrow \text{Hom}(A, A) \rightarrow \text{Hom}(A, A/tA)$ and  $0 \rightarrow \text{Hom}(A/tA, A/tA) \rightarrow \text{Hom}(A, A/tA) \rightarrow \text{Hom}(tA, A/tA) = 0$ and Hom (A/tA, A/tA) is countable. Consequently, Hom (A, A) is countable, as desired.

COROLLARY 3.7. Suppose that A is a self-small mixed abelian group. Then for all primes, p,  $(tA)_p$  is finite or zero; the cardinality of A/pA is finite for all primes p with  $(tA)_p \neq 0$ ; and if B is a torsion free subgroup of A such that A/B is torsion, then A/B is p-divisible for all but a finite number of primes with  $(tA)_p \neq 0$ .

4. Locally A-projective groups. We first remind the reader of the homological setting described in [1]. Let  $\mathscr{G}$  be the category of abelian groups,  $A \in \mathscr{G}$ ,  $R = \operatorname{End}(A)$  and  $\mathcal{M}_R$  the category of right *R*-modules. There is a left exact functor  $H: \mathscr{G} \to \mathcal{M}_R$  defined by  $H(G) = \operatorname{Hom}_Z(A, G)$  and a right exact functor  $T: \mathcal{M}_R \to \mathscr{G}$  defined by  $T(M) = M \otimes_R A$ , where A is regarded as a left *R*-module in the obvious fashion.

There are natural transformations  $\theta: TH \to 1_{\mathscr{G}}$  and  $\phi: 1_{\mathscr{M}_R} \to HT$ , where  $\theta_G: \operatorname{Hom}_Z(A, G) \bigotimes_R A \to G$  is defined by  $\theta_G(f \otimes a) = f(a)$  and  $\phi_M: M \to \operatorname{Hom}_Z(A, M \otimes_R A)$  is defined by  $\phi_M(x)(a) = x \otimes a$ .

A group, G, is finitely A-projective if G is isomorphic to a direct summand of the direct sum of a finite number of copies of A.

THEOREM 4.1. Let A be an abelian group.

(a) the category of finitely A-projective groups is equivalent to the category of finitely generated projective right End(A)-modules.

(b) If A is self-small, then the category of A-projective groups is equivalent to the category of projective right End(A)-modules.

**Proof.** The following observations suffice to prove the theorem (all of which are easy to verify): (i) if G is *finitely A-free* (i.e., a finite direct sum of copies of A) or if G is A-free and A is self-small, then H(G) is a free right R-module; (ii) If M is a free right R-module, then T(M) is A-free; (iii)  $\theta_A: TH(A) \rightarrow A$  and  $\phi_R: R \rightarrow HT(R)$  are isomorphisms and (iv)  $\theta$  and  $\phi$  are natural transformations (details are given in [1]; see also Warfield [15] for the case that A is torsion free of rank 1).

COROLLARY 4.2. Let A be an abelian group.

(a) If A is self-small, then every A-projective group is A-free iff every projective right End(A)-module is free.

(b) Every locally A-projective group is locally A-free iff every finite generated projective right End(A)-module is free.

COROLLARY 4.3. Let  $\mathcal{F} = \{A_i\}_{i \in I}$  be a class of countable self-small groups together with a partial ordering on the index set I such that  $i \leq j$ iff Hom $(A_i, A_j) \neq 0$ . Define  $\mathcal{F}_{\epsilon}$  to be the class of groups isomorphic to direct sums of groups in  $\mathcal{F}$ . If every projective right End $(A_i)$ -module is free for all  $A_i \in \mathcal{F}$ , then  $\mathcal{F}_{\epsilon}$  is closed under direct summands.

*Proof.* Theorems due to Kulikov-Fuchs (see Charles [2]) and Kaplansky (see Fuchs [6], p. 49) reduce the argument to the  $A_{i-}$  projective case. Now apply Corollary 4.2.a.

COROLLARY 4.4. Suppose that A is torsion free and that End(A) is a principal ideal domain. If B is a subgroup of an A-free group G and  $S_A(B) = B$ , then B is A-free.

**Proof.** A is self-small by (3.2.a) and so H(B) is an End(A)-submodule of the free End(A)-module H(G). Now use the naturality of  $\theta$  to show  $\theta_G: T \cdot H(G) \rightarrow G$  is an isomorphism (details are given in [1]).

We observe that Corollary 4.3 includes the Baer-Kulikov-Kaplansky theorem (i.e., direct summands of completely decomposable groups are completely decomposable) results of Murley [10] and Arnold-Lady [1] as special cases. Corollary 4.4 is a generalization of theorems by Baer-Kolettis for the case that A has rank 1 (see Fuchs [6], p. 114].

LEMMA 4.5. Suppose that A is an abelian group such that  $\operatorname{Hom}_{Z}(A, G)$  is a locally projective right  $\operatorname{End}(A)$ -module for all locally A-projective groups G. Then the category of locally A-projective abelian groups is equivalent to the category of locally projective right  $\operatorname{End}(A)$ -modules.

**Proof.** Let  $\mathscr{L}_A$  be the category of locally A-projective abelian groups and  $\mathscr{L}_R$  the category of locally projective right R-modules, where R = End(A). The hypotheses guarantee that  $H: \mathscr{L}_A \to \mathscr{L}_R$  is well defined. To show that  $T: \mathscr{L}_R \to \mathscr{L}_A$  is well defined let  $M \in \mathscr{L}_R$  and  $y_1, \dots, y_m \in T(M) = M \otimes_R(A)$ , where  $y_i = \sum m_{ij} \otimes a_{ij}$  with  $m_{ij} \in M$ ,  $a_{ij} \in A$ . Now  $\{m_{ij}\}$  is contained in a finitely generated projective summand, P, of M, say  $M = P \oplus L$ . But  $T(M) = T(P) \oplus T(L)$  and  $\{y_1, \dots, y_m\} \subset T(P)$  a finitely A-projective summand of T(M) (Theorem 4.1.a). Next we prove that HT is naturally equivalent to the identity functor on  $\mathscr{L}_{\mathbb{R}}$ , i.e., if  $M \in \mathscr{L}_{\mathbb{R}}$ , then  $\phi_{\mathbb{M}} \colon M \to HT(M)$  is an isomorphism. Assume that  $x \in \ker \phi_{\mathbb{M}}$  and embed x in a finitely generated projective summand P of M. Naturality of  $\phi$  gives a commutative diagram.



The bottom row is exact since P is a summand of M. Since P is finitely generated and projective,  $\phi_P$  is an isomorphism. But  $x \in P$  and  $\phi_M(x) = 0$ , so x = 0. Consequently,  $\phi_M$  is monic.

To prove that  $\phi_M$  is epic, let  $f \in HT(M)$ . Since  $HT(M) \in \mathscr{L}_R$ , f is an element of some finitely generated projective summand,  $P_1$ , of HT(M). Let  $P = \{x \in M | \phi_M(x) \in P_1\}$ . Since  $\phi$  is natural,  $P_1$  is the image of HT(P) in HT(M). Moreover,  $\phi_P : P \to HT(P)$  is epic and  $f \in P_1$ , so  $f \in \text{image } \phi_M$  as desired.

Finally, we prove that  $\theta_G: TH(G) \to G$  is an isomorphism for all  $G \in \mathscr{L}_A$ . Let  $g \in G$  and embed g in a finitely A-projective summand B of G. Now  $\theta_B: TH(B) \to B$  is an isomorphism,  $g \in \operatorname{im} \theta_B$ , and  $\theta$  is natural, so  $g \in \operatorname{image} \theta_G$ , i.e.,  $\theta_G$  is epic. To show that  $\theta_G$  is monic let  $y = \sum_{i=1}^n f_i \otimes a_i \in \ker \theta_G$ , where  $f_i \in H(G)$  and  $a_i \in A$ . Since  $H(G) \in \mathscr{L}_R$ ,  $\{f_1, \dots, f_n\}$  is contained in a finitely generated projective R-summand, P, of H(G). Let  $B = \{\theta_G(x) | x \in P \otimes_R A \subset TH(G)\}$ . It now follows that  $P \otimes_R A = \operatorname{Hom}(A, B) \otimes_R A$  and B is finitely A-projective so that  $\theta_B: TH(B) \to B$  is an isomorphism and  $y \in \ker \theta_B$ , i.e., y = 0. The proof is now complete.

**Proof of Theorem III.** In view of Lemma 4.5, it suffices to prove that if G is an locally A-projective group and if  $f_1, \dots, f_n \in \text{Hom}(A, G)$ , then  $\{f_1, \dots, f_n\}$  is contained in a finitely generated End(A)-projective summand of Hom(A, G).

Let B be a finitely generated subgroup of A with  $B^* = 0$ . By the hypotheses,  $f_1(B) + \cdots + f_n(B)$  is contained in a finitely A-projective summand,  $G_1$ , of G, say  $G = G_1 \oplus G'_1$ . Since Hom $(A, G_1)$  is a finitely generated projective End(A)-summand of Hom(A, G), it is enough to prove that  $f_1, \cdots, f_n \in$  Hom $(A, G_1)$ . Suppose not; choose  $a \in A \setminus B$ and  $i \leq n$  such that  $\prod' f_i(a) \neq 0$ , where  $\prod' : G \to G'_1$  is the projection map. Now  $\prod' f_i(a)$  is contained in a finitely A-projective summand  $G_2$ of G, say  $G = G_2 \oplus G'_2$ . Furthermore,  $\sigma \prod' f_n(a) \neq 0$  where  $\sigma : G \to G_2$ is the projection map. Since  $G_2$  is A-projective, there is some  $\delta : G_2 \to A$  with  $g(a) \neq 0$  where  $g = \delta \sigma \prod' f_n \in$  End(A). Thus  $g \neq 0$  and g(B) = 0 (since  $f_i(B) \subset G_1$ ), a contradiction. We apply the preceding results in the case that End(A) is a principal ideal domain (hence discrete by Proposition 3.2.a).

Proof of Corollary IV. (a) Since End(B) is discrete, there is a finite subset X of B with  $X^* = 0 \in \text{End}(B)$ . Embed X in a finitely A-free summand, C of G say  $G = C \oplus D$ . We prove that  $B \subseteq$ C. Suppose not and let  $\delta: G \to D$  be the projection map. There is some  $\psi: D \to A$  with  $\psi \delta(B) \neq 0$  (D is A-free since G is A-free and R = End(A) is a principal ideal domain). But  $S_A(B) = B$  so there is  $\phi: A \to B$  with  $0 \neq \phi \psi \delta \in \text{End}(B)$ . Consequently,  $0 \neq \phi \psi \delta \in X^*$ (since  $X \subseteq C$ ) a contradiction.

One can easily show that Hom(A, B) is a pure *R*-submodule of Hom(A, C) (since *B* is pure in *C* and *A/im f* is torsion for all  $0 \neq f \in R$ ). But Hom(A, C) is a finitely generated free *R*-module and thus Hom(A, B) is a summand of Hom(A, C) (recall that *R* is a principal ideal domain). The naturality of  $\phi$  and Theorem 4.1.a guarantees that  $\phi_B$ :  $\text{Hom}(A, B) \otimes_R A \to B$  is an isomorphism and that *B* is an *A*-free summand of *C*, hence of *G*.

(b)  $(\Rightarrow)$  Let  $\mathscr{F}$  be the set of projections of the locally A-free group, G, onto A. For each  $\phi \in \mathscr{F}$  define  $A_{\phi} = \phi(A)$  and let  $e: G \to \prod_{\phi \in \mathscr{F}} A_{\phi}$  be given by  $e(x) = (\phi(x))_{\phi \in \mathscr{F}}$ . Clearly, e is monic and  $G = S_A(G) \subseteq S_A(\prod_{\phi \in \mathscr{F}} A_{\phi})$  (observe that G is locally A-free and that  $S_A$ commutes with direct sums). The purity of G is routine (observe that the characteristic of x in G is the minimum of the characteristics of  $\phi(x)$  in  $A_{\phi}$  as  $\phi$  varies over  $\mathscr{F}$ ).

( $\Leftarrow$ ) We first prove that  $P = S_A(\prod_{i \in I} A_i)$  is locally A-free. Note that Hom (A, P) is R-isomorphic to  $\prod$  Hom  $(A, A_i)$ , a locally free R-module (Chase [3]). By Theorem III, it suffices to prove that  $\theta_p$ : Hom  $(A, P) \otimes_{\mathbb{R}} A \to P$  is an isomorphism. Clearly,  $\theta_p$  is epic since  $S_A(P) = P$ . Suppose that ker  $\theta_p \neq 0$ ; let  $0 \neq x \in f_1 \otimes a_1 + \cdots + f_n \otimes a_n \in \ker \theta_p$  with n minimal and  $f_i: A \to P$ ,  $a_i \in A$ . Since n is minimal, R is a principal ideal domain, and A is a torsion free R-module (Proposition 3.2.a) it follows that  $\{a_1, \dots, a_n\}$  is an R-independent subset of A. Now  $\theta_p(x) = f_1(a_1) + \cdots + f_n(a_n) = 0$ . For each  $i \in I$ , let  $\prod_i: P \to A_i = A$  be the projection map so that  $\prod_i f_j \in R$  for all  $i \leq j \leq n$ . But  $0 = \prod_i f_1(a_1) + \cdots + \prod_i f_n(a_n)$  so that  $\prod_i f_j = 0$ for all  $i \in I$ ,  $i \leq j \leq n$ . Consequently,  $f_1 = f_2 = \cdots = f_n = 0$  and x = 0, a contradiction. This proves that  $\theta_p$  is monic.

Now Hom (A, G) is a pure *R*-submodule of Hom (A, P) (since *G* is pure in *P* and *A*/*im f* is torsion for all  $0 \neq f \in R$ ), so Hom (A, G) is locally *R*-free (Chase [3]). Naturality of  $\theta$  guarantees that  $\theta_G$ : Hom  $(A, G) \otimes_R A \to G$  is an isomorphism so that *G* is locally *A*-free (by Theorem III).

(c) A consequence of (a) and (b).

(d) Let G be a countable locally A-free group and write  $G = \{x_i | i = 1, 2, 3, \dots\}$ . There is an A-free summand  $G_1$  of G, say  $G = G_1 \bigoplus H_1$ , with  $x_1 \in G_1$ . By (c),  $H_1$  is locally A-free. Let  $y_2$  be the

projection of  $x_2$  in  $H_1$ . If  $y_2 = 0$  define  $G_2 = G_1$ . Otherwise, let  $G_2$  be an A-free summand of  $H_1$  containing  $y_2$ . Proceed inductively to construct  $G_n$  for each n > 0 so that  $G = \sum_{n=1}^{\infty} \bigoplus G_n$  and G is A-free.

**REMARK.** P is locally A-free if End(A) is a principal ideal domain, but is not A-free if I is infinite and A is reduced. Corollary IV.d. includes Theorem 1 of Murley [10].

The discussion in this section has been restricted to Abelian groups in order to obtain sharpness in examples and applications. However, much remains true for R-modules. Let R be a commutative ring with 1 and A be an R-module. Define A-projective [A-free] in the obvious way and note that the homological setting remains intact. It follows that Theorem 4.1 and Corollary 4.2 hold for an *R*-module *A*. Suppose **R** and  $\operatorname{End}_{R}(A)$  are Dedekind domains. Then it is easily seen that any  $\operatorname{End}_{R}(A)$ -injective module is an *R*-injective (= R - divisible)module. This observation together with the arguments involved show that Proposition 3.2.a and Corollary 4.4 hold for an *R*-module A with R a Dedekind domain.

### 5. Examples.

**EXAMPLE 5.1.** There is a self-small torsion free abelian group, A, such that End(A) is not discrete in the finite topology.

**Proof.** Let p be a prime,  $Z_p$  the localization of Z at p, and let  $A_p$  be a free  $Z_p$ -module of rank p. Then  $A = \prod_p A_p$  is self-small (Corollary 1.3). Assume that the finite topology on End(A) is discrete, i.e.,  $X^* = 0$  for some finite subset, X, of A. Suppose that n is the cardinality of X. For each prime p, let  $\pi_p: A \to A_p$  be the projection map; the cardinality of  $\pi_p(X) \leq n$ . Now  $0 = (\pi_p(X))^* \subseteq \text{End}(A_p)$ . On the other hand, for each p > n,  $\pi_p(X)$  is contained in a  $Z_p$ -free summand of  $A_p$  of rank at most n, so  $0 \neq (\pi_p(X))^* \subseteq \text{End}(A_p)$ , a contradiction. Thus End(A) is not discrete.

EXAMPLE 5.2. There is a countable torsion free abelian group, A, such that End(A) is discrete and End(A) fails to have the minimum condition on left annihilator ideals. (compare Proposition 2.4.a)

**Proof.** Corner [4], constructs a countable torsion free abelian group, A, such that End(A) is countable and A has no nonzero indecomposable summands. Now apply Corollary 2.3 and Proposition 2.4.c.

EXAMPLE 5.3. There is a countable self-small torsion free abelian group, A, such that the cardinality of A/pA is infinite for all primes p (compare Corollary 3.3.b).

**Proof.** Let R = Z[X], the polynomial ring in an indeterminate X. By Corner [4], there is a countable torsion free abelian group A with End(A) = R. Clearly, A/pA is infinite for all primes, p.

### REFERENCES

1. D. M. Arnold and E. L. Lady, Endomorphism rings and direct sums of torsion free Abelian groups, to appear.

2. B. Charles, *Sous-groupes Fonctoriels et Topologies*, Studies on Abelian Groups, 75–92 (Paris), 1968.

3. S. U. Chase, Locally free modules and a problem of Whitehead, Illinois J. Math., 6 (1962), 682–699.

4. A. L. S. Corner, Every countable reduced Torsion-free ring is an endomorphism ring, Proc. London Math. Soc., 13 (1963), 687-710.

5. \_\_\_\_\_, Endomorphism rings of torsion-free Abelian groups, Proc. Internat. Conf. Theory Groups, (New York), 1967, 59-69.

6. L. Fuchs, Infinite Abelian Groups, Vol. I, Academic Press (New York, 1970.

7. \_\_\_\_, Infinite Abelian Groups, Vol. II, Academic Press (New York, 1973.

8. I. Kaplansky, Modules over Dedekind domains and valuation rings, Trans. Amer. Math. Soc., 73 (1952), 327-340.

9. C. Murley, The classification of certain classes of torsion free Abelian groups, Pacific J. Math., 40 (1972), 647–665.

10. \_\_\_\_\_, Direct products and sums of torsion free Abelian groups, Proc. Amer. Math. Soc., 38 (1973), 235-241.

11. J. Reid, On the Ring of Quasi-endomorphisms of a Torsion-free Group, Topics in Abelian Groups, (Chicago), 1963, 51-68.

12. F. Richman, A class of rank two torsion free groups, Studies on Abelian Groups, (Paris), 1968 327-333.

13. T. Szele, On a topology in endomorphism rings of Abelian groups, Publ. Math. Debrecen, 5 (1957), 1-4.

14. E. Walker, Quotient Categories and Quasi-isomorphisms of Abelian Groups, Proc. Colloq, Abelian Groups, (Budapest), 1964, 147–162.

15. R. B. Warfield, Jr., Homomorphisms and duality for torsion free groups, Math. Z., 107 (1968), 189–200.

Received November 14, 1973 and in revised form December 4, 1973. The second author was partially supported in this research by the National Research Council of Canada; Grant No. A-8073.

NEW MEXICO STATE UNIVERSITY AND UNIVERSITY OF VICTORIA

# Pacific Journal of Mathematics Vol. 56, No. 1 November, 1975

Shimshon A. Amitsur, Central embeddings in semi-simple rings	1
David Marion Arnold and Charles Estep Murley, Abelian groups, A, such	
that $Hom(A,)$ preserves direct sums of copies of A	7
Martin Bartelt, An integral representation for strictly continuous linear	
operators	21
Richard G. Burton, Fractional elements in multiplicative lattices	35
James Alan Cochran, Growth estimates for the singular values of	
square-integrable kernels	51
C. Martin Edwards and Peter John Stacey, On group algebras of central group extensions	59
Peter Fletcher and Pei Liu, <i>Topologies compatible with homeomorphism</i>	
groups	77
George Gasper, Jr., <i>Products of terminating</i> ${}_{3}F_{2}(1)$ <i>series</i>	87
Leon Gerber, The orthocentric simplex as an extreme simplex	97
Burrell Washington Helton, A product integral solution of a Riccati	
equation	113
Melvyn W. Jeter, On the extremal elements of the convex cone of	
superadditive n-homogeneous functions	131
R. H. Johnson, Simple separable graphs	143
Margaret Humm Kleinfeld, More on a generalization of commutative and	
alternative rings	159
A. Y. W. Lau, <i>The boundary of a semilattice on an n-cell</i>	171
Robert F. Lax, The local rigidity of the moduli scheme for curves	175
Glenn Richard Luecke, A note on quasidiagonal and quasitriangular	
operators	179
Paul Milnes, On the extension of continuous and almost periodic	
functions	187
Hidegoro Nakano and Kazumi Nakano, Connector theory.	195
James Michael Osterburg, Completely outer Galois theory of perfect	
rings	215
Lavon Barry Page, Compact Hankel operators and the F. and M. Riesz	
theorem	221
Joseph E. Quinn, Intermediate Riesz spaces	225
Shlomo Vinner, Model-completeness in a first order language with a	
generalized quantifier	265
Jorge Viola-Prioli, On absolutely torsion-free rings	275
Philip William Walker, A note on differential equations with all solutions of	
integrable-square	285
Stephen Jeffrey Willson, Equivariant maps between representation	
spheres	291