

Pacific Journal of Mathematics

**COMPACT HANKEL OPERATORS AND THE F. AND M. RIESZ
THEOREM**

LAVON BARRY PAGE

COMPACT HANKEL OPERATORS AND THE F. AND M. RIESZ THEOREM

LAVON PAGE

The F. and M. Riesz theorem asserts that every complex Borel measure on the unit circle whose Fourier coefficients with negative index vanish is necessarily absolutely continuous with respect to Lebesgue measure.

The purpose of this note is to give a new proof of Hartman's theorem on compact Hankel operators which clarifies the general context of the theorem. The proof depends only on a few simple operator-theoretic results, Nehari's characterization of bounded Hankel operators, and the aforementioned theorem of F. and M. Riesz.

A Hankel operator on the Hardy space \mathcal{H}^2 of functions on the unit circle is an operator of the form

$$H_\phi : f(e^{it}) \rightarrow P_+ \phi(e^{-it}) f(e^{-it})$$

where ϕ is a fixed function in \mathcal{L}^∞ and P_+ is the orthogonal projection of \mathcal{L}^2 onto \mathcal{H}^2 . P. Hartman proved in 1958 that if H_ϕ above is compact, then ϕ can be chosen to be continuous [2].

Let S be the usual unilateral shift on \mathcal{H}^2 , $S : f(e^{it}) \rightarrow e^{it} f(e^{it})$. Nehari had proved earlier that every bounded solution to the operator equation $S^*H = HS$ is of the form $H = H_\phi$, where $\phi \in \mathcal{L}^\infty$, and had further shown that ϕ could be chosen so that $\|H\| = \|\phi\|_\infty$ [3].

R. N. Hevener [3] and D. Sarason [5] have previously used the Riesz theorem in obtaining Hartman's characterization of compact Hankel operators. Their procedure involves the factorization of analytic functions. T. L. Kriete motivated this paper by suggesting that Lemma 1 below might be used to make more precise the relationship between compact Hankel operators and the F. and M. Riesz theorem.

Let \mathcal{C} denote the space of complex-valued continuous functions on the unit circle, and \mathcal{A}_0 the subspace of \mathcal{C} consisting of those functions which extend continuously to an analytic function on the unit disk vanishing at the origin. With the F. and M. Riesz theorem and the Riesz representation theorem for bounded linear functionals on \mathcal{C} , one easily shows that the Hardy space \mathcal{H}^1 is the dual space of $\mathcal{C}/\mathcal{A}_0$. Let \mathcal{H}_0^∞ denote the class of \mathcal{H}^∞ functions which vanish at zero. Since \mathcal{L}^∞ is

the dual of \mathcal{L}^1 , $\mathcal{L}^\infty/\mathcal{H}_0^\infty$ is the dual of \mathcal{H}^1 . (This is the case because \mathcal{H}_0^∞ is the annihilator of \mathcal{H}^1 .)

Now let Ψ be the mapping from \mathcal{L}^∞ to the space $\mathcal{H}_\mathfrak{B}$ of bounded Hankel operators on \mathcal{H}^2 given by $\Psi: \phi \rightarrow H_\phi$. It is elementary to see that $\ker \Psi = \mathcal{H}_0^\infty$. Thus the Nehari theorem states that the mapping $\Psi^-: \mathcal{L}^\infty/\mathcal{H}_0^\infty \rightarrow \mathcal{H}_\mathfrak{B}$ given by $\phi + \mathcal{H}_0^\infty \rightarrow H_\phi$ is an isomorphism of Banach spaces.

The converse of Hartman's theorem is easy, that is H_ϕ is compact if ϕ is continuous. The reason is that if ϕ is continuous, then ϕ is a uniform limit of trigonometric polynomials $\{\phi_n\}$, in which case $H_{\phi_n} \rightarrow H_\phi$ in operator norm. Compactness of H_ϕ now follows from the observation that each H_{ϕ_n} is of finite rank.

We now let θ denote the restriction of Ψ to \mathcal{C} . Since H_ϕ is compact when ϕ is continuous, θ maps \mathcal{C} into $\mathcal{H}_\mathfrak{C}$, the space of compact Hankel operators. Also $\ker \theta = \mathcal{A}_0$, so $\theta^-: \mathcal{C}/\mathcal{A}_0 \rightarrow \mathcal{H}_\mathfrak{C}$ defined by $\theta^-: \phi + \mathcal{A}_0 \rightarrow H_\phi$ is one-to-one. Furthermore, if $\phi \in \mathcal{C}$, the norm of the coset $\phi + \mathcal{H}_0^\infty$ in $\mathcal{L}^\infty/\mathcal{H}_0^\infty$ is the same as the norm of the operator H_ϕ . (This is the Nehari theorem.) But since the natural mapping of $\mathcal{C}/\mathcal{A}_0$ into $\mathcal{L}^\infty/\mathcal{H}_0^\infty$ is an isometry, $\|H_\phi\|$ is equal to the norm of $\phi + \mathcal{A}_0$ in $\mathcal{C}/\mathcal{A}_0$. Thus the mapping θ^- is an isometry mapping $\mathcal{C}/\mathcal{A}_0$ into $\mathcal{H}_\mathfrak{C}$. We now have the following commutative diagram:

$$\begin{array}{ccc}
 \text{canonical embedding} & & \text{inclusion} \\
 \mathcal{C}/\mathcal{A}_0 & \longrightarrow & \mathcal{L}^\infty/\mathcal{H}_0^\infty \\
 \theta^- \downarrow & & \downarrow \Psi^- \\
 \mathcal{H}_\mathfrak{C} & \longrightarrow & \mathcal{H}_\mathfrak{B}
 \end{array}$$

The mapping Ψ^- is an isomorphism. From the two lemmas below it now follows that the range of θ^- is $\mathcal{H}_\mathfrak{C}$, and this is precisely Hartman's result.

LEMMA 1. *The Banach space $\mathcal{H}_\mathfrak{B}$ is isomorphic to the second dual space of $\mathcal{H}_\mathfrak{C}$ in such a way that the inclusion mapping $\mathcal{H}_\mathfrak{C} \rightarrow \mathcal{H}_\mathfrak{B}$ corresponds to the canonical embedding of a normed space in its second dual.*

Proof. This follows from an abstract result due to Gellar and Page [1]. Primarily, it is just a matter of checking that every Hankel operator is the limit in the weak operator topology of a sequence of compact Hankel operators. The important point is that this can be checked *without* prior knowledge of which Hankel operators are compact. See Theorems 1 and 2 of [1].

LEMMA 2. Let \mathcal{X} and \mathcal{Y} be normed linear spaces, and let $A : \mathcal{X} \rightarrow \mathcal{Y}$ and $B : \mathcal{X}^{**} \rightarrow \mathcal{Y}^{**}$ be linear isometries such that the diagram :

canonical embedding canonical embedding

$$\begin{array}{ccc} \mathcal{X} & \longrightarrow & \mathcal{X}^{**} \\ A \downarrow & & \downarrow \\ \mathcal{Y} & \longrightarrow & \mathcal{Y}^{**} \end{array}$$

is commutative.

If the range of B is \mathcal{Y}^{**} , then the range of A is \mathcal{Y} .

Proof. This is a simple consequence of the Hahn-Banach Theorem and the fact that a normed linear space is weak-star dense in its second dual.

REFERENCES

1. R. Gellar and L. Page, *A new look at some familiar spaces of intertwining operators*, Pacific J. Math., **47** (1973), 435-441.
2. P. Hartman, *On completely continuous Hankel matrices*, Proc. Amer. Math. Soc., **9** (1958), 862-866.
3. R. Hevener, *A functional analytic approach to Hankel and Toeplitz matrices*, Thesis, University of Virginia, 1965.
4. Z. Nehari, *On bounded bilinear forms*, Ann. of Mathematics, **65** (1957), 153-162.
5. D. Sarason, *Generalized interpolation in \mathcal{H}^∞* , Trans. Amer. Math. Soc., **127** (1967), 179-203.

Received September 25, 1973.

NORTH CAROLINA STATE UNIVERSITY

Pacific Journal of Mathematics

Vol. 56, No. 1

November, 1975

Shimshon A. Amitsur, <i>Central embeddings in semi-simple rings</i>	1
David Marion Arnold and Charles Estep Murley, <i>Abelian groups, A, such that $\text{Hom}(A, - - -)$ preserves direct sums of copies of A</i>	7
Martin Bartelt, <i>An integral representation for strictly continuous linear operators</i>	21
Richard G. Burton, <i>Fractional elements in multiplicative lattices</i>	35
James Alan Cochran, <i>Growth estimates for the singular values of square-integrable kernels</i>	51
C. Martin Edwards and Peter John Stacey, <i>On group algebras of central group extensions</i>	59
Peter Fletcher and Pei Liu, <i>Topologies compatible with homeomorphism groups</i>	77
George Gasper, Jr., <i>Products of terminating ${}_3F_2(1)$ series</i>	87
Leon Gerber, <i>The orthocentric simplex as an extreme simplex</i>	97
Burrell Washington Helton, <i>A product integral solution of a Riccati equation</i>	113
Melvyn W. Jeter, <i>On the extremal elements of the convex cone of superadditive n-homogeneous functions</i>	131
R. H. Johnson, <i>Simple separable graphs</i>	143
Margaret Humm Kleinfeld, <i>More on a generalization of commutative and alternative rings</i>	159
A. Y. W. Lau, <i>The boundary of a semilattice on an n-cell</i>	171
Robert F. Lax, <i>The local rigidity of the moduli scheme for curves</i>	175
Glenn Richard Luecke, <i>A note on quasidiagonal and quasitriangular operators</i>	179
Paul Milnes, <i>On the extension of continuous and almost periodic functions</i>	187
Hidegoro Nakano and Kazumi Nakano, <i>Connector theory</i>	195
James Michael Osterburg, <i>Completely outer Galois theory of perfect rings</i>	215
Lavon Barry Page, <i>Compact Hankel operators and the F. and M. Riesz theorem</i>	221
Joseph E. Quinn, <i>Intermediate Riesz spaces</i>	225
Shlomo Vinner, <i>Model-completeness in a first order language with a generalized quantifier</i>	265
Jorge Viola-Prioli, <i>On absolutely torsion-free rings</i>	275
Philip William Walker, <i>A note on differential equations with all solutions of integrable-square</i>	285
Stephen Jeffrey Willson, <i>Equivariant maps between representation spheres</i>	291