

Pacific Journal of Mathematics

INTEGRAL REPRESENTATIONS OF WEAKLY COMPACT OPERATORS

A. KATSARAS AND DAR-BIAU LIU

INTEGRAL REPRESENTATIONS OF WEAKLY COMPACT OPERATORS

A. KATSARAS AND D. B. LIU

Let X be a completely regular space and E, F locally convex spaces. Denote by $C_{rc} = C_{rc}(X, E)$ the space of all continuous functions f from X into E for which $f(X)$ is relatively compact. Uniformly continuous weakly compact operators from C_{rc} into F are represented by integrals with respect to $\mathcal{L}(E, F)$ valued measures on the algebra generated by the zero sets. Necessary and sufficient conditions for an operator to be continuous, with respect to certain topologies, are obtained. A sufficient condition for extending a measure to all Baire sets is given.

Introduction. In [5] D. Lewis represented weakly compact operators from the space $C(S)$ of all scalar-valued continuous functions on a compact space into a locally convex space. The representation was given by means of integrals with respect to vector-valued measures on the Borel field. In [1] Bartle, Dunford, and Schwartz gave a similar representation for operators from $C(S)$ into a Banach space. Also Grothendieck [2] noted that the family of all weakly compact operators from $C(S)$ into a locally convex space E corresponds exactly to the family of E -valued measures on the Baire algebra. In this paper we will give integral representations for weakly compact operators from C_{rc} into F by means of integrals with respect to $\mathcal{L}(E, F)$ valued measures on the algebra generated by the zero sets. Necessary and sufficient conditions for an operator to be continuous with respect to certain locally convex topologies are given. Also a result is obtained on the extension of measures to all Baire sets.

1. Definitions and preliminaries. Let X be a completely regular Hausdorff space and let $B = B(X)$ denote the algebra of subsets of X generated by the zero sets. By $Ba = Ba(X)$ and $Bo = Bo(X)$ we will denote the σ -algebras of Baire and Borel sets respectively. Let $M(X)$ denote the space of all bounded finitely-additive regular (with respect to the zero sets) measures on B (see Varadarajan [8]). The spaces of all σ -additive and all τ -additive members of $M(X)$ will be denoted by $M_\sigma(X)$ and $M_\tau(X)$ respectively. The set $M_\sigma(Ba)$ is the space of all real-valued Baire measures while $M_\tau(Bo)$ denotes the space of all bounded real-valued regular Borel measures m with the property

that $m(G\alpha) \rightarrow 0$ for every net $\{G_\alpha\}$ of closed sets which decreases to the empty set.

Let E be a real locally convex Hausdorff space. For p a continuous seminorm on E , we define $M_p(B, E')$ as the set of all E' -valued (E' is the dual of E) finitely-additive measures m on B with the following two properties:

(1) For every $s \in E$, the function ms , from B into the reals $R, G \rightarrow m(G)s$, is in $M(X)$.

(2) $\|m\|_p = m_p(X) < \infty$, where for G in B the $m_p(G)$ is defined to be the supremum of all $|\sum m(G_i)s_i|$ for all finite B -partitions $\{G_i\}$ of G , i.e., $G_i \in B$, and all finite collections $s_i \in B_p = \{s \in E : p(s) \leq 1\}$. The set $M_{\sigma,p}(B, E')$ consists of those m in $M_p(B, E')$ for which $ms \in M_\sigma(X)$ for all s in E . The spaces $M_{\tau,p}(B, E')$, $M_{\sigma,p}(Ba, E')$, and $M_{\tau,p}(Bo, E')$ are defined similarly. As shown in [3] if m is in any one of the spaces $M_p(B, E')$, $M_{\sigma,p}(B, E')$, $M_{\tau,p}(B, E')$, $M_{\sigma,p}(Ba, E')$, $M_{\tau,p}(Bo, E')$, then m_p belongs to $M(X)$, $M_\sigma(X)$, $M_\tau(X)$, $M_\sigma(Ba)$, $M_\tau(Bo)$ respectively. Every $m \in M_{\sigma,p}(B, E')$ [$m \in M_{\tau,p}(B, E')$] has a unique extension μ to a member of $M_{\sigma,p}(Ba, E')$ [to a member of $M_{\tau,p}(Bo, E')$]. Moreover, the restriction of μ_p to B coincides with m_p . Let $\{p : p \in I\}$ be a generating family of continuous seminorms on E which is directed, i.e., given p_1, p_2 in I there exists $p \in I$ with $p \geq p_1, p_2$. Let $M(B, E') = \cup \{M_p(B, E') : p \in I\}$ with analogous definitions for $M_\sigma(B, E')$, $M_\tau(B, E')$, $M_\sigma(Ba, E')$ and $M_\tau(Bo, E')$.

Denote by $C_{rc} = C_{rc}(X, E)$ the space of all continuous functions f from X into E for which $f(X)$ is relatively compact. Every f in C_{rc} has a unique continuous extension \hat{f} to all of the Stone Cech compactification βX . By $C^b(X)$ we denote the space of all bounded continuous real-valued functions on X . Let Ω and Ω_1 be, respectively, the class of all compact and all zero sets in βX disjoint from X . Let $Q \in \Omega$. We define β_Q to be the locally convex topology generated by the family of seminorms $f \rightarrow \|fg\|_p = \sup\{p(f(x)g(x)) : x \in X\}$ where $p \in I$ and $g \in B_Q = \{h \in C^b : \hat{h}(x) = 0 \text{ for } x \text{ in } Q\}$. The topologies β and β_1 on C_{rc} are defined to be the inductive limits of the topologies β_Q as Q ranges over Ω and Ω_1 respectively. For a fixed $p, \beta_{p,Q}$ is the locally convex topology on C_{rc} generated by the seminorms $f \rightarrow \|gf\|_p, g \in B_Q$. As shown in [3], $\beta_{p,Q}$ is the finest locally convex topology on C_{rc} which agrees with $\beta_{p,Q}$ on p -bounded sets. Let β_p and $\beta_{1,p}$ denote the inductive limits of the topologies $\beta_{p,Q}$ as Q ranges over Ω and Ω_1 respectively. The topologies β' and β'_1 are the projective limits of the topologies β_p and $\beta_{1,p}$, respectively, as p ranges over I . If u is the uniform topology, then $\beta' \leq \beta \leq \beta_1 \leq u$ and $\beta'_1 \leq \beta_1$.

For G in B and $m \in M_p(B, E')$ we define $\int_G f dm = \lim \sum m(G_i)f(x_i)$ where the limit is taken over the directed set of all

finite B -partitions $\{G_i\}$ of G and $x_i \in G_i$. The map $f \rightarrow T_m(f) = \int_X f dm$ is a uniformly continuous linear functional on C_{rc} . Moreover, $\|m\|_p = \sup\{|T_m(f)| : \|f\|_p \leq 1\}$. The mapping $m \rightarrow T_m$ is a one-to-one linear map from $M(B, E')$ into $(C_{rc}, u)'$. The space $M_\sigma(B, E')$ is the dual space of each of the topologies β_1 and β'_1 while $M_\tau(B, E')$ is the dual space of each of the topologies β and β' . Given any $m \in M_p(B, E')$ there exists a unique \hat{m} in $M_{\tau,p}(Bo(\beta X), E')$ such that $\int_X f dm = \int_{\beta X} \hat{f} d\hat{m}$ for all f in C_{rc} . As shown in [3], m is σ -additive iff $\hat{m}_p(Z) = 0$ for all Z in Ω_1 . Similarly m is τ -additive iff $\hat{m}_p(Q) = 0$ for all Q in Ω . Moreover, if m is σ -additive or τ -additive, then $\hat{m}(Q) = m(Q \cap X)$ and $\hat{m}_p(Q) = \hat{m}_p(Q \cap X)$ for all Q in $B(\beta X)$.

Let now F be another real locally convex Hausdorff space and let $\{q : q \in J\}$ be a generating directed family of continuous seminorms on F . Let $\mathcal{L}(E, F)$ denote the space of all continuous operators from E into F . We define $M(B, \mathcal{L}(E, F))$ to be the space of all finitely-additive $\mathcal{L}(E, F)$ valued measures m on B with the following two properties:

(1) For each $x' \in F'$ the set function $x'm : B \rightarrow E'$, $(x'm)(G)s = x'(m(G)s)$, $s \in E$, is in $M(B, E')$.

(2) Given $q \in J$ there exists p in I such that for all x' in the polar B_q^0 of B_q in F' the $x'm$ is in $M_p(B, E')$ and $\|m\|_{p,q} = m_{p,q}(X) < \infty$ where for Q in B , $m_{p,q}(Q) = \sup\{(x'm)_p(Q) : x' \in B_q^0\}$. We define $M_\sigma(B, \mathcal{L}(E, F))$, $M_\tau(B, \mathcal{L}(E, F))$, $M_\sigma(Ba, \mathcal{L}(E, F))$ and $M_\tau(Bo, \mathcal{L}(E, F))$ analogously. Let $m \in M(B, \mathcal{L}(E, F))$ and f a function from X into E . We say that f is m -integrable over G in B if

(i) For each $x' \in F'$, the integral $\int_G f d(x'm)$ exists

(ii) there exists a vector in F denoted by $\int_G f dm$ such that for all $x' \in F'$ we have $x' \left(\int_G f dm \right) = \int_G f d(x'm)$.

Since F is a locally convex Hausdorff space, the $\int_G f dm$ is unique whenever it exists. If f is m -integrable over all G in B , we say that f is m -integrable.

2. Continuous linear operators from C_{rc} into F . Let $E, F, \{p : p \in I\}, \{q : q \in J\}$ be as in paragraph 1. Recall that a linear operator T from a topological vector space A into another B is weakly compact if it maps bounded subsets of A into weakly relatively compact subsets of B . We need the following lemma due to Grothendieck [2].

LEMMA 1. *Let T be an operator from a topological vector space A into another B and let T' and T'' denote, respectively, the transpose and the second transpose of T . The following are equivalent:*

- (1) *T is weakly compact*
- (2) *T'' maps A'' into B*
- (3) *If B' is equipped with the Mackey topology $m(B', B)$ and A' with the strong topology $\beta(A', A)$, then T' is continuous.*

LEMMA 2. *Let f_0 be in C_{rc} and G in B . Define ϕ on $M(B, E')$ by $\phi(m) = \int_G f_0 dm$. Then ϕ belongs to the $(C_{rc}, u)''$.*

Proof. Let $A = \{f \in C_{rc} : \|f\|_p \leq \|f_0\|_p \text{ for all } p \text{ in } I\}$. Then A is u -bounded and hence the polar A^0 in $(C_{rc}, u)'$ is a strong neighborhood of zero. We will finish the proof by showing that ϕ is bounded on A^0 . To this end consider an arbitrary m in A^0 . Let $\epsilon > 0$ be given. There exists a B -partition G_1, G_2, \dots, G_n of G and $x_i \in G_i$ such that $\left| \int_G f_0 dm \right| \leq |\sum m(G_i)s_i| + \epsilon$, $s_i = f_0(x_i)$. By the regularity of ms_i we can find zero sets $Z_i \subset G_i$ such that $|\sum m(G_i)s_i| \leq |\sum m(Z_i)s_i| + \epsilon$. Again by the regularity of $|ms_i|$ ($|ms_i|$ is the absolute variation of ms_i) we can find pairwise disjoint cozero sets U_1, \dots, U_n , $Z_i \subset U_i$ such that $\sum |ms_i|(U_i - Z_i) < \epsilon$. For each i choose $h_i \in C^b$, with $0 \leq h_i \leq 1$, such that $h_i = 1$ on Z_i and $h_i = 0$ on $X - U_i$. Set $h = \sum h_i s_i$. Then $h \in A$ and hence $\left| \int_X h dm \right| \leq 1$. But

$$\begin{aligned} \left| \int_X h dm \right| &\geq \left| \sum \int_{Z_i} h_i s_i dm \right| - \left| \sum \int_{U_i - Z_i} h_i d_i ms_i \right| \\ &\geq \left| \sum m(Z_i)s_i \right| - \epsilon \geq \left| \int f_0 dm \right| - 3\epsilon. \end{aligned}$$

Since $\epsilon > 0$ was arbitrary we conclude that $\left| \int_G f_0 dm \right| \leq 1$ and this completes the proof.

THEOREM 3. *If T is a continuous weakly compact operator from (C_{rc}, u) into F , then there exists a unique $m \in M(B, \mathcal{L}(E, F))$ such that:*

- (1) *Every f in C_{rc} is m -integrable and $\int_X f dm = T(f)$*
- (2) *If $p \in I$ and $q \in J$ are such that $\|T\|_{p,q} = \sup\{q(T(f)) : \|f\|_p \leq 1\} \leq \infty$, then $\|m\|_{p,q} = \|T\|_{p,q}$.*
- (3) *For every $x' \in F'$, we have $T'x' = x'm$*

(4) For every bounded set S in E the set $V_{m,S} = \{\Sigma m(G_i) s_i : \{G_i\} \text{ is a finite } B\text{-partition of } X, s_i \in S\}$ is weakly relatively compact. Conversely, if $m \in M(B, \mathcal{L}(E, F))$ is such that

(5) holds, then every f in C_{rc} is m -integrable and the operator $T(f) = \int_X f dm$ is u -continuous and weakly compact.

Proof. Suppose that T is u -continuous and weakly compact. By Lemma 1, T'' maps $(C_{rc}, u)''$ into F . If $f \in C_{rc}$ and G in B , the function $f\chi_G$ (χ_G is the characteristic function of G) defines an element of $(C_{rc}, u)''$ by $\langle \mu, f\chi_G \rangle = \int_G f d\mu$, $\mu \in M(B, E') = (C_{rc}, u)'$. Thus we may consider $f\chi_G$ as an element of $(C_{rc}, u)''$. Define $m(G): E \rightarrow F$ by $m(G)s = T''(\chi_G s)$, G in B . It is easy to see that $m(G) \in \mathcal{L}(E, F)$. In this way we define a map $m: B \rightarrow \mathcal{L}(E, F)$ which is clearly finitely additive. If $x' \in F'$ and s in E , then $(x'm)(G)s = x'(T''(\chi_G s)) = \langle T'x', \chi_G s \rangle = T'x'(G)s$. Thus $x'm = T'x' \in M(B, E')$. Let $q \in J$. Since T is u -continuous there exists $p \in I$ such that $\|T\|_{p,q} < \infty$. Let $x' \in B_q^0$. Then for f in C_{rc} with $\|f\|_p \leq 1$ we have $|\langle f, x'm \rangle| = |\langle f, T'x' \rangle| \leq |\langle Tf, x' \rangle| \leq \|T\|_{p,q}$. Thus $\|x'm\|_p \leq \|T\|_{p,q}$ which proves that $\|m\|_{p,q} \leq \|T\|_{p,q}$ and so m is in $M(B, \mathcal{L}(E, F))$. Let G be in B and $f \in C_{rc}$. For $x' \in F'$ we have $x'(T''(\chi_G f)) = \langle T'x', \chi_G f \rangle = \langle x'm, \chi_G f \rangle = \int_G f d(x'm)$. This shows that $\int_G f dm = T''(\chi_G f) \in F$. Taking $G = X$ we get $\int_X f dm = T''(f) = T(f)$. For $f \in C_{rc}$ with $\|f\|_p \leq 1$ and $x' \in B_q^0$ we have $|x'(T(f))| = |\int f d(x'm)| \leq \|x'm\|_p \leq \|m\|_{p,q}$. This proves that $\|T\|_{p,q} \leq \|m\|_{p,q}$. For the uniqueness of m , suppose m_1 is another element in $M(B, \mathcal{L}(E, F))$ such that $\int_X f dm_1 = T(f)$ for all $f \in C_{rc}$. Then for $x' \in F'$ we have $\int_X f d(x'm) = \int_X f d(x'm_1)$ for all f in C_{rc} . This implies that $x'm = x'm_1$ and hence $m = m_1$ since F is a locally convex Hausdorff space. Finally, let S be a bounded subset of E and $W = V_{m,S}$. Let $A = \{f \in C_{rc} : f(X) \subset S\}$. Then A is u -bounded and therefore $T(A)$ is weakly relatively compact. We will finish the proof of (4) by showing that E is contained in the weak closure of $T(A)$. Let G_1, \dots, G_n be a B -partition of X and s_1, \dots, s_n in S . Let $x'_1, \dots, x'_n \in F'$. There exist $q \in J$ and $M > 0$ such that $x'_i \in MB_q^0$. Let $p \in I$ be such that $\|T\|_{p,q} < \infty$. Since S is bounded, $d = \sup\{p(s) : s \in E\} < \infty$. By the regularity of $(x'_j m)_p$ we can find zero sets Z_1, \dots, Z_n with $\sum_{i=1}^n (x'_j m)_p(G_i - Z_i) < \epsilon/2d$ (where $\epsilon > 0$ is arbitrary) for $j = 1, \dots, N$. Next, again by regularity, we can find

pairwise disjoint cozero sets U_1, \dots, U_n with $Z_i \subset U_i$ such that for each $j, 1 \leq j \leq n$, we have $\sum_{i=1}^n (x'_j m)_p (U_i - Z_i) < \epsilon/2d$. For each i between 1 and n we pick a function $h_i \in C^b$ with $0 \leq h_i \leq 1$, such that $h_i = 1$ on Z_i and $h_i = 0$ on the complement of U_i . The function $h = \sum_{i=1}^n h_i s_i$ is in A and hence $T(h) \in T(A)$. Moreover

$$\begin{aligned} & \left| x'_j(T(h) - \sum m(G_i) s_i) \right| \\ &= \left| x'_j \left(\sum m(Z_i) s_i - \sum m(G_i) s_i + \sum \int_{U_i - Z_i} h_i s_i dm \right) \right| < \epsilon/2 + \epsilon/2 = \epsilon. \end{aligned}$$

This shows that $\sum m(G_i) s_i$ is in the weak closure of $T(A)$ and the proof of (4) is complete. Conversely, suppose that $m \in M(B, \mathcal{L}(E, F))$ satisfies (4). Let $G \in B$ and $f \in C_{rc}$. Denote by D_G the set of all $\alpha = \{G_1, \dots, G_n; x_1, \dots, x_n\}$ where $\{G_i\}$ is a B -partition of G and $x_i \in G_i$. For α, γ in D_G we write $\alpha \cong \gamma$ if the B -partition of G for α is a refinement of the one in γ . Then D_G becomes a directed set.

For $\alpha = \{G_1, \dots, G_n; x_1, \dots, x_n\}$ in D_G we define $z_\alpha = \sum m(G_i) f(x_i)$. By (4) the net $\{z_\alpha\}$ is contained in a weakly compact set. Hence there exists a subnet which converges weakly to a vector z in F . But for each $x' \in F'$ we have $x'(z_\alpha) \rightarrow \int_G f d(x' m)$. Thus

$$x'(z) = \int_G f d(x' m) \text{ which shows that } \int_G f dm = z. \text{ Define } T: C_{rc} \rightarrow F,$$

$$T(f) = \int_X f dm. \text{ Then } T \text{ is } u\text{-continuous and weakly compact. For}$$

the continuity, let $q \in J$. Choose $p \in I$ such that $\|m\|_{p,q} < \infty$. If

$$x' \in B_q^0 \text{ and } \|f\|_p \leq 1, \text{ we have } |x'(T(f))| = \left| \int_X f d(x' m) \right| \leq \|x' m\|_p \leq$$

$$\|m\|_{p,q}. \text{ It follows that } \|T\|_{p,q} \leq \|m\|_{p,q} \text{ and the continuity of } T \text{ is}$$

established. To prove the weak compactness consider an arbitrary

bounded set A in C_{rc} and let S denote the convex circled hull of $\cup \{f(X): f \in A\}$. Then S is bounded in E . Let $W = V_{m,S}$. Clearly W

is convex and circled. By hypothesis W is also weakly relatively compact. It follows that the polar W^0 of W in F' is a $m(F', F)$

neighborhood of zero. We will show that $T'(W^0) \subset A^0$. Let $x' \in W^0$

and $f \in A$. If G_1, \dots, G_n is a B -partition of X and $x_i \in G_i$, then

$$|x'(\sum_{i=1}^n m(G_i) f(x_i))| \leq 1. \text{ This implies that } |x'(f dm)| \leq 1. \text{ Thus}$$

$$|\langle T'x', f \rangle| = |\langle x', T(f) \rangle| \leq 1 \text{ which proves that } T'x' \in A^0. \text{ Now the}$$

result follows Lemma 1.

By the preceding theorem, given a continuous weakly compact operator T from C_{rc} into F there exists $m \in M(B, \mathcal{L}(E, F))$ which represents T . Since the operator $\hat{T}: C(\beta X, E) \rightarrow F, \hat{T}(\hat{f}) = T(f)$, is also

weakly compact and since the dual of $C(\beta X, E)$ (with the uniform topology) is $M_c(B_0(\beta X), E')$ we can find, using an argument analogous to that of Theorem 2, an $\hat{m} \in M_c(B_0(\beta X), \mathcal{L}(E, F))$ representing \hat{m} . The next theorem gives necessary and sufficient conditions on m and \hat{m} so that T is β'_1 continuous.

THEOREM 4. *Let T be a u -continuous and weakly compact operator from C_{rc} into F and let m and \hat{m} be as above. The following are equivalent:*

- (1) T is β'_1 continuous
- (2) Given $q \in J$ there exists p in I with $\|T\|_{p,q} < \infty$ such that $m_{p,q}(Z_n) \rightarrow 0$ whenever $\{Z_n\}$ is a sequence of zero sets decreasing to the empty set.
- (3) Given $q \in J$ there exists $p \in I$ with $\|T\|_{p,q} < \infty$ such that for each Z in Ω_1 we have $\inf\{\hat{m}_{p,q}(V) : V \text{ cozero set, } V \supset Z\} = 0$.

Proof. (1 \Rightarrow 3). Since T is β'_1 -continuous there exists $p \in I$ such that $T^{-1}(B_q)$ is a $\beta_{1,p}$ neighborhood of zero. Let now Z be in Ω_1 . Then there exists $g \in C^b(X)$ with $\hat{g}(Z) = 0$ such that $W = \{f \in C_{rc} : \|gf\|_p \leq 1\} \subset T^{-1}(B_q)$. Let $\epsilon > 0$ be given and set $V = \{x \in \beta X : |\hat{g}(x)| < \epsilon\}$. Then V is a cozero set containing Z . For a given $\delta > 0$ there exist $x' \in B_q^0$, a $B_0(\beta X)$ partition G_1, \dots, G_N of V and s_i in E with $p(s_i) \leq 1$ such that $|\sum x' \hat{m}(G_i) s_i| > \hat{m}_{p,q}(V) - \delta$. Next we choose compact sets $Z_i \subset G_i$ and pairwise disjoint open sets 0_i with $Z_i \subset 0_i \subset V$ such that $|\sum x' \hat{m}(G_i) s_i - \sum x' \hat{m}(Z_i) s_i| < \delta$ and $\sum (x' \hat{m})_p(0_i - Z_i) < \delta$. For each $i, 1 \leq i \leq n$, we pick $h_i \in C^b(X)$ with $0 \leq h_i \leq 1, \hat{h}_i = 1$ on Z_i and $\hat{h}_i = 0$ in the complement of 0_i in βX . Set $h = \sum h_i s_i$. Then $1/\epsilon h \in W$ and so $q(Th) \leq \epsilon$. Thus

$$\begin{aligned} \hat{m}_{p,q}(V) < \delta + \sum |x' \hat{m}(G_i) s_i| &\leq \delta + \delta + \sum \left| \int_{0_i - Z_i} \hat{h}_i s_i d(x' \hat{m}) \right| \\ &+ |x'(T(h))| \leq 3\delta + \epsilon. \end{aligned}$$

Since $\delta > 0$ was arbitrary we conclude that $\hat{m}_{p,q}(V) \leq \epsilon$. (3 \Rightarrow 2). Let $x' \in F'$. If $x' \in MB_q^0$ for some $q \in J$ and if $p \in I$ is as in (2), then from the fact that $(x' \hat{m})_p(Z) = 0$ for all Z in Ω_1 and from $\int_X f d(x' m) = \int_{\beta X} \hat{f} d(x' \hat{m})$, which holds for all f in C_{rc} , it follows that $x' m$ is σ -additive and hence $(x' \hat{m})_p(A) = (x' m)_p(A \cap X)$ for each A in $B(\beta X)$. Let now $\{Z_n\}$ be a sequence of zero sets in X which decreases to the empty set. For each n there exists a zero set F_n in βX such that $F_n \cap X = Z_n$. Let $\epsilon > 0$ be given. By (3) there exists a cozero set V in

βX containing $\cap F_n$ such that $\hat{m}_{p,q}(V) < \epsilon$. Since $(\cap F_n) \cap (\beta x - V) = \emptyset$ there exists N such that $F_1 \cap \dots \cap F_N \subset V$.

Now it follows that for $n \geq N$ we have

$$m_{p,q}(Z_n) \leq m_{p,q}(Z_N) = \hat{m}_{p,q}(F_1 \cap \dots \cap F_N) < \epsilon.$$

(2 \Rightarrow 1). Let $q \in J$ and choose $p \in I$ satisfying (2). For $x' \in B_q^0$ and $Z_n \downarrow \emptyset$ we have $(x'm)_p(Z_n) \leq m_{p,q}(Z_n) \rightarrow 0$ which implies that $x'm$ is σ -additive and so $(x'\hat{m})_p(A) = (x'm)_p(A \cap X)$ for each A in $B(\beta X)$. Let Z be in Ω_1 . There exists $h \geq 0$ in C^b such that $Z = \{x \in \beta X : \hat{h}(x) = 0\}$. For each n set $F_n = \{x \in \beta X : \hat{h}(x) \leq 1/n\}$. Then $Z_n = F_n \cap X$ is a zero set in X and $Z_n \downarrow \emptyset$. Given $r > 0$ there exists n such that $m_{p,q}(Z_n) < 1/2r$. Choose $g \in C^b$, $0 \leq g \leq 1$ with $\hat{g} = 0$ on Z and $\hat{g} = 1$ on the complement of V in βX , where $V = \{x \in \beta X : \hat{h}(x) < 1/n\}$. Let now $f \in C_{rc}$ with $\|f\|_p \leq r$ and $\|fg\|_p \leq \delta = 1/2\|m\|_{p,q}$. If

$$\begin{aligned} x' \in B_q^0, |x' \int f dm| &= |x' \int \hat{f} d\hat{m}| \leq \left| x' \int_V \hat{f} dm \right| \\ &+ \left| x' \int_{\beta X - V} \hat{g} \hat{f} d\hat{m} \right| \leq r \cdot 1/2r + \delta \|m\|_{p,q} \leq 1. \end{aligned}$$

This shows that $q(\int f dm) \leq 1$. Thus $\{f \in C_{rc} : \|f\|_p \leq r, \|fg\|_p \leq \delta\} \subset T^{-1}(B_q)$ and $s_0 T^{-1}(B_q)$ is a $\beta_{p,Z}$ neighborhood of zero. Since this is true for all Z in Ω_1 it follows that $T^{-1}(B_q)$ is a $\beta_{1,p}$ neighborhood of zero which proves that T is β'_1 continuous.

We have an analogous theorem for β' with a similar proof.

THEOREM 5. *Let T, m and \hat{m} be as in Theorem 3. The following are equivalent:*

- (1) T is β' -continuous
- (2) Given $g \in J$ there exists $p \in I, \|T\|_{p,q} < \infty$ such that for each G in Ω we have $\inf\{\hat{m}_{p,q}(V) : V \text{ cozero set, } G \subset V\} = 0$.
- (3) Given $g \in J$ there exists $p \in I$ with $\|T\|_{p,q} < \infty$ that $m_{p,q}(Z_\alpha) \rightarrow 0$ for each net $\{Z_\alpha\}$ of zero sets in X which decreases to the empty set.

THEOREM 6. *Suppose T is a linear operator from C_{rc} into F which is β_1 -continuous and that every weakly closed bounded subset of F is weakly sequentially complete. Then there exists $m \in M(B, \mathcal{L}(E, F))$, with respect to which each f in C_{rc} is integrable, such that $T(f) = \int f dm$ for all f in C_{rc} . Moreover, if T is β'_1 continuous, given $q \in J$ there exists $p \in I$ with $\|m\|_{p,q} < \infty$ such that $m_{p,q}(Z_n) \rightarrow 0$ whenever $\{Z_n\}$ is a sequence of zero sets which decreases to the empty set.*

Proof. Since T is β_1 -continuous, T' maps F' into the space $M_\sigma(B, E') = (C_{rc}, \beta_1)'$. Let Z be a zero set in X . There exists $g \in C^b$ such that $Z = \{x : g(x) = 0\}$. For each n let

$$V_n = \{x \in X : |g(x)| < 1/n\}.$$

Choose f_n in $C^b, 0 \leq f_n \leq 1$ with $f_n = 1$ on Z and $f_n = 0$ on $X - V_n$. Then $f_n \rightarrow \chi_Z$ pointwise. An arbitrary element of $B(X)$ can be written as a finite disjoint union of sets of the form $Z - F$ where $F \subset Z$ and F, Z are zero sets. It follows that for G in $B(X)$ there exists a bounded sequence $\{f_n\}$ in C^b which converges pointwise to χ_G . For μ in $M_G(B, E')$ and $s \in E$ we have $\langle \mu, f_n s \rangle = \int f_n d(\mu s) \rightarrow \int \chi_G d(\mu s) = \langle \mu, \chi_G s \rangle$. Thus $f_n s \rightarrow \chi_G s$ in the $\sigma((C_{rc}, \beta_1)'', M_\sigma(B, E'))$ topology and hence $T''(f_n s) \rightarrow T''(\chi_G s)$ in the $\sigma(F'', F')$ sense. But $T''(f_n s) = T(f_n s)$ and the set $\{T(f_n s) : n = 1, 2, \dots\}$ is $\sigma(F, F')$ bounded. Also the sequence $\{T''(f_n s)\}$ is weakly Cauchy. By hypothesis there exists $a \in F$ that $T''(f_n s) \rightarrow a$ in the $\sigma(F, F')$ topology. This implies that $T''(\chi_G s) = a \in F$. Define $m(G)s = T''(\chi_G s)$. It is easy to see that $m \in M(B, \mathcal{L}(E, F))$ and that $T(f) = \int f dm$ for all f in C_{rc} . Assume next that T is β_1 -continuous. Let $\hat{T} : C(\beta X, E) \rightarrow F, \hat{T}(f) = T(f)$. As in the case of T we can find $\bar{m} \in M(B(\beta X), \mathcal{L}(E, F))$ such that $\hat{T}(f) = \int f d\bar{m}$ for all f in C_{rc} . Now to complete the proof we use an argument similar to that of Theorem 4.

If $m \in M_\sigma(Ba, \mathcal{L}(E, F))$, then the restriction of m to B is in $M_\sigma(B, \mathcal{L}(E, F))$. The following result is a partial converse.

THEOREM 7. *Let $m \in M_\sigma(B, \mathcal{L}(E, F))$ be such that for any $s \in E$ the set $(ms)(B)$ is weakly relatively compact in F . Then there exists a unique \bar{m} in $M_\sigma(Ba, \mathcal{L}(E, F))$ whose restriction to B coincides with m . Moreover, if $\|m\|_{p,q} < \infty$, then $\|\bar{m}\|_{p,q} = \|m\|_{p,q}$.*

Proof. Let $G \in Ba$ and set $W = \{Z : Z \subset G, Z \text{ a zero set}\}$. If we order W by inclusion, it becomes a directed set. For $s \in E, \{m(Z)s : Z \in W\}$ is a net in F . By hypothesis there exists a subnet which converges weakly to some a in F . For $x' \in F', x'm$ is σ -additive and thus has a unique extension to a member $\mu_{x'}$ of $M_\sigma(Ba, E')$. Moreover $x'm(Z)s \rightarrow \mu_{x'}(G)s$. Thus $x'(a) = \mu_{x'}(G)s$. Define $\bar{m}(G)s = x'(a)$. Then $x'\bar{m} = \mu_{x'} \in M_\sigma(Ba, E')$. Furthermore $\bar{m}(G) \in \mathcal{L}(E, F)$. Indeed if $\|m\|_{p,q} < \infty$, then for $x' \in B_q^0$ we have $|x'\bar{m}(G)s| = |\mu_{x'}(G)s| \leq p(s)\|\mu_{x'}\|_p = p(s)\|x'm\|_p \leq p(s)\|m\|_{p,q}$. Thus $q(\bar{m}(G)s) \leq p(s)\|m\|_{p,q}$ which proves that $\bar{m}(G) \in \mathcal{L}(E, F)$. Also $\|x'\bar{m}\|_p = \|\mu_{x'}\|_p = \|x'm\|_p$ implies that $\|\bar{m}\|_{p,q} = \|m\|_{p,q}$. Finally suppose λ is another extension. Then for each x' in F' both $x'\lambda$ and $x'\bar{m}$ are extensions of $x'm$ and hence they are equal. This implies that $\lambda = \bar{m}$.

REFERENCES

1. R. G. Bartle, N. Dunford, and J. Schwartz, *Weak compactness and vector measures*, Canad. J. Math., **7** (1955), 289–305.
2. A. Grothendieck, *Sur les applications lineaires faiblement compactes d'espaces du type $C(K)$* , Canad. J. Math., **V**, **2** (1953), 129–173.
3. A. Katsaras, *Vector-valued measures on topological spaces*, Doctoral Dissertation, Southern Illinois University (1973).
4. ———, *Spaces of vector-valued measures*, Trans. Amer. Math. Soc., (to appear).
5. D. R. Lewis, *Integration with respect to vector measures*, Pacific J. Math., **33** (1970).
6. B. J. Pettis, *On Integration in vector spaces*, Trans. Amer. Math. Soc., **44** (1938), 277–304.
7. H. H. Schaefer, *Topological Vector Spaces*, (Macmillan, New York (1966)).
8. V. Varadarajan, *Measures on topological spaces*, Amer. Math. Soc. Transl., (2), **48** (1965), 161–228.

Received December 20, 1973 and in revised form April 10, 1974.

UNIVERSIDADE ESTADUAL DE CAMPINAS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$ 72.00 a year (6 Vols., 12 issues). Special rate: \$ 36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1975 Pacific Journal of Mathematics
All Rights Reserved

Ralph Alexander, <i>Generalized sums of distances</i>	297
Zvi Arad and George Isaac Glauberan, <i>A characteristic subgroup of a group of odd order</i>	305
B. Aupetit, <i>Continuité du spectre dans les algèbres de Banach avec involution</i>	321
Roger W. Barnard and John Lawson Lewis, <i>Coefficient bounds for some classes of starlike functions</i>	325
Roger W. Barnard and John Lawson Lewis, <i>Subordination theorems for some classes of starlike functions</i>	333
Ladislav Bican, <i>Preradicals and injectivity</i>	367
James Donnell Buckholtz and Ken Shaw, <i>Series expansions of analytic functions. II</i>	373
Richard D. Carmichael and E. O. Milton, <i>Distributional boundary values in the dual spaces of spaces of type \mathcal{S}</i>	385
Edwin Duda, <i>Weak-unicoherence</i>	423
Albert Edrei, <i>The Padé table of functions having a finite number of essential singularities</i>	429
Joel N. Franklin and Solomon Wolf Golomb, <i>A function-theoretic approach to the study of nonlinear recurring sequences</i>	455
George Isaac Glauberan, <i>On Burnside's other $p^a q^b$ theorem</i>	469
Arthur D. Grainger, <i>Invariant subspaces of compact operators on topological vector spaces</i>	477
Jon Craig Helton, <i>Mutual existence of sum and product integrals</i>	495
Franklin Takashi Iha, <i>On boundary functionals and operators with finite-dimensional null spaces</i>	517
Gerald J. Janusz, <i>Generators for the Schur group of local and global number fields</i>	525
A. Katsaras and Dar-Biau Liu, <i>Integral representations of weakly compact operators</i>	547
W. J. Kim, <i>On the first and the second conjugate points</i>	557
Charles Philip Lanski, <i>Regularity and quotients in rings with involution</i>	565
Ewing L. Lusk, <i>An obstruction to extending isotopies of piecewise linear manifolds</i>	575
Saburou Saitoh, <i>On some completenesses of the Bergman kernel and the Rudin kernel</i>	581
Stephen Jeffrey Willson, <i>The converse to the Smith theorem for Z_p-homology spheres</i>	597