

Pacific Journal of Mathematics

**ON CAUCHY'S THEOREM FOR REAL ALGEBRAIC CURVES
WITH BOUNDARY**

NORMAN LARRABEE ALLING

ON CAUCHY'S THEOREM FOR REAL ALGEBRAIC CURVES WITH BOUNDARY

NORMAN L. ALLING

On a real algebraic curve with a nonempty boundary, one must orient the several boundary components in order to pose the question considered in Cauchy's theorem for analytic differentials. It is proved that the conclusion of Cauchy's theorem is true, in this context, if and only if the orientation in question is induced by an orientation of the interior of the curve.

Let \mathfrak{Y} be a real algebraic curve (i.e., a compact Klein surface [3, 4]), whose boundary ∂Y has r components, where $r > 0$. Let g be the algebraic genus of \mathfrak{Y} : i.e., the genus of the field E of meromorphic "functions" on \mathfrak{Y} that are real valued on ∂Y ; then g is the first Betti number of Y , the underlying space of \mathfrak{Y} , and the Euler characteristic χ of Y is $1 - g$ [4, 2]. Y is—of course—characterised topologically by knowing g, r and whether or not Y is orientable. In [2] the author investigated some sheaves that arise from analytic problems on \mathfrak{Y} , whose cohomology groups reflect the orientability of Y ; however these sheaves and groups seemed rather remote from analytic function theory on \mathfrak{Y} . This paper is an outgrowth of the search for a simple analytic question which could be posed on \mathfrak{Y} , whose answer would reflect the orientability of Y . What analytic question on \mathfrak{Y} is, after all, more basic than Cauchy's theorem?

In order to pose the question considered in Cauchy's Theorem on \mathfrak{Y} , we must orientate the r -components of ∂Y ; there are 2^r ways to do this. If Y is orientable, then two of these 2^r orientations are engendered by the two possible orientations of Y ; these will be called *indigenous orientations* of ∂Y . If Y is nonorientable, then ∂Y has no indigenous orientations. Let \mathcal{O} be an orientation of ∂Y .

Next we must have a space of analytic differentials on \mathfrak{Y} to integrate along ∂Y , as oriented by \mathcal{O} . A space $\Omega_{\mathfrak{Y}}$ of analytic "differentials" on \mathfrak{Y} was defined in [4] which is the natural generalization of the space of Schottky differentials on a bordered Riemann surface, in that they are real on ∂Y . (The space of meromorphic differentials on \mathfrak{Y} [4, 1.10] is also very natural from the point of view of the algebraic geometry of E .) Even though $\omega \in \Omega_{\mathfrak{Y}}$ is called a "differential" its integral along an oriented Jordan curve, or arc Γ , need not have an invariant meaning! If $\Gamma \subset \partial Y$ then it does, and $\int_{\Gamma} \omega \in \mathbb{R}$. The real

part of $\int_r \omega$ always has an invariant meaning. If Γ is contained in an orientable tubular neighborhood, then $\int_r \omega$ is always invariantly defined, up to complex conjugation. The real dimension of $\Omega_{\mathfrak{Y}}$ is g [2]. (The reason it is so small is that the condition that ω be real on ∂Y allows it to be extended to the complex double.)

$\lambda_{\mathcal{O}}: \omega \in \Omega_{\mathfrak{Y}} \mapsto \int_{\partial Y, \mathcal{O}} \omega \in \mathbf{R}$ is a well defined \mathbf{R} -linear functional on $\Omega_{\mathfrak{Y}}$ whose image is either of dimension 0 or 1. Accordingly we will say that *the conclusion of Cauchy's theorem holds on \mathfrak{Y} for \mathcal{O}* , or *does not hold on \mathfrak{Y} for \mathcal{O}* .

THEOREM 1. *The conclusion of Cauchy's theorem holds on \mathfrak{Y} for an orientation \mathcal{O} of $\partial Y (\neq \emptyset)$ if and only if \mathcal{O} is an indigenous orientation of ∂Y (i.e., one induced by an orientation of Y). Thus Y is orientable (resp. nonorientable) if and only if there exists 2 (resp. 0) orientations \mathcal{O} of ∂Y for which the conclusion of Cauchy's theorem holds on \mathfrak{Y} .*

Proof. Assume first that \mathcal{O} is an indigenous orientation of ∂Y ; then, by definition, \mathcal{O} is induced by an orientation of Y . In the dianalytic structure on Y that gives \mathfrak{Y} we may choose an analytic structure and thus consider the bordered Riemann surface structure \mathfrak{Y}_1 on Y that engenders \mathcal{O} on ∂Y [4]. Then $\Omega_{\mathfrak{Y}}$ becomes the space of Schottky differentials on \mathfrak{Y}_1 . In this context the Cauchy theorem is known to hold. (This can be shown directly by triangulating Y and using the Cauchy theorem in C .) Assume, henceforth that \mathcal{O} is nonindigenous.

It is well known (see e.g., [6]), that a topological model of Y can be built up from a closed unit disc D , by adjoining various strips and handles to it. Adopting a different construction suited to our purposes, first adjoin to D $r - 1$ untwisted strips—glued to the boundary of D —to form D' so that $\partial D'$ has r components. Let D' be imbedded in C (in some way), let $\Gamma_1, \dots, \Gamma_{r-1}$ be the components of $\partial D'$ that bound bounded components of $C - D'$, and let them be positively oriented relative to D' (as oriented by C). The Euler characteristic of D' , $\chi(D')$, is $2 - r$, which is—necessarily—not smaller than $\chi(Y)$. Next choose the largest integer h such that $2 - r - 2h \geq \chi(Y)$, in the orientable case or $2 - r - 2h \geq \chi(Y) + 1$ in the nonorientable. Adjoin h handles to D' —by removing $2h$ open discs from D' , whose boundaries do not meet $\partial D'$, and attaching h handles (each on the same side of D'), to some choice of h -pairs of these circular boundaries—to form D_0 . If Y is orientable, then Y and D_0 are homeomorphic. In this event let Y and D_0 be

identified. If Y is nonorientable, let D_1 be formed by adjoining a half twisted strip to D_0 ; then $\chi(D_1) = 2 - r - 2h - 1$, which is either $\chi(Y)$ or is $\chi(Y) + 1$. In the first case Y and D_1 are homeomorphic, and are to be identified. In the second let one more half twisted strip be adjoined to D_1 to form D_2 , a space that is homeomorphic to Y ; then let D_2 and Y be identified. In general, $\chi(Y) = 2 - r - 2h - m$, where $m = 0, 1$, or 2 , and Y and D_m are identified. Let Γ_r be $\partial Y - (\Gamma_1 \cup \dots \cup \Gamma_{r-1})$, and let it be oriented in such a way that Γ_j and Γ_r have opposing orientations on the untwisted strip adjoined to D to form D' , which gives rise to Γ_j , for each $1 \leq j < r$.

The main analytic technique we will use, that of doubling \mathfrak{Y} , goes back—essentially to Schottky and Schwarz—and explicitly, in this context, to Klein. Let (\mathfrak{X}, τ, p) be the complex double of \mathfrak{Y} [4, 1.6]: i.e., \mathfrak{X} is a compact Riemann surface (without boundary), τ is an anti-analytic involution of \mathfrak{X} , and p is an analytic map of \mathfrak{X} onto \mathfrak{Y} (i.e., a morphism [4, 1.4]). $p^{-1}(y)$ has one (resp. two) points in it, for $y \in Y$, if and only if $y \in \partial Y$ (resp. $y \in Y - \partial Y$). For $1 \leq j \leq r$, let Δ_j be the pullback of Γ_j to X , endowed with the orientation induced on it by the orientation of Γ_j ; thus Δ_j is an oriented Jordan curve in X . For $1 \leq j < r$, let $a_j \equiv \Delta_j$. It is easy to see that these, regarded as elements in $H_1(X, \mathbb{Z})$, are part of the usual a -paths (see e.g., [7, Chapt. 10] for details), which arise from doubling the $r - 1$ untwisted strips that were adjoined to D to form D' . τ induces an involution σ on $\Omega_{\mathfrak{X}}$, the \mathbb{C} -space of analytic differentials on \mathfrak{X} , which is \mathbb{R} -linear, such that $\sigma(i\omega) = -i\sigma(\omega)$, for each $\omega \in \Omega_{\mathfrak{X}}$. $\Omega_{\mathfrak{X}}$ is then the direct sum of the \mathbb{R} -space, $\Omega_{\mathfrak{X},s}$, of symmetric elements of $\Omega_{\mathfrak{X}}$, and the \mathbb{R} -space $\Omega_{\mathfrak{X},a}$ of anti-symmetric elements of $\Omega_{\mathfrak{X}}$. Further $\Omega_{\mathfrak{X},a} = i\Omega_{\mathfrak{X},s}$; thus the real dimensions of $\Omega_{\mathfrak{X},a}$ and $\Omega_{\mathfrak{X},s}$ are the same, namely g , the genus of \mathfrak{X} —which is also the algebraic genus of \mathfrak{Y} . Given $\omega \in \Omega_{\mathfrak{X}}$, let $\omega = \rho + \zeta$ where ρ is symmetric and ζ is anti-symmetric. This convention will hold throughout the paper. Further $\Omega_{\mathfrak{Y}}$ can be naturally identified with $\Omega_{\mathfrak{X},s}$ (see [4], [1], and [2] for more details).

LEMMA. *Let a be an oriented Jordan curve (arc) in X such that $\tau(a) = a$, and assume that $\int_a \omega \equiv t$ is real; then $\int_a \rho = t$ and $\int_a \zeta = 0$.*

Proof. Using [1, 3.1], we see that $\int_a \rho = \kappa \circ \int_{\tau(a)} \rho = \kappa \circ \int_a \rho$, so $\int_a \rho \in \mathbb{R}$. $\zeta = i\eta$ for some $\eta \in \Omega_{\mathfrak{X},s}$, so $\int_a \zeta = i \int_a \eta \in i\mathbb{R}$, proving the lemma.

Returning now to the proof of Theorem 1, first let us treat the (trivial) case in which Y is orientable, or equivalently in which $m =$

0. Then Y and D_0 are identified, and $\Gamma_1 + \dots + \Gamma_r = 0$ in $H_1(Y, Z)$. The orientation \mathcal{O} , of ∂Y , can be given by choosing $e_j \in \{\pm 1\}$, $1 \leq j \leq r$, and by assigning each e_j to Γ_j . Since \mathcal{O} is nonindigenous, not all the e_j 's are alike. $\{a_1, \dots, a_{r-1}\}$ is contained in a basis $B_0 \equiv \{a_1, b_1, \dots, a_g, b_g\}$ of $H_1(X, Z)$ such that

$$\text{a) } a_1 \cdot b_1 \cdot a_1^{-1} \cdot b_1^{-1} \cdot \dots \cdot a_g \cdot b_g \cdot a_g^{-1} \cdot b_g^{-1} = 1, \text{ in } \pi_1(X).$$

Let $\{\omega_1, \dots, \omega_g\}$ be a basis of Ω_x over C such that $\int_{a_j} \omega_k = \delta_{jk}$, $1 \leq j, k \leq g$. (See e.g., [7, Chapt. 10] for details.) As noted above there exists k , $1 \leq k < r$, such that $e_k \neq e_r$. Using, among other things, the lemma, we find that

$$\int_{\partial Y, \mathcal{O}} \rho_k = \sum_{j=1}^r e_j \int_{\Delta_j} \rho_k = e_k \int_{\Delta_k} \rho_k + e_r \int_{\Delta_r} \rho_k.$$

Since $\Gamma_1 + \dots + \Gamma_r = 0$ in $H_1(Y, Z)$, $\Delta_r = -\Delta_1 - \dots - \Delta_{r-1}$ in $H_1(X, Z)$; thus $e_r \int_{\Delta_r} \rho_k = -e_r \int_{\Delta_k} \rho_k$, by Cauchy's theorem, and thus is $-e_r$, and so $\int_{\partial Y, \mathcal{O}} \rho_k = e_k - e_r \neq 0$, disposing of the proof, if Y is orientable. Before going on to the nonorientable cases, let us consider a useful example.

EXAMPLE. Let $\alpha \equiv 1/2 + bi$, where $b > 0$, and consider $G \equiv Z \oplus \alpha Z$ in C . Let G act as a set of conformal maps of C , by translation. Note that G is invariant under κ , where κ is complex conjugation. Let $\mathfrak{X} \equiv C/G$, and let τ be the anti-analytic involution on \mathfrak{X} induced by κ . The parallelogram whose vertices are $0, \bar{\alpha}, 1$, and α can be taken as a fundamental domain for \mathfrak{X} . The interval $[0, 1]$ is the set of fixed points of this domain under the action of κ . Let $\mathfrak{Y} \equiv \mathfrak{X}/\{1, \tau\}$; then \mathfrak{Y} is a Möbius strip, the image of $[0, 1]$ in Y being its boundary. The isosceles triangle, whose vertices are $0, 1$ and α , may be taken as the fundamental domain for \mathfrak{Y} . Given β in the straight line segment $[0, \alpha]$, it will be identified—when passing to \mathfrak{Y} —with $\bar{\beta} + \alpha$ in $[\alpha, 1]$. dz induces a basis $\{dz\}$ of $\Omega_{\mathfrak{Y}}$ over R . Since $\int_0^1 dz = 1$, we see that the conclusion of Cauchy's theorem is never true for \mathfrak{Y} . (It is also not hard to show that all dianalytic Möbius strips occur in this way.)

Returning again to the proof of Theorem 1, note that the triangle, described in the above example and identified as indicated, is a Möbius strip. Assume now that Y is nonorientable: i.e., $m > 0$. We can modify the construction of D_m from D_0 as follows. Let one end of a strip be glued to $[1/3, 2/3]$ in the Möbius strip above. The resulting space will be referred to as a *Christmas tree*. Glue the other end of the strip that forms the *trunk* of the Christmas tree to the edge

of D_0 to form D_1 . Repeat the procedure on D_1 to form D_2 . Let A_r and A_{r+m-1} be the boundaries of the Möbius strip—in the form given in the example—before the trunk is glued on, and then glued to D_0 , and D_{m-1} . Let them be oriented to agree with the orientation of Γ_r . In doubling \mathfrak{Y} to form \mathfrak{X} , the Christmas tree doubles to a torus with a tube running from a hole in it, back to the rest of X . A_j lifts to a (nonunique) path A_j in X , $r \leq j \leq r+m-1$. Let these paths be oriented by the paths onto which they map. Let $a_r \equiv A_r$ and let $a_{r+m-1} \equiv A_{r+m-1}$. $\{a_1, \dots, a_r, a_{r+m-1}\}$ is contained in a basis $B_m \equiv \{a_1, b_1, \dots, a_g, b_g\}$ of $H_1(X, Z)$ that satisfies condition a) above. Thus there exists a basis $\{\omega_1, \dots, \omega_g\}$ of $\Omega_{\mathfrak{X}}$ over C such that $\int_{a_j} \omega_k = \delta_{jk}$, $1 \leq j, k \leq g$. (See e.g., [7, Chapt. 10] again.) Let ρ_k be the symmetric component of ω_k , and let $\rho \equiv \sum_{k=1}^m \rho_{r+k-1}$; then $\rho \in \Omega_{\mathfrak{Y}}$. There exist $e_j \in \{\pm 1\}$, $1 \leq j \leq r$, such that $\int_{\partial Y, \mathcal{O}} \rho = \sum_{j=1}^r e_j \int_{\Gamma_j} \rho$. Since—for $1 \leq j < r - \tau(a_j) = a_j$, and since $\int_{a_j} \omega_{r+k-1} = 0$, $k = 1$ or 2 , for such j 's, we may apply the lemma and conclude that $\int_{a_j} \rho = 0$; thus $\sum_{j=1}^r e_j \int_{\Gamma_j} \rho = e_r \int_{\Gamma_r} \rho$. Assume now that $m = 1$. Since we may invoke Cauchy's theorem on an orientable sub-domain of \mathfrak{Y} , we find that $\int_{\Gamma_r} \rho = \pm \int_{a_r} \rho$. Let a_r be re-oriented so that positive sign above holds. Since $\int_{a_r} \omega_r = 1$, we may reason as we did above. Since $\int_{\Gamma_r} \omega_r = \int_{a_r} \omega_r = 1$, and since $\tau(\Gamma_r) = \Gamma_r$, we may apply the lemma and conclude that $\int_{\Gamma_r} \rho = 1$. Thus $\int_{\partial Y, \mathcal{O}} \rho = e_r \neq 0$. Assume lastly that $m = 2$. Reasoning as above $\int_{\Gamma_r} \omega_r + \omega_{r+1} = \int_{\pm a_r \pm a_{r+1}} \omega_r + \omega_{r+1}$, the signs being independent, one of the other. If necessary, re-orient a_r or a_{r+1} so that the plus sign holds twice above; thus $\int_{\Gamma_r} \omega_r + \omega_{r+1} = 2$. Since $\tau(\Gamma_r) = \Gamma_r$, we may apply the lemma and conclude that $\int_{\partial Y, \mathcal{O}} \rho = e_r \int_{\Gamma_r} \rho = 2e_r \neq 0$, proving the theorem.

Greenleaf and Read considered a related question in [5]. Given an orientation \mathcal{O} of ∂Y they defined the notion of an analytic differential ρ as being *positive at y* (in ∂Y), *relative to \mathcal{O}* as follows: if given $f \in E$ —a local uniformizer at y —that is increasing near y , relative to \mathcal{O} then $0 < h(y)$, where $\rho = hdf$ for a (unique) $h \in E$. (This definition is independent of the choice of f , as may easily be seen.) ρ is said to be *positive relative to \mathcal{O}* if it is positive at y , relative to \mathcal{O} , for each $y \in \partial Y$. Greenleaf and Read proved that if \mathcal{O} is indigenous, then no positive analytic differentials exist in $\Omega_{\mathfrak{Y}}$, that if \mathcal{O} is non-indigenous and \mathfrak{Y} is *elliptic* or *hyperelliptic* then positive analytic differentials always exist; and then they went on to conjecture that the condition that \mathfrak{Y} be elliptic or hyperelliptic can

be dropped while preserving the conclusion above. If the Greenleaf-Read conjecture is correct, then it would imply our theorem; thus our theorem may lend additional credence to their conjecture.

Having set up this machinery, let us use it to draw some additional conclusions. As noted before, the real dimension of $\Omega_{\mathfrak{y}}$ is g and $g = r - 1 + m + 2h$. Given $\rho \in \Omega_{\mathfrak{y}}$, let

$$\varphi_1(\rho) \equiv \int_{\Gamma_1} \rho, \dots, \varphi_{r-1}(\rho) \equiv \int_{\Gamma_{r-1}} \rho.$$

Clearly these maps are real valued. If $m \geq 1$, let $\varphi_r(\rho) \equiv \int_{A_r} \rho$. If $m = 2$, let $\varphi_{r+1}(\rho) \equiv \int_{A_{r+1}} \rho$. Let $r \leq k \leq r + m - 1$. By [1, 3.1], $\int_{\tau(\Delta_k)} \rho = \kappa(\phi_k(\rho))$. Since Δ_k and $\tau(\Delta_k)$ are homologous, $\int_{\tau(\Delta_k)} \rho = \phi_k(\rho)$, proving that $\phi_k(\rho)$ is real. $\phi_k(\rho)$ can also be computed by integrating allong Λ_k . Note that Λ_k lifts to Δ_k and to $\tau(\Delta_k)$, and that the integral of ρ allong each of these paths is the same. Let Θ_j be an oriented α -path about the j^{th} handle adjoined to D' to form D_0 , for $1 \leq j \leq h$.

For each such path we may choose an analytic structure in a tubular neighborhood; then integration of ρ along each such is well defined. As remarked before the real parts of these integrals are a priori well defined, whereas the imaginary parts are a priori well defined only up to sign. If $h > 0$, let $\varphi_g(\rho) \equiv \text{Re} \int_{\Theta_1} \rho$, $\varphi_{g-1}(\rho) \equiv \pm \text{Im} \int_{\Theta_1} \rho$, \dots , $\varphi_{g-2h+2}(\rho) \equiv \text{Re} \int_{\Theta_h} \rho$, and $\varphi_{g-2h+1}(\rho) \equiv \pm \text{Im} \int_{\Theta_h} \rho$, the signs above being independent of one another.

THEOREM 2. *Given $\rho \in \Omega_{\mathfrak{y}}$, $\rho = 0$ if and only if $\varphi_1(\rho) = \dots = \varphi_g(\rho) = 0$.*

Proof. Since each of the φ_j 's is an R -linear functional, $\rho = 0$ implies that each $\varphi_j(\rho) = 0$. Conversely, let $\rho \in \Omega_{\mathfrak{y}}$ such that $\varphi_j(\rho) = 0$, for each j , $1 \leq j \leq g$. As noted in the proof of Theorem 1, Γ_j lifts to $\alpha_j \in H_1(X, Z)$ for $1 \leq j \leq r - 1$. If $m > 0$, Λ_r lifts to α_r , and if $m = 2$, Λ_{r+1} lifts to α_{r+1} . For $1 \leq j \leq h$, Θ_j lifts to two oriented α -paths on X , α_{g-2j+2} and α_{g-2j+1} , which are permuted by τ . Finally, $\{\alpha_1, \dots, \alpha_g\}$ is contained in a basis $B \equiv \{\alpha_1, b_1, \dots, \alpha_g, b_g\}$ of $H_1(X, Z)$ such that condition a) holds. By [1, 3.1], the integral of ρ about α_{g-2j+2} and about α_{g-2j+1} are complex conjugates of each other. Because of this symmetry, to know that ρ is zero it suffices to know that its periods with respect to $\alpha_1, \dots, \alpha_{r-1}, \dots, \alpha_{r-1+m}, \alpha_g, \alpha_{g-2}, \dots, \alpha_{g-2h+4}$, and α_{g-2h+2} are all zero: i.e., the periods of ρ with respect to $\Gamma_1, \dots, \Gamma_{r-1}, \dots, \Lambda_{r-1+m}, \Theta_1, \Theta_2, \dots, \Theta_{h-1}, \Theta_h$ are all zero; but this is implied by the condition that $\varphi_1(\rho) = \dots = \varphi_g(\rho) = 0$, proving the theorem.

COROLLARY. *There exists a unique basis $\{\rho_1, \dots, \rho_g\}$ in $\Omega_{\mathfrak{y}}$ of $\Omega_{\mathfrak{y}}$ over \mathbf{R} (resp. in $\Omega_{\mathfrak{y}}$ of $\Omega_{\mathfrak{x}}$ over \mathbf{C}), such that $\varphi_j(\rho_i) = \delta_{jk}$, for all $1 \leq j, k \leq g$.*

Bibliographic note. See also [8, 9] for related results on the period matrix of a symmetric Riemann surface.

REFERENCES

1. N. L. Alling, *Analytic and harmonic obstruction on nonorientable Klein surfaces*, Pacific J. Math., **36** (1971), 1-19.
2. ———, *Analytic geometry on real algebraic curves*, Math. Ann., **207** (1974), 23-46.
3. N. L. Alling and N. Greenleaf, *Klein surfaces and real algebraic function fields*, Bull. Amer. Math. Soc., **75** (1969), 869-872.
4. ———, *Foundations of the Theory of Klein Surfaces and Real Algebraic Function Fields*, Springer-Verlag Lecture Notes in Mathematics, vol. **219** (1971).
5. N. Greenleaf and W. Read, *Position holomorphic differentials on Klein surfaces*, Pacific J. Math., **32** (1970), 711-713.
6. W. S. Massey, *Algebraic Topology: An Introduction*, Harcourt, Brace and World, 1967.
7. G. Springer, *Introduction to Riemann Surfaces*, Addison-Wesley, 1957.
8. G. Weichold, *Über symmetrische Riemannsche Flächen und die Periodizitätsmoduln der zugehörigen Abelschen Normalintegrale erster Gattung*, Zeitschrift f. Math. u. Phys., **28** (1883), 321-351.
9. R. Zarrow, *Symmetric and quasi-symmetric Riemann surfaces*, unpublished dissertation, Cornell University, 1972.

Received November 16, 1973. The author wishes to thank Walter Read, the Referee, for his careful reading and many constructive criticisms which have been incorporated in this paper.

THE UNIVERSITY OF ROCHESTER

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 57, No. 2

February, 1975

Norman Larrabee Alling, <i>On Cauchy's theorem for real algebraic curves with boundary</i>	315
Daniel D. Anderson, <i>A remark on the lattice of ideals of a Prüfer domain</i>	323
Dennis Neal Barr and Peter D. Miletta, <i>A necessary and sufficient condition for uniqueness of solutions to two point boundary value problems</i>	325
Ladislav Beran, <i>On solvability of generalized orthomodular lattices</i>	331
L. Carlitz, <i>A three-term relation for some sums related to Dedekind sums</i>	339
Arthur Herbert Copeland, Jr. and Albert Oscar Shar, <i>Images and pre-images of localization maps</i>	349
G. G. Dandapat, John L. Hunsucker and Carl Pomerance, <i>Some new results on odd perfect numbers</i>	359
M. Edelstein and L. Keener, <i>Characterizations of infinite-dimensional and nonreflexive spaces</i>	365
Francis James Flanigan, <i>On Levi factors of derivation algebras and the radical embedding problem</i>	371
Harvey Friedman, <i>Provable equality in primitive recursive arithmetic with and without induction</i>	379
Joseph Braucher Fugate and Lee K. Mohler, <i>The fixed point property for tree-like continua with finitely many arc components</i>	393
John Norman Ginsburg and Victor Harold Saks, <i>Some applications of ultrafilters in topology</i>	403
Arjun K. Gupta, <i>Generalisation of a "square" functional equation</i>	419
Thomas Lee Hayden and Frank Jones Massey, <i>Nonlinear holomorphic semigroups</i>	423
V. Kannan and Thekkedath Thrivikraman, <i>Lattices of Hausdorff compactifications of a locally compact space</i>	441
J. E. Kerlin and Wilfred Dennis Pepe, <i>Norm decreasing homomorphisms between group algebras</i>	445
Young K. Kwon, <i>Behavior of Φ-bounded harmonic functions at the Wiener boundary</i>	453
Richard Arthur Levaro, <i>Projective quasi-coherent sheaves of modules</i>	457
Chung Lin, <i>Rearranging Fourier transforms on groups</i>	463
David Lowell Lovelady, <i>An asymptotic analysis of an odd order linear differential equation</i>	475
Jerry Malzan, <i>On groups with a single involution</i>	481
J. F. McClendon, <i>Metric families</i>	491
Carl Pomerance, <i>On multiply perfect numbers with a special property</i>	511
Mohan S. Putcha and Adil Mohamed Yaqub, <i>Polynomial constraints for finiteness of semisimple rings</i>	519
Calvin R. Putnam, <i>Hyponormal contractions and strong power convergence</i>	531
Douglas Conner Ravenel, <i>Multiplicative operations in BP^*BP</i>	539
Judith Roitman, <i>Attaining the spread at cardinals which are not strong limits</i>	545
Kazuyuki Saitô, <i>Groups of $*$-automorphisms and invariant maps of von Neumann algebras</i>	553
Brian Kirkwood Schmidt, <i>Homotopy invariance of contravariant functors acting on smooth manifolds</i>	559
Kenneth Barry Stolarsky, <i>The sum of the distances to N points on a sphere</i>	563
Mark Lawrence Teply, <i>Semiprime rings with the singular splitting property</i>	575
J. Pelham Thomas, <i>Maximal connected Hausdorff spaces</i>	581
Charles Thomas Tucker, II, <i>Concerning σ-homomorphisms of Riesz spaces</i>	585
Rangachari Venkataraman, <i>Compactness in abelian topological groups</i>	591
William Charles Waterhouse, <i>Basically bounded functors and flat sheaves</i>	597
David Westreich, <i>Bifurcation of operator equations with unbounded linearized part</i>	611
William Robin Zame, <i>Extendibility, boundedness and sequential convergence in spaces of holomorphic functions</i>	619