

# Pacific Journal of Mathematics

**ON SOLVABILITY OF GENERALIZED ORTHOMODULAR  
LATTICES**

LADISLAV BERAN

## ON SOLVABILITY OF GENERALIZED ORTHOMODULAR LATTICES

LADISLAV BERAN

**The purpose of this paper is to establish a universal criterion for a generalized orthomodular lattice to belong to a primitive class of lattices.**

The starting point for the investigation is the description of the reflection and the coreflection. These lattices can be determined by two lattice congruences defined by means of alleles on any lattice.

The main results permit to characterize a generalized orthomodular lattice solvable in a nontrivial primitive class  $\mathcal{E}$  of lattices as a lattice belonging to  $\mathcal{E}$ . The characterization is further used to show that such a lattice belongs to  $\mathcal{E}$  if and only if its commutator sublattice is a lattice of  $\mathcal{E}$ .

1. Preliminaries. Basic facts on orthomodular lattices used in this paper may be found in [4], and also most of the notation and terminology will be taken from that book. We assume familiarity with the results of Marsden [8] on solvability of generalized orthomodular lattices. The notation concerning the projectivity of quotients in a lattice is essentially the same as in [7].

Let  $\Omega$  be a nonempty set of quotients of a lattice  $\mathcal{L}$ . A quotient  $b/a$  is called an  $\Omega$ -allele of  $\mathcal{L}$  if there exists  $n \in \mathbb{N}$  and a sequence  $b_0/a_0 = b/a, b_1/a_1, \dots, b_n/a_n$  of quotients of  $\Omega$  such that  $b_i/a_i \sim b'_{i+1}/a'_{i+1}$  for every  $i = 0, 1, \dots, n-1$  where  $[a'_{i+1}, b'_{i+1}] \subset [a_{i+1}, b_{i+1}]$  and  $b \leq a_n$  or  $a \geq b_n$ . The set  $A_\Omega(\mathcal{L})$  of all  $\Omega$ -alleles of  $\mathcal{L}$  will be called the  $\Omega$ -allelomorph of the lattice  $\mathcal{L}$ .

For the special case  $\Omega = \Omega_0 = \{b/a \mid a \prec b \text{ (} b \text{ covers } a)\}$ , the  $\Omega_0$ -alleles of a submodular lattice have been studied in the paper [1]. Other results concerned with  $\Omega$ -alleles can be found in [3].

In the present paper we shall investigate the  $\Omega_1$ -alleles where  $\Omega_1$  is the set of all quotients of  $\mathcal{L}$  and we shall omit reference to  $\Omega_1$ ; e.g., we shall refer to an  $\Omega_1$ -allele as an allele; similarly,  $A(\mathcal{L})$  (or simply  $A$ ) will be the set of all alleles. Hence,  $b/a \in A(\mathcal{L})$  if and only if there exists  $d/c$  such that  $b/a$  is weakly projective into  $d/c$ , i.e.  $b/a \approx_w d/c$ , and  $c \geq b$  or  $d \leq a$ .

2. Reflection and coreflection. For each lattice  $\mathcal{L}$  there exists a smallest congruence  $\Delta$  such that  $\mathcal{L}/\Delta$  is distributive. Let  $\Delta^*$  be the pseudocomplement of  $\Delta$ . We define the reflection of  $\mathcal{L}$ , denoted by  $\text{Ref } \mathcal{L}$ , to be the lattice  $\mathcal{L}/\Delta^*$ ; the lattice  $\mathcal{L}/\Delta$  is denoted by

Coref  $\mathcal{L}$  and it is called the *coreflection* of  $\mathcal{L}$ . A well-known result asserts that a nontrivial primitive class of lattices contains the class of distributive lattices. Since  $\mathcal{L}$  is a subdirect product of Ref  $\mathcal{L}$  and Coref  $\mathcal{L}$ , it follows that a lattice belongs to a nontrivial class  $\mathcal{E}$  of lattices if and only if its reflection belongs to  $\mathcal{E}$ .

The following lemma (which appears in [9, p. 95] as Lemma 2) is useful:

**LEMMA 2.1.** *Let  $\mathcal{L}$  be a lattice and  $\theta$  a congruence of  $\mathcal{L}$ . If  $[b]/[a] \approx_w [d]/[c]$  in  $\mathcal{L}/\theta$ , then there exists  $\backslash a \in [a]$ ,  $\backslash b \in [b]$ ,  $\backslash c \in [c]$ ,  $\backslash d \in [d]$  such that  $\backslash b/\backslash a \approx_w \backslash d/\backslash c$  in  $\mathcal{L}$ .*

*Proof.* Suppose  $[b]/[a] = [b_0]/[a_0] \sim_w [b_1]/[a_1] \sim_w \dots \sim_w [b_n]/[a_n] = [d]/[c]$  with  $a_0 = a \leq b = b_0$  and  $a_n = c \leq d = b_n$ . Let  $\backslash d = \backslash b_n = d$  and  $\backslash c = \backslash a_n = c$ , and suppose that  $\backslash a_j, \backslash b_j$  for  $i < j < n$  have already been defined so that  $\backslash a_j \in [a_j]$ ,  $\backslash b_j \in [b_j]$ ,  $\backslash a_j \leq \backslash b_j$ , and  $\backslash b_j/\backslash a_j \sim_w \backslash b_{j+1}/\backslash a_{j+1}$ . Then if  $[b_i]/[a_i] \nearrow_w [b_{i+1}]/[a_{i+1}]$ , let  $\backslash a_i = \backslash a_{i+1} \wedge \backslash b_i$ ,  $\backslash b_i = \backslash b_{i+1} \wedge \backslash b_i$ , and if  $[b_i]/[a_i] \searrow_w [b_{i+1}]/[a_{i+1}]$ , let  $\backslash a_i = \backslash a_{i+1} \vee \backslash a_i$ ,  $\backslash b_i = \backslash b_{i+1} \vee \backslash a_i$ . So we get  $\backslash a = \backslash a_0 \in [a]$ ,  $\backslash b = \backslash b_0 \in [b]$ , and  $\backslash b/\backslash a \approx_w \backslash d/\backslash c$ . Note that if  $[a] \neq [b]$ , we must have  $\backslash a \neq \backslash b$ .

**COROLLARY 2.2.** *Let  $\mathcal{L}$  be a lattice and  $\theta$  a congruence of  $\mathcal{L}$ . If  $[b]/[a]$  is an allele of  $\mathcal{L}/\theta$ , then there exists  $\backslash a \in [a]$ ,  $\backslash b \in [b]$  such that  $\backslash b/\backslash a$  is an allele of  $\mathcal{L}$ .*

*Proof.* Let  $[b]/[a] \approx_w [d]/[c]$  with  $a \leq b \leq c \leq d$ . Choose  $\backslash a, \backslash b, \backslash c, \backslash d$  as in Lemma 2.1. It suffices to prove that  $\backslash b \leq \backslash c$ . But we may assume that  $[b]/[a] = [b_0]/[a_0] \nearrow_w [b_1]/[a_1]$ , so  $\backslash b = \backslash b_0 = \backslash b_1 \wedge \backslash b_0 = \backslash b_1 \wedge b \leq \backslash b \leq c = \backslash c$ .

**LEMMA 2.3.** (i) *Let  $\mathcal{L}$  be any lattice and let  $a < b, r < s$  be elements of  $\mathcal{L}$ . Then if  $b/a \approx_w s/r$  and  $r = r_0 < r_1 < \dots < r_n = s$ , there exist  $a_j, j = 0, 1, \dots, n$  such that  $a = a_0 \leq a_1 \leq \dots \leq a_n = b$  and  $a_{j+1}/a_j \approx_w r_{j+1}/r_j$  for each  $j$ .*

(ii) *Let  $\mathcal{L}$  be a lattice and let  $\gamma$  be the binary relation defined on  $L$  by*

$$a \equiv b(\gamma) \Leftrightarrow \exists n \in \mathbb{N} \exists a_1, a_2, \dots, a_n$$

$$a \wedge b = a_0 \leq a_1 \leq \dots \leq a_n = a \vee b,$$

and

$$a_{j+1}/a_j \in \mathbf{A}(\mathcal{L}) \text{ for every } j = 0, 1, \dots, n - 1.$$

*If  $b/a \approx_w q/p$  and  $p \equiv q(\gamma)$ , then  $a \equiv b(\gamma)$ .*

*Proof.* (i) It is enough to consider the case  $b/a \sim_w s/r$ . Assume  $b/a \searrow_w s/r$ ; then  $a_j = a \vee r_j, j = 0, 1, \dots, n$  are the required elements.

(ii) We will first treat the following two cases:

*Case I.*  $b/a \searrow_w q/p, q/p \approx_w s/r$  and  $r \geq q$ . Let  $r_0 = r, r_1 = r \vee (b \wedge s)$  and  $r_2 = s$ . Since  $r_1/r \searrow b \wedge s/b \wedge r$  and  $b/q \searrow a/a \wedge q$ , we get that  $r_1/r_0 \approx_w a/a \wedge q$ . We also have  $r_2/r_1 \approx_w b \vee s/b$ . Since  $b/a \approx_w s/r$ , (i) implies that there exist  $a_0 = a \leq a_1 \leq a_2 = b$  such that  $a_1/a_0 \approx_w a/a \wedge q$  and  $a_2/a_1 \approx_w b \vee s/b$ . Thus  $a_1/a$  and  $b/a_1$  belong to the allelomorph and so  $a \equiv b(\gamma)$ .

*Case II.*  $b/a \nearrow_w q/p, q/p \approx_w s/r$  and  $r \geq q$ . However, here we have  $b/a \approx_w s/r$  and  $r \geq q \geq p \vee b \geq b$ . This yields  $b/a \in \mathbf{A}$  and, consequently,  $a \equiv b(\gamma)$ .

By the same argument as in the proof of (i) it is clear that  $b/a \sim_w q/p$  and  $p \equiv q(\gamma)$  implies  $a \equiv b(\gamma)$ . The general case now follows by induction.

Our next theorem provides much more information on  $\gamma$ .

**THEOREM 2.4.** *Let  $\mathcal{L}$  be a lattice. Then  $\gamma$  is a congruence relation of  $\mathcal{L}$ .*

*Proof.* If  $x \leq y, x \equiv y(\gamma)$  and  $t \in L$ , then  $y \wedge t/x \wedge t \approx_w y/x$ . By Lemma 2.3 (ii) it follows that  $x \wedge t \equiv y \wedge t(\gamma)$ . The conclusion is now immediate from [6, Lemma 8, p. 24].

The following theorem can be proved by similar arguments; we omit the proof.

**THEOREM 2.5.** *Let  $\mathcal{L}$  be a lattice and let  $\beta$  be the binary relation defined on  $L$  by*

$$a \equiv b(\beta) \Leftrightarrow \{(d/c \approx_w a \vee b/a \wedge b \ \& \ d/c \in \mathbf{A}(\mathcal{L})) \Rightarrow c = d\}.$$

*Then  $\beta$  is a congruence of  $\mathcal{L}$ . Moreover,  $\beta$  is the pseudocomplement of  $\gamma$ .*

**COROLLARY 2.6.** *Let  $\mathcal{L}$  be a weakly modular lattice. Then*

$$a \equiv b(\beta) \Leftrightarrow \{([m, n] \subset [a \wedge b, a \vee b] \ \& \ n/m \in \mathbf{A}(\mathcal{L})) \Rightarrow m = n\}.$$

Observe that the description of  $\beta$  as given in Corollary 2.6 is a direct consequence of the proof of [7, Theorem III. 4. 10].

**REMARK.** If  $\mathcal{L}$  is a relatively complemented lattice, then [5,

p. 355]  $y/x \approx_w b/a$  implies  $y/x \approx b'/a'$  where  $[a', b'] \subset [a, b]$ . We say that  $b/a$  has a *close allele*  $d/c$  if there exists a quotient  $d/c$  such that  $b/a \approx d/c$  &  $(b \leq c$  or  $a \geq d)$ . Thus, in such a lattice  $b/a \in \mathbf{A}$  if and only if  $b/a$  has a close allele and the congruences  $\beta$  and  $\gamma$  of  $\mathcal{L}$  can be characterized in terms of close alleles.

**PROPOSITION 2.7.** *Let  $\mathcal{L}$  be a lattice and let  $\theta$  be a congruence relation of  $\mathcal{L}$ . Then the quotient lattice  $\mathcal{L}/\theta$  is distributive if and only if  $\theta \supset \gamma$ .*

*Proof.* If  $\mathcal{L}/\theta$  is distributive and if  $a \equiv b(\gamma)$ , then there are elements  $x_i$  such that

$$a \wedge b = x_0 \leq x_1 \leq \dots \leq x_s = a \vee b$$

and such that  $x_{i+1}/x_i \in \mathbf{A}(\mathcal{L})$  for every  $i = 0, 1, \dots, s-1$ . But then  $[x_{i+1}]/[x_i] \in \mathbf{A}(\mathcal{L}/\theta)$  and by [7, Lemma III. 2.7] we have  $[x_{i+1}] = [x_i]$ . Thus  $[a] = [b]$ , i.e.,  $a \equiv b(\theta)$  and so  $\gamma \subset \theta$ .

To prove the converse, suppose  $\mathcal{L}/\gamma$  is not distributive; then there exists an allele  $[b]/[a]$  in  $\mathcal{L}/\gamma$  such that  $[a] \neq [b]$ . By Corollary 2.2 there exists  $\backslash a \in [a]$ ,  $\backslash b \in [b]$  such that  $\backslash b/\backslash a$  is an allele in  $\mathcal{L}$ ; hence  $\backslash a \equiv \backslash b(\gamma)$  and so we get  $[a] = [b]$ , a contradiction.

The following is an application of Proposition 2.7.

**COROLLARY 2.8.** *If  $\mathcal{L}$  is a lattice, then  $\text{Coref } \mathcal{L} = \mathcal{L}/\gamma$  and  $\text{Ref } \mathcal{L} = \mathcal{L}/\beta$ .*

**3. Solvability in classes of lattices.** If  $a, b$  are two elements of an ortholattice  $\mathcal{L} = (L, \vee, \wedge, ', 0, 1)$ , then

$$\text{com}_{\mathcal{L}}(a, b) = (a \vee b) \wedge (a \vee b') \wedge (a' \vee b) \wedge (a' \vee b')$$

is called the *commutator* of  $a, b$  (cf. [8]). The  $n$ -th *commutator sublattice*  $\mathcal{G}_n$  of a generalized orthomodular lattice  $\mathcal{G}$  is defined by induction in the following way:  $\mathcal{G}_0 = \mathcal{G}$  and  $\mathcal{G}_n (n \geq 1)$  is by definition the  $p$ -ideal generated in  $\mathcal{G}_{n-1}$  by all the commutators of the generalized orthomodular lattice  $\mathcal{G}_{n-1}$ . The lattice  $\mathcal{G}$  is said to be *solvable* (in the sense of Marsden) if there exists  $n \in \mathbf{N}$  such that  $\mathcal{G}_n = \langle 0 \rangle$ . Recall that [8, Theorem 9, p. 361]  $\mathcal{G}$  is solvable if and only if it is distributive.

It is easily verified that if we define  $\mathcal{G}^{(0)} = \mathcal{G}$  and if  $\mathcal{G}^{(n)} (n \geq 1)$  equals to the ideal of the generalized orthomodular lattice  $\mathcal{G}$  generated by all the commutators of  $\mathcal{G}^{(n-1)}$ , then  $\mathcal{G}_n = \mathcal{G}^{(n)}$  for every  $n = 0, 1, \dots$ . A generalized orthomodular lattice  $\mathcal{G}$  is said to be *solvable in a class*  $\mathcal{C}$  of lattices if and only if there exists  $n \in \mathbf{N}$  such that

$\mathcal{E}^{(n)}$  belongs to  $\mathcal{E}$ .

Since it is known [8, Theorem 9, p. 361] that  $\mathcal{E}_1 = \mathcal{E}_2 = \dots$ , it is clear that in the definitions above we can suppose  $n = 1$ . We remark that it is a simple matter to verify that a lattice  $\mathcal{E}$  is solvable in the class of distributive lattices if and only if it is solvable in the sense of Marsden.

The solvability of generalized orthomodular lattices in the class of modular lattices has been investigated in [2].

**PROPOSITION 3.1.** *Let  $\mathcal{E}$  be a generalized orthomodular lattice. Then an element  $a$  of  $\mathcal{E}$  belongs to  $\mathcal{E}'$  if and only if  $a \equiv 0(\gamma)$ .*

*Proof.* By [8, Theorems 6 and 7, p. 360], the commutator sublattice  $\mathcal{E}'$  is the kernel of the congruence  $\Delta$ . The conclusion  $\gamma = \Delta$  is now immediate from Proposition 2.7.

**THEOREM 3.2.** *The commutator sublattice  $\mathcal{E}'$  of a generalized orthomodular lattice  $\mathcal{E}$  is a dually distributive ideal.*

*Proof.* To prove the assertion, it is sufficient to show that if  $I_1, I_2$  are ideals of  $\mathcal{E}$ , then necessarily  $G' \cap (I_1 \vee I_2) \subset (G' \cap I_1) \vee (G' \cap I_2)$ . Suppose  $m \in G'$  and  $m \leq i_1 \vee i_2$ ,  $i_s \in I_s (s = 0, 1)$ . Let  $x'$  denote the orthocomplement of an element  $x \in [0, i_1 \vee i_2]$  in the orthomodular lattice determined by the interval  $[0, i_1 \vee i_2]$ .

Set

$$\begin{aligned} k_1 &= i_1 \wedge (i'_1 \vee m \vee i_2) \wedge (i'_1 \vee m \vee i'_2), \\ k_2 &= i_2 \wedge (i'_2 \vee m \vee i_1) \wedge (i'_2 \vee m \vee i'_1) \end{aligned}$$

so that  $k_s = i_s \wedge c \in I_s$  where

$$c = (i'_1 \vee m \vee i_2) \wedge (i'_1 \vee m \vee i'_2) \wedge (i'_2 \vee m \vee i_1).$$

Let  $w = i_1 \wedge (i'_1 \vee m)$  and  $v = i_1 \wedge m$ . Since  $v \leq m \in G'$ ,  $v \in G'$ . Now we have  $w \wedge v' = i_1 \wedge \text{com}(i_1, m) \in G'$  and so  $w = v \vee (w \wedge v') \in G'$ .

Let  $w^+ = w' \wedge k_1$ . Since  $i_1$  commutes with  $i'_1$  and  $i_1 \wedge m'$ , it follows that  $[i'_1 \vee (i_1 \wedge m')] \wedge i_1 = i_1 \wedge m'$ . Thus  $w^+ \leq \text{com}(i'_1 \vee m, i_2)$  and this yields  $w^+ \in G'$ . Therefore  $k_1 = w \vee w^+ \in G' \cap I_1$  and, by symmetry,  $k_2 \in G' \cap I_2$ .

It is clear that  $m \leq (i_1 \vee i_2) \wedge c$  and that every  $i_s$  commutes with each element  $(\dots)$  of the definition of the element  $c$ . Therefore,  $i_1$  and  $i_2$  commute with  $c$  and this gives

$$m \leq (i_1 \vee i_2) \wedge c = (i_1 \wedge c) \vee (i_2 \wedge c) = k_1 \vee k_2.$$

This shows that  $m \in (G' \cup I_1) \vee (G' \cap I_2)$ , completing the proof.

**COROLLARY 3.3.** *The ideal  $\mathcal{S}'$  of a generalized orthomodular lattice  $\mathcal{S}$  is neutral.*

*Proof.* By Proposition 3.1 and Theorem III. 3.10 of [7], the ideal  $\mathcal{S}'$  is standard. By Theorem 3.2 and Theorem III. 2.5 of [7] we see that  $\mathcal{S}'$  is neutral.

For an element  $g \in G$  let  $[g]$  denote the corresponding element of the quotient lattice  $\mathcal{S}/\beta$  where  $\mathcal{S}$  is a generalized orthomodular lattice and let  $K([g]) = (g) \cap G'$ . If  $g_1 \in [g]$  and  $k \in K([g])$ , then  $k \equiv k \wedge g_1 \wedge g(\gamma)$ . Since  $k/k \wedge g_1 \wedge g \nearrow_w g/g_1 \wedge g$  and  $g \equiv g_1 \wedge g(\beta)$ , we have  $k \equiv k \wedge g_1 \wedge g(\beta)$ . By Theorem 2.5 this implies  $k \leq g_1 \wedge g \leq g_1$ . The set  $K([g])$  is therefore well-defined.

**LEMMA 3.4.** *Under the convention made above we have*

- (i)  $[g] = \sup_{\mathcal{S}/\beta} \{[k] \mid k \in K([g])\}$ ;
- (ii)  $K([g_1 \wedge g_2]) = K([g_1]) \wedge K([g_2])$   
 $K([g_1 \vee g_2]) = K([g_1]) \vee K([g_2])$

where the symbols  $\vee, \wedge$  of the right-hand side denote the join and the meet in the ideal lattice  $\mathcal{S}$  of the lattice  $\mathcal{S}'$ ;

(iii) the mapping  $f: [g] \mapsto K([g])$  is an isomorphism of the lattice  $\mathcal{S}/\beta$  onto a sublattice of the lattice  $\mathcal{S}$ .

*Proof.* (i) If  $[h]$  is such that  $[h] < [g]$  with  $h < g$ , then, by Corollary 2.6, there are  $v, w$  satisfying  $h \leq v < w \leq g$  and  $w/v \in \mathbf{A}$ . Let  $z$  denote a complement of  $v$  in  $[0, w]$ . Then  $v \equiv w(\gamma)$  implies  $z \in G'$ . If  $[z] \leq [h]$ , then  $[v] = [w]$  and so  $v \equiv w(\beta \cap \gamma)$ . Therefore, by Theorem 2.5,  $v = w$ , a contradiction. Since  $z \in K([g])$ , we see that  $[h]$  is not an upper bound for the set  $\{[k] \mid k \in K([g])\}$ . Conclusion (i) results.

(ii) This is immediate by Theorem 3.2.

Assertion (iii) now follows directly from (i) and (ii) above.

**THEOREM 3.5.** *Let  $\mathcal{E}$  be a primitive class of lattices which contains a lattice with more than one element. Then a generalized orthomodular lattice is a lattice of  $\mathcal{E}$  if and only if its commutator sublattice belongs to  $\mathcal{E}$ .*

**COROLLARY 3.6.** *A generalized orthomodular lattice is solvable in a primitive class  $\mathcal{E}$  which contains a lattice with more than one element if and only if it belongs to the class  $\mathcal{E}$ .*

*Proof.* If  $\mathcal{S}$  belongs to  $\mathcal{E}$ , then its sublattice  $\mathcal{S}'$  belongs also to  $\mathcal{E}$ .

Conversely, suppose  $\mathcal{G}' \in \mathcal{E}$ . By [6, Lemma 8, p. 34], the ideal lattice  $\mathcal{I}$  of  $\mathcal{G}'$  also belongs to the class  $\mathcal{E}$ . Since  $\mathcal{E}$  is primitive, every isomorphic image of a sublattice of  $\mathcal{I}$  belongs to  $\mathcal{E}$ . It, therefore, follows from Lemma 3.4 (iii) that  $\text{ref } \mathcal{G} \in \mathcal{E}$ . It is now immediate that  $\mathcal{G}$  belongs to  $\mathcal{E}$ .

The author wishes to thank the referee for suggestions on the formulation of the theorems in § 2 and particularly for his pointing out the consequence stated in Corollary 3.3 as well as for his many other valuable comments.

#### REFERENCES

1. L. Beran, *Treillis sous-modulaires*, Séminaire Dubreil-Pisot: Algèbre et théorie des nombres, Fac. Sci. Paris, 21e année, (1967/68), n° 13, p. 13.01-13.17.
2. ———, *Modularity in generalized orthomodular lattices*, Comment. Math. Univ. Carolinae **15**, No. 1 (1974), 189-193.
3. ———, *Reflection and coreflection in generalized orthomodular lattices* (to appear).
4. G. Birkhoff, *Lattice Theory*, Amer. Math. Soc. Colloq. Publ. Vol. XXV, Providence, R. I. (1967).
5. R. P. Dilworth, *The structure of relatively complemented lattices*, Ann. of Math., **51** (1950), 348-359.
6. G. Grätzer, *Lattice Theory (First Concepts and Distributive Lattices)*, W. H. Freeman, San Francisco 1971.
7. ———, *Lattice Theory (Second Part)* (to appear).
8. E. L. Marsden, Jr., *The commutator and solvability in a generalized orthomodular lattice*, Pacific J. Math., **33** (1970), 357-361.
9. R. Wille, *Primitive subsets of lattices*, Algebra Universalis, **2** (1972), 95-98.

Received February 28, 1974 and in revised form November 11, 1974.

CHARLES UNIVERSITY, PRAGUE  
SOKOLOVSKÁ 83 CZECHOSLOVAKIA





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

# Pacific Journal of Mathematics

Vol. 57, No. 2

February, 1975

Norman Larrabee Alling, <i>On Cauchy's theorem for real algebraic curves with boundary</i> . . . . .	315
Daniel D. Anderson, <i>A remark on the lattice of ideals of a Prüfer domain</i> . . . . .	323
Dennis Neal Barr and Peter D. Miletta, <i>A necessary and sufficient condition for uniqueness of solutions to two point boundary value problems</i> . . . . .	325
Ladislav Beran, <i>On solvability of generalized orthomodular lattices</i> . . . . .	331
L. Carlitz, <i>A three-term relation for some sums related to Dedekind sums</i> . . . . .	339
Arthur Herbert Copeland, Jr. and Albert Oscar Shar, <i>Images and pre-images of localization maps</i> . . . . .	349
G. G. Dandapat, John L. Hunsucker and Carl Pomerance, <i>Some new results on odd perfect numbers</i> . . . . .	359
M. Edelstein and L. Keener, <i>Characterizations of infinite-dimensional and nonreflexive spaces</i> . . . . .	365
Francis James Flanigan, <i>On Levi factors of derivation algebras and the radical embedding problem</i> . . . . .	371
Harvey Friedman, <i>Provable equality in primitive recursive arithmetic with and without induction</i> . . . . .	379
Joseph Braucher Fugate and Lee K. Mohler, <i>The fixed point property for tree-like continua with finitely many arc components</i> . . . . .	393
John Norman Ginsburg and Victor Harold Saks, <i>Some applications of ultrafilters in topology</i> . . . . .	403
Arjun K. Gupta, <i>Generalisation of a "square" functional equation</i> . . . . .	419
Thomas Lee Hayden and Frank Jones Massey, <i>Nonlinear holomorphic semigroups</i> . . . . .	423
V. Kannan and Thekkedath Thrivikraman, <i>Lattices of Hausdorff compactifications of a locally compact space</i> . . . . .	441
J. E. Kerlin and Wilfred Dennis Pepe, <i>Norm decreasing homomorphisms between group algebras</i> . . . . .	445
Young K. Kwon, <i>Behavior of <math>\Phi</math>-bounded harmonic functions at the Wiener boundary</i> . . . . .	453
Richard Arthur Levaro, <i>Projective quasi-coherent sheaves of modules</i> . . . . .	457
Chung Lin, <i>Rearranging Fourier transforms on groups</i> . . . . .	463
David Lowell Lovelady, <i>An asymptotic analysis of an odd order linear differential equation</i> . . . . .	475
Jerry Malzan, <i>On groups with a single involution</i> . . . . .	481
J. F. McClendon, <i>Metric families</i> . . . . .	491
Carl Pomerance, <i>On multiply perfect numbers with a special property</i> . . . . .	511
Mohan S. Putcha and Adil Mohamed Yaqub, <i>Polynomial constraints for finiteness of semisimple rings</i> . . . . .	519
Calvin R. Putnam, <i>Hyponormal contractions and strong power convergence</i> . . . . .	531
Douglas Conner Ravenel, <i>Multiplicative operations in <math>BP^*BP</math></i> . . . . .	539
Judith Roitman, <i>Attaining the spread at cardinals which are not strong limits</i> . . . . .	545
Kazuyuki Saitô, <i>Groups of <math>*</math>-automorphisms and invariant maps of von Neumann algebras</i> . . . . .	553
Brian Kirkwood Schmidt, <i>Homotopy invariance of contravariant functors acting on smooth manifolds</i> . . . . .	559
Kenneth Barry Stolarsky, <i>The sum of the distances to <math>N</math> points on a sphere</i> . . . . .	563
Mark Lawrence Teply, <i>Semiprime rings with the singular splitting property</i> . . . . .	575
J. Pelham Thomas, <i>Maximal connected Hausdorff spaces</i> . . . . .	581
Charles Thomas Tucker, II, <i>Concerning <math>\sigma</math>-homomorphisms of Riesz spaces</i> . . . . .	585
Rangachari Venkataraman, <i>Compactness in abelian topological groups</i> . . . . .	591
William Charles Waterhouse, <i>Basically bounded functors and flat sheaves</i> . . . . .	597
David Westreich, <i>Bifurcation of operator equations with unbounded linearized part</i> . . . . .	611
William Robin Zame, <i>Extendibility, boundedness and sequential convergence in spaces of holomorphic functions</i> . . . . .	619