

Pacific Journal of Mathematics

**LATTICES OF HAUSDORFF COMPACTIFICATIONS OF A
LOCALLY COMPACT SPACE**

V. KANNAN AND THEKKEDATH THRIVIKRAMAN

LATTICES OF HAUSDORFF COMPACTIFICATIONS OF A LOCALLY COMPACT SPACE

V. KANNAN AND T. THRIVIKRAMAN

This paper gives a lattice theoretic characterization of (complete) lattices which are lattices of Hausdorff compactifications of locally compact spaces. This is accomplished via a characterization of the lattices of closed equivalence relations on T_1 spaces.

NOTATIONS. L is a complete dually atomic lattice.
 D is the set of all dual atoms of L .

1. DEFINITION. Let $p \in L$. Then the set $H(p) = \{d \in D \mid d \geq p\}$ is called the hull of p .

NOTE. $H(1) = \emptyset$.

2. DEFINITION. Let $E \subset D$. The kernel of E denoted by $\text{Ker } E$ is defined as $\bigwedge_{d \in E} d$.

3. DEFINITION. Let $p \in L$. p is said to be a primary element if whenever q and r are two elements in $H(p)$ such that $\text{Card. } H(q \wedge r) \neq 3$ and $s \in D$ is such that $\text{Card. } H(s \wedge q) = 3 = \text{Card. } H(s \wedge r)$, it is true that $s \in H(p)$.

NOTE. Trivially, 1 as well as any dual atom is primary.

4. DEFINITION. A star of L is defined as a subset S of D which is maximal with respect to the following property: $d, d' \in S \Rightarrow d \wedge d'$ is a primary element and if (d, d') and (d_1, d_2) are distinct pairs of elements of S , then $d \wedge d' \neq d_1 \wedge d_2$.

5. DEFINITION. Let p, q be primary elements. Then a primary rectangle is defined as $H(p, q) = \{(S_1, S_2) \mid S_1, S_2 \text{ are stars such that } S_1 \cap H(p) \neq \emptyset \neq S_2 \cap H(q)\}$.

6. DEFINITION. Let $A \subset D$. $\alpha(A)$ is defined as the set of all unordered pairs (S, S') of stars such that $S \cap S' \cap A \neq \emptyset$.

7. THEOREM. *A complete, dually atomic lattice L is isomorphic to the lattice of closed equivalence relations of a T_1 space X if and*

only if the following are true:

- (i) $\bigvee_{i \in J} p_i$ is a primary element for any collection $\{p_i | i \in J\}$ of primary element in L .
- (ii) (a) If $d \in D$, then d belongs to exactly two stars.
 (b) Any two stars intersect in a singleton.
- (iii) $H \subset D$ is a hull if and only if
 - (a) if $d_1, d_2 \in H$ and if $d \in D$ such that $d \geq d_1 \wedge d_2$, then $d \in H$.
 - (b) $\alpha(H)$ is an intersection of finite unions of primary rectangles.
- (iv) $a = \text{Ker}(H(a))$ for every $a \in L$.

Proof (Necessity). Easily checked, bearing in mind the discussion in §1 of [3].

(Sufficiency). Let X be the set of all stars in L . From (ii) (a) and (ii) (b), there exists a bijection θ between the set D of all dual atoms of L and the set of all unordered pairs of distinct stars.

From (i), and noticing that $0 \in L$ is a primary element, it follows that primary elements of L form a complete lattice P under the same order. Now D is precisely the set of all dual atoms of P . We can form the hull-kernel topology for D in the lattice. This topology can then be translated to X as follows:

A set $A \subset X$ is closed if and only if $\theta^{-1}(A \times A)$ is a hull of a member of P . We show now that this defines a T_1 topology on X . Clearly $\theta^{-1}(\emptyset \times \emptyset) = \emptyset$ and $\theta^{-1}(X \times X) = D$ so that \emptyset and X are closed. If $S \in X$, then $\theta^{-1}(\{S\} \times \{S\}) = \emptyset = H(1)$ so that every singleton is closed. If $S \neq S'$, $S, S' \subset X$, $\theta^{-1}(\{S\} \times \{S'\})$ is a singleton. So any two-element set is closed.

Now let $A, B \subset X$ be closed, each containing at least two elements. Then $A \cup B$ is closed. For, let p and q be the primary elements determined by A and B respectively. Then consider the primary rectangles $H(p, p)$, $H(q, q)$, $H(p, q)$, $H(q, p)$. Now if $C = \theta^{-1}((A \cup B) \times (A \cup B))$, then $\alpha(C)$ is the union of these primary rectangles. For, if $(S, S') \in \alpha(C)$, then there exists $d \in C$ such that $S \cap S' = \{d\}$; now $\theta(d) = (S, S') \in ((A \cup B) \times (A \cup B))$ so that $S, S' \in A \cup B$; it is easy to check that if S, S' both belong to A (respectively B), then $(S, S') \in H(p, p)$ (respectively $H(q, q)$). If $S \in A$ and $S' \in B$, then $(S, S') \in H(p, q)$ and if $S \in B$ and $S' \in A$, then $(S, S') \in H(q, p)$. On the other hand, that all these four primary rectangles are subsets of $\alpha(C)$ is easily verified.

Hence by condition (iii), C is a hull set. Note that condition (iii) (a) is satisfied here, by the maximality in the definition of stars.

Let K be the kernel of C . It can be seen that K is primary. It follows that $A \cup B$ is closed.

Let $A_i \subset X$ be closed for every $i \in J$ and let $A = \bigcap A_i$. Then

$\theta^{-1}(A \times A)$ is the intersection $\cap \theta^{-1}(A_i \times A_i)$ and so is a hull set. For, by condition (iii), intersection of hull sets is again a hull set. Let r be its kernel. Then by using condition (iv), it can be proved that $r = \vee p_i$ where p_i is the primary element corresponding to A_i . So it follows from condition (i) that r is primary. So A is closed. Thus X is a T_1 space.

Next we show that L is isomorphic to $L(X)$, the lattice of closed equivalence relations for this space X . Let us define a map $\eta: L \rightarrow L(X)$ as follows: If $x \in L$, $\eta(x)$ is the relation defined on X as below: $S_1 \eta(x) S_2$ if and only if either $S_1 = S_2$ or $S_1 \cap S_2 \cap H(x) \neq \emptyset$. This is an equivalence relation. For, let $S_1 \eta(x) S_2$ and $S_2 \eta(x) S_3$; also let $S_1 \cap S_2 = \{d_3\}$, $S_2 \cap S_3 = \{d_1\}$, and $S_3 \cap S_1 = \{d_2\}$; then by the maximality in the definition of stars, it can be proved that $d_2 \geq d_1 \wedge d_3$ and so by (iii) (a), $d_2 \in H(x)$ since $H(x)$ is a hull set. So $S_1 \eta(x) S_3$. Thus transitivity is proved. Reflexivity and symmetry of $\eta(x)$ are trivial.

Now we show that $\eta(x) \in L(X)$. That is, $\eta(x)$ is a closed equivalence relation on X . The relation $\eta(x)$ viewed as a subset of $X \times X$ is precisely $\alpha(H(x))$. So by (iii)(b), it is an intersection of finite union of primary rectangles. Each primary rectangle is a closed subset of $X \times X$ since $H(p, q) = A \times B$ where A and B are the closed subsets of X determined by p and q respectively. So $\eta(x)$ is a closed subset of $X \times X$.

Thus $\eta: L \rightarrow L(X)$ is well defined.

Now η is injective. For, let $x \neq y$. Then by condition (iv) we get that $H(x) \neq H(y)$. Let $d \in H(x) - H(y)$ (say). Let S, S' be the two stars containing d . Then $(S, S') \in \alpha(H(x))$ but $\notin \alpha(H(y))$. So $\eta(x) \neq \eta(y)$. Further η is surjective. For, let R be a closed equivalence relation on X . Then R is an intersection of finite unions of closed rectangles, whereas each closed rectangle is a primary rectangle. Let $H = \{d \in D \mid \text{the pair } (S_1, S_2) \text{ of stars containing } d \text{ belongs to } R\}$. Then H satisfies conditions (iii) (a) and (iii) (b) and hence by (iii), H is a hull set. If x is its kernel, it follows that $\eta(x) = R$.

η preserves arbitrary unions. For, let $x = \bigvee_{\alpha} x_{\alpha}$. Now $\cap H(x_{\alpha})$ is a hull set by (iii). It ought to contain $H(x)$. Let y be its kernel. Then $y \geq x_{\alpha}$ for each α . So $y \geq x$. Therefore, $H(y) \subset H(x)$. So $\eta(y) = \eta(x)$. But $\eta(y) = \bigvee_{\alpha} \eta(x_{\alpha})$ since it can be easily seen that $S_1 \bigvee_{\alpha} \eta(x_{\alpha}) S_2$ if and only if $S_1 \cap S_2 \cap (\cap H(x_{\alpha})) \neq \emptyset$, while $\cap H(x_{\alpha}) = H(y)$.

η preserves intersections. This can be seen as in the case of unions, by considering the union of the hulls.

Thus η is an isomorphism.

Thus sufficiency is proved.

8. THEOREM. *Let L be a complete, dually atomic lattice. Then L is the lattice of T_2 compactifications of a locally compact space if*

and only if it satisfies conditions (i) through (iv) above and also:

(v) Given any two primary elements $p_1, p_2 \in L$, there exist primary elements $q_1, q_2 \in L$ such that $p_1 \vee q_1 = p_2 \vee q_2 = 1$ and $q_1 \wedge q_2 = 0$.

(vi) Given any collection of primary elements $\{p_\alpha\}_{\alpha \in J}$ such that $\bigwedge_{\alpha \in K} p_\alpha$ is a primary element for any finite subset K of J , then $\bigwedge_{\alpha \in J} p_\alpha$ is also a primary element.

Proof. Notice that (v) is equivalent to saying that the space X uniquely specified by L , is normal.

Also (vi) is equivalent to compactness.

It can be proved that the lattice of all closed equivalence relations of $\beta X - X$ is isomorphic to the lattice of all Hausdorff compactifications of X , when X is locally compact and Hausdorff.

Thus the theorem.

9. REMARK. When X is not locally compact, the upper semilattice $K(X)$ of all Hausdorff compactifications of X is not necessarily a lattice (cf. [5]). Now the problem arises whether the semilattice $K(X)$ determines the space $\beta X - X$. When X is locally compact, the answer is in the affirmative (cf. [1]). A method to construct the space $\beta X - X$ from $K(X)$ is given in [2]. But when x is not locally compact, $K(X)$ does not determine $\beta X - X$ (cf. [2]). A study in this direction forms a part of [4].

REFERENCES

1. K. D. Magill, Jr., *The lattice of compactifications of a locally compact space*, Proc. London Math. Soc., **18** (1968), 231-244.
2. T. Thrivikraman, *On compactifications of Tychonoff spaces*, Yokohama Math. J., **20** (1972), 99-106.
3. ———, *On Hausdorff quotients of spaces and Magill's theorem*, Monat. für-Math., **76** (1972), 345-355.
4. ———, *Studies in Compactifications and Other Extensions*, Doctoral Thesis submitted to Madurai University, July, 1971.
5. J. Visliseni and J. Flachsmeyer, *The power and structure of the lattice of all compactifications of a completely regular space*, Doklady Nauk SSSR (N. S.), **165** (1965), 1423-1425.

Received July 16, 1973 and in revised form April 30, 1974.

MADURAI UNIVERSITY, MADURAI-21

AND

MAR ATHANASTUS COLLEGE, KOTHAMANGALAM, KERALA, INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 57, No. 2

February, 1975

Norman Larrabee Alling, <i>On Cauchy's theorem for real algebraic curves with boundary</i>	315
Daniel D. Anderson, <i>A remark on the lattice of ideals of a Prüfer domain</i>	323
Dennis Neal Barr and Peter D. Miletta, <i>A necessary and sufficient condition for uniqueness of solutions to two point boundary value problems</i>	325
Ladislav Beran, <i>On solvability of generalized orthomodular lattices</i>	331
L. Carlitz, <i>A three-term relation for some sums related to Dedekind sums</i>	339
Arthur Herbert Copeland, Jr. and Albert Oscar Shar, <i>Images and pre-images of localization maps</i>	349
G. G. Dandapat, John L. Hunsucker and Carl Pomerance, <i>Some new results on odd perfect numbers</i>	359
M. Edelstein and L. Keener, <i>Characterizations of infinite-dimensional and nonreflexive spaces</i>	365
Francis James Flanigan, <i>On Levi factors of derivation algebras and the radical embedding problem</i>	371
Harvey Friedman, <i>Provable equality in primitive recursive arithmetic with and without induction</i>	379
Joseph Braucher Fugate and Lee K. Mohler, <i>The fixed point property for tree-like continua with finitely many arc components</i>	393
John Norman Ginsburg and Victor Harold Saks, <i>Some applications of ultrafilters in topology</i>	403
Arjun K. Gupta, <i>Generalisation of a "square" functional equation</i>	419
Thomas Lee Hayden and Frank Jones Massey, <i>Nonlinear holomorphic semigroups</i>	423
V. Kannan and Thekkedath Thrivikraman, <i>Lattices of Hausdorff compactifications of a locally compact space</i>	441
J. E. Kerlin and Wilfred Dennis Pepe, <i>Norm decreasing homomorphisms between group algebras</i>	445
Young K. Kwon, <i>Behavior of Φ-bounded harmonic functions at the Wiener boundary</i>	453
Richard Arthur Levaro, <i>Projective quasi-coherent sheaves of modules</i>	457
Chung Lin, <i>Rearranging Fourier transforms on groups</i>	463
David Lowell Lovelady, <i>An asymptotic analysis of an odd order linear differential equation</i>	475
Jerry Malzan, <i>On groups with a single involution</i>	481
J. F. McClendon, <i>Metric families</i>	491
Carl Pomerance, <i>On multiply perfect numbers with a special property</i>	511
Mohan S. Putcha and Adil Mohamed Yaqub, <i>Polynomial constraints for finiteness of semisimple rings</i>	519
Calvin R. Putnam, <i>Hyponormal contractions and strong power convergence</i>	531
Douglas Conner Ravenel, <i>Multiplicative operations in BP^*BP</i>	539
Judith Roitman, <i>Attaining the spread at cardinals which are not strong limits</i>	545
Kazuyuki Saitô, <i>Groups of $*$-automorphisms and invariant maps of von Neumann algebras</i>	553
Brian Kirkwood Schmidt, <i>Homotopy invariance of contravariant functors acting on smooth manifolds</i>	559
Kenneth Barry Stolarsky, <i>The sum of the distances to N points on a sphere</i>	563
Mark Lawrence Teply, <i>Semiprime rings with the singular splitting property</i>	575
J. Pelham Thomas, <i>Maximal connected Hausdorff spaces</i>	581
Charles Thomas Tucker, II, <i>Concerning σ-homomorphisms of Riesz spaces</i>	585
Rangachari Venkataraman, <i>Compactness in abelian topological groups</i>	591
William Charles Waterhouse, <i>Basically bounded functors and flat sheaves</i>	597
David Westreich, <i>Bifurcation of operator equations with unbounded linearized part</i>	611
William Robin Zame, <i>Extendibility, boundedness and sequential convergence in spaces of holomorphic functions</i>	619