

Pacific Journal of Mathematics

**ON MULTIPLY PERFECT NUMBERS WITH A SPECIAL
PROPERTY**

CARL POMERANCE

ON MULTIPLY PERFECT NUMBERS WITH A SPECIAL PROPERTY

CARL POMERANCE

If m is a multiply perfect number and $m = p^a n$ where p is prime and $n \mid \sigma(p^a)$, then $m = 120, 672, 523776$, or m is an even perfect number.

1. Introduction. Suppose p is a prime, a, n are natural numbers, and

$$(1.1) \quad p^a \mid \sigma(n), \quad n \mid \sigma(p^a)$$

where σ is the sum of the divisors function. Then $1 = (p^a, \sigma(p^a)) = (p^a, n)$, so that $p^a n \mid \sigma(p^a)\sigma(n) = \sigma(p^a n)$; that is $p^a n$ is a multiply perfect number. In this paper we identify all multiply perfect numbers which arise in this fashion.

Let M be the set of Mersenne exponents, that is, $M = \{k: 2^k - 1 \text{ is prime}\}$. We shall prove

THEOREM 1.1. *If p, a, n is a solution of (1.1) where p is prime, then either*

$$(1.2) \quad p^a = 2^k - 1, \quad n = 2^{k-1} \quad \text{for some } k \in M$$

$$(1.3) \quad p^a = 2^{k-1}, \quad n = 2^k - 1 \quad \text{for some } k \in M$$

$$(1.4) \quad p^a = 2^3, \quad n = 15$$

$$(1.5) \quad p^a = 2^5, \quad n = 21$$

$$(1.6) \quad p^a = 2^9, \quad n = 1023.$$

COROLLARY 1.1. *If m is a multiply perfect number and $m = p^a n$ where p is prime and $n \mid \sigma(p^a)$, then $m = 120, 672, 523776$, or m is an even perfect number.*

Note that in [2] all solutions of (1.1) with $p^a = \sigma(n)$ are enumerated: they are (1.2) and (1.5). Hence in the proof of Theorem 1.1, we may assume $p^a < \sigma(n)$.

We recall that a natural number n is said to be super perfect if $\sigma(\sigma(n)) = 2n$. In [2] and Suryanarayana [8] it is shown that if n is super perfect and if either n or $\sigma(n)$ is a prime power, then $n = 2^{k-1}$ for $k \in M$. Here we will say n is *super multiply perfect* if $\sigma(\sigma(n))/n$ is an integer.

COROLLARY 1.2. *If n is super multiply perfect, and if n or $\sigma(n)$ is a prime power, then $n = 8, 21, 512$, or $n = 2^{k-1}$ for some $k \in M$.*

If p is a prime, denote by $\sigma_p(n)$ the sum of all those divisors of n which are powers of p . Then $\sigma_p(n) \mid \sigma(n)$.

COROLLARY 1.3. *If $n > 1$ and $n \mid \sigma_p(\sigma(n))$ for some prime p , then $p = 2$ and $n = 15, 21$, or 1023 or $p = 2^k - 1$ for some $k \in M$ and $n = 2^{k-1}$.*

We remark that in general the super multiply perfect numbers appear to be quite intractable. Partly complicating matters is that for every K , $\sigma(\sigma(n))/n \geq K$ on a set of density 1. Professor David E. Penney of the University of Georgia, in a computer search, found that there are exactly 37 super multiply perfect numbers ≤ 150000 . Of these, the only odd ones are 1, 15, 21, 1023, and 29127.

Recently, Guy and Selfridge [4], p. 104, published a proof of a stronger version of Theorem 1.1 for the special case $p = 2$.

2. Preliminaries. If n is a natural number, we let $\omega(n)$ be the number of distinct prime factors of n , and we let $\tau(n)$ be the number of natural divisors of n . If a, b are natural numbers with $(a, b) = 1$, we let $\text{ord}_a(b)$ be the least positive integer k for which $a \mid b^k - 1$. If p is a prime and x is a natural number, then $\sigma(p^x) = (p^{x+1} - 1)/(p - 1) < (p/(p - 1))p^x$.

THEOREM 2.1 (Bang [1]). *If p is a prime, a is a natural number, and $1 < d \mid a + 1$, then there is a prime $q \mid \sigma(p^a)$ with $\text{ord}_q(p) = d$, unless*

- (i) $p = 2$ and $d = 6$, or
- (ii) p is a Mersenne prime and $d = 2$.

COROLLARY 2.1.

$$\omega(\sigma(p^a)) \geq \begin{cases} \tau(a + 1) - 2, & \text{if } p = 2 \text{ and } 6 \mid a + 1 \\ \tau(a + 1), & \text{if } p > 2 \text{ is not Mersenne and} \\ & 2 \mid a + 1 \\ \tau(a + 1) - 1, & \text{otherwise.} \end{cases}$$

The following is a weaker form of a lemma from [2].

LEMMA 2.1. *Suppose p, q are primes with $q > 2$ and x, y, b, c are natural numbers with $\sigma(q^x) = p^y$ and $q^b \mid \sigma(p^c)$. Then $q^{b-1} \mid c + 1$.*

3. The start of the proof. Suppose p, a, n is a solution of (1.1) where p is prime. Then there are integers s, t with

$$\sigma(n) = sp^a, \quad \sigma(p^a) = tn.$$

As we remarked, we have already studied these equations in the case $s = 1$ (in [2]), so here we assume $s > 1$. We have

$$(3.1) \quad st = \frac{\sigma(p^a)}{p^a} \cdot \frac{\sigma(n)}{n},$$

Considering the unique prime factorization of n , we write n_1 for the product of those prime powers q^b for which $\sigma(q^b)$ is divisible by a prime $\neq p$, and we write n_2 for the product of those prime powers q^b for which $\sigma(q^b)$ is a power of p . Then $(n_1, n_2) = 1, n_1 n_2 = n$, and $\sigma(n_2)$ is a power of p . Let ω_i be the number of distinct odd prime factors of n_i for $i = 1, 2$. Let ω_3 be the number of distinct prime factors of t which do not divide n . Hence

$$(3.2) \quad \omega(\sigma(p^a)) = \omega(tn) = \begin{cases} \omega_1 + \omega_2 + \omega_3, & \text{if } n \text{ is odd} \\ 1 + \omega_1 + \omega_2 + \omega_3, & \text{if } n \text{ is even.} \end{cases}$$

We write

$$n_1 = 2^{k_1} \prod_{i=1}^{\omega_1} p_i^{a_i}, \quad n_2 = 2^{k_2} \prod_{i=1}^{\omega_2} q_i^{b_i}$$

where $k_1 k_2 = 0$ and the p_i and q_i are distinct odd primes.

4. The case $p > 2$. Since each $\sigma(q_i^{b_i})$ is a power of p , and since p is odd, we have each b_i even. Since also each $q_i^{b_i} \mid \sigma(p^a)$, Lemma 2.1 implies

$$\prod_{i=1}^{\omega_2} q_i \mid a + 1.$$

Suppose n is even. Then also $2 \mid a + 1$, so that $\tau(a + 1) \geq 2^{\omega_2 + 1}$. It follows from (3.2) and Corollary 2.1 that

$$(4.1) \quad \omega_1 + \omega_3 \geq 2^{\omega_2 + 1} - \omega_2 - 2.$$

Suppose $k_1 > 0$. Then $(\sigma(2^{k_1}), s) \geq 3$ and for

$$1 \leq i \leq \omega_1, \quad (\sigma(p_i^{a_i}), s) \geq 2.$$

Then $s \geq 3 \cdot 2^{\omega_1}$. Also every prime counted by ω_3 is odd, so $t \geq 3^{\omega_3}$. Hence from (3.1) we have

$$\begin{aligned}
3 \cdot \left(\frac{5}{4}\right)^{3\omega_1+4\omega_3} &< 3 \cdot 2^{\omega_1} \cdot 3^{\omega_3} \\
&\leq st = \frac{\sigma(p^a)}{p^a} \cdot \frac{\sigma(n)}{n} < \frac{p}{p-1} \cdot 2 \cdot \prod_{i=1}^{\omega_1} \frac{p_i}{p_i-1} \cdot \prod_{i=1}^{\omega_2} \frac{q_i}{q_i-1} \\
&\leq 3 \cdot \left(\frac{5}{4}\right)^{\omega_1+\omega_2}
\end{aligned}$$

so that

$$\omega_2 > 2\omega_1 + 4\omega_3 \geq 2(\omega_1 + \omega_3).$$

Hence (4.1) implies that

$$\omega_2 > 2^{\omega_2+2} - 2\omega_2 - 4$$

which fails for all $\omega_2 \geq 0$. This contradiction shows $k_1 = 0$.

Suppose $k_2 > 0$. Then $\sigma(2^{k_2})$ is a power of p , so that $\sigma(2^{k_2}) = p$ (Gerono [3]). Now $2 \mid a + 1$, so $2^{k_2+1} = \sigma(p) \mid \sigma(p^a)$. Hence $2 \mid t$, so that $t \geq 2 \cdot 3^{\omega_3}$. Also $(\sigma(p_i^{t_i}), s) \geq 2$, so $s \geq 2^{\omega_1}$. Hence

$$\begin{aligned}
\left(\frac{5}{4}\right)^{3\omega_1+4\omega_3} &< \frac{1}{2} st < \frac{p}{p-1} \cdot \prod \frac{p_i}{p_i-1} \cdot \prod \frac{q_i}{q_i-1} \leq \frac{3}{2} \left(\frac{5}{4}\right)^{\omega_1+\omega_2} \\
&< \left(\frac{5}{4}\right)^{\omega_1+\omega_2+2}
\end{aligned}$$

so that

$$(4.2) \quad \omega_2 > 2\omega_1 + 4\omega_3 - 2 \geq 2(\omega_1 + \omega_3) - 2.$$

It follows from (4.1) that

$$\omega_2 > 2^{\omega_2+2} - 2\omega_2 - 6,$$

which implies $\omega_2 \leq 1$. Then (4.2) implies $\omega_1 \leq 1$. Since $s > 1$ and $2 \mid t$, we have

$$4 \leq st < \frac{\sigma(2^{k_2})}{2^{k_2}} \cdot \frac{p}{p-1} \cdot \frac{p_1}{p_1-1} \cdot \frac{q_1}{q_1-1} < \frac{2pp_1q_1}{(p-1)(p_1-1)(q_1-1)}$$

so that $\max\{p, p_1, q_1\} = 13$. But $\sigma(2^{k_2}) = p$, so $k_2 \leq 2$. Then

$$4 < \frac{\sigma(2^2)}{2^2} \cdot \frac{3}{2} \cdot \frac{5}{4} \cdot \frac{7}{6} < 4,$$

so $k_2 = 0$.

Thus we have n odd, so $p^a n$ is an odd multiply perfect number. It follows from Hagis [5] and McDaniel [6] that

$$(4.3) \quad 1 + \omega_1 + \omega_2 = 1 + \omega(n) = \omega(p^a n) \geq 8.$$

From (3.2) and Corollary 2.1 we have

$$(4.4) \quad \omega_1 + \omega_3 \geq 2^{\omega_2} - \omega_2 - 1 .$$

Now $s \geq 2^{\omega_1}$, $t \geq 2^{\omega_3}$ so that

$$\begin{aligned} \left(\frac{5}{4}\right)^{3\omega_1+3\omega_3} &< st < \frac{p}{p-1} \cdot \prod \frac{p_i}{p_i-1} \cdot \prod \frac{q_i}{q_i-1} \\ &\leq \frac{3}{2} \left(\frac{5}{4}\right)^{\omega_1+\omega_2} < \left(\frac{5}{4}\right)^{\omega_1+\omega_2+2} . \end{aligned}$$

Hence

$$(4.5) \quad \omega_2 > 2\omega_1 + 3\omega_3 - 2 \geq 2(\omega_1 + \omega_3) - 2 ,$$

so that (4.4) implies

$$\omega_2 > 2^{\omega_2+1} - 2\omega_2 - 4 ,$$

which implies $\omega_2 \leq 2$. Then (4.5) implies $\omega_1 \leq 1$, contradicting (4.3).

5. The case $p = 2$. Since $\sigma(n_2)$ is a power of 2, it follows that n_2 is a product of distinct Mersenne primes (Sierpiński [7]), say

$$n_2 = \prod_{i=1}^{\omega_2} (2^{c_i} - 1)$$

where each c_i and $q_i = 2^{c_i} - 1$ is prime, and $c_1 < c_2 < \dots < c_{\omega_2}$.

Suppose $n_1 = 1$. Then s is a power of 2, say $s = 2^c$. Then

$$2^{c+a} = \sigma(n) = \sigma(n_2) = 2^{\sum c_i}$$

so that $c + a = \sum c_i$. But $2^{c_i} - 1 \mid \sigma(2^a)$, so $c_i \mid a + 1$. Since $c_1, c_2, \dots, c_{\omega_2}$ are distinct primes, $\prod c_i \mid a + 1$. Since $1 < s = 2^c$, we have $c \geq 1$. Hence $\prod c_i \leq a + 1 \leq \sum c_i$, so

$$\prod_{i=1}^{\omega_2} c_i - \sum_{i=1}^{\omega_2} c_i \leq 0 .$$

.....ls only for $\omega_2 = 1$, which gives solution (1.3).

We now assume $n_1 > 1$. Then s is divisible by an odd prime; in fact, $s \geq 3^{\omega_1} \geq 3$. Also t is odd, so $t \geq 3^{\omega_3}$. As above, $\prod c_i \mid a + 1$, so $\tau(a + 1) \geq 2^{\omega_2}$. Hence from (3.2) and Corollary 2.1 we have

$$(5.1) \quad \omega_1 + \omega_3 \geq 2^{\omega_2} - \omega_2 - 2 .$$

Also from (3.1) we have

$$\begin{aligned} \left(\frac{5}{4}\right)^{4\omega_1+4\omega_3-4} &< 3^{\omega_1-1} \cdot 3^{\omega_3} \leq \frac{1}{3} st = \frac{1}{3} \cdot \frac{\sigma(2^a)}{2^a} \cdot \frac{\sigma(n)}{n} \\ &< \frac{1}{3} \cdot 2 \cdot \prod \frac{p_i}{p_i-1} \cdot \prod \frac{q_i}{q_i-1} \leq \left(\frac{5}{4}\right)^{\omega_1+\omega_3-1}, \end{aligned}$$

so that

$$(5.2) \quad \omega_2 > 3\omega_1 + 4\omega_3 - 3 \geq 3(\omega_1 + \omega_3) - 3.$$

Then (5.1) implies

$$\omega_2 > 3 \cdot 2^{\omega_2} - 3\omega_2 - 9$$

so that $\omega_2 \leq 2$. Then from (5.2) and the fact that $\omega_1 \geq 1$, we have $\omega_1 = 1$, $\omega_3 = 0$, and $\omega_2 > 0$. Hence we have two choices for $\omega_1, \omega_2, \omega_3$: 1, 1, 0 and 1, 2, 0. Also since

$$5 > \frac{2}{1} \cdot \frac{3}{2} \cdot \frac{5}{4} \cdot \frac{7}{6} > \frac{\sigma(2^a)}{2^a} \cdot \frac{\sigma(n)}{n} = st$$

and since $s \geq 3$, $s \neq 4$, we have $s = 3$, $t = 1$.

Suppose $\omega_2 = 1$. Then $\sigma(2^a) = p_1^{a_1}(2^{c_1} - 1)$. Then c_1 is a proper divisor of $a + 1$. But $\omega(\sigma(2^a)) = 2$, so Corollary 2.1 implies $a + 1 = 6$ or $a + 1 = c_1^2$. The first choice gives $n = 63$, but $\sigma(63) \neq 3 \cdot 2^5$. Hence $a + 1 = c_1^2$. Then Theorem 2.1 implies $\text{ord}_{p_1}(2) = c_1^2$, so that $p_1 \equiv 1 \pmod{c_1^2}$. If $c_1 \geq 3$, then $p_1 \geq 19$, $q_1 = 2^{c_1} - 1 \geq 7$, so that

$$3 = st < \frac{2}{1} \cdot \frac{7}{6} \cdot \frac{19}{18} < 3,$$

a contradiction. Hence $c_1 = 2$, $a + 1 = 4$, $n = 15$, and we have solution (1.4).

Our last case is $\omega_2 = 2$. Then $\sigma(2^a) = p_1^{a_1}(2^{c_1} - 1)(2^{c_2} - 1)$, so that $c_1 c_2 \mid a + 1$. Now $\omega(\sigma(2^a)) = 3$, so that Corollary 2.1 implies $c_1 c_2 = a + 1$, where $c_1 c_2 \neq 6$. We also have $\sigma(p_1^{a_1}(2^{c_1} - 1)(2^{c_2} - 1)) = 3 \cdot 2^a$. Then $\sigma(p_1^{a_1})$ is 3 times a power of 2. Now $\sigma(p_1^{a_1}) \neq 3$, so $\sigma(p_1^{a_1})$ is even. Hence $2 \mid a_1 + 1$. Now Theorem 2.1 implies $\text{ord}_{p_1}(2) = c_1 c_2$, a composite number. Hence p_1 is not Mersenne. Also, $p_1 \equiv 1 \pmod{c_1 c_2}$. From Corollary 2.1 and the fact that $\omega(\sigma(p_1^{a_1})) = 2$ we have $a_1 = 1$. Hence for some d we have $p_1 = 3 \cdot 2^d - 1$. If $c_1 > 2$, then $q_1 = 2^{c_1} - 1 \geq 7$, $q_2 = 2^{c_2} - 1 \geq 31$, $p_1 \geq 2c_1 c_2 + 1 \geq 31$. Then

$$3 = st < \frac{2}{1} \cdot \frac{31}{30} \cdot \frac{31}{30} \cdot \frac{7}{6} < 3,$$

so that we must have $c_1 = 2$. Then

$$2^{2c_2} - 1 = (3 \cdot 2^d - 1)(2^2 - 1)(2^{c_2} - 1)$$

where $c_2 \geq 3$. Looking at this equation mod 8, we obtain $3 \cdot 2^d - 1 \equiv 2^2 - 1 \pmod{8}$. Hence $d = 2$, $p_1 = 11$. Then $a + 1 = 2c_2 = \text{ord}_{p_1}(2) = 10$. This gives solution (1.6).

REFERENCES

1. A. S. Bang, *Taltheoretiske Undersøgelser*, Tidsskrift Math., 5 IV (1886), 70-80, 130-137.
2. G. G. Dandapat, J. L. Hunsucker, and C. Pomerance, *Some new results on odd perfect numbers*, Pacific J. Math., to appear.
3. C. G. Gerono, *Note sur la résolution en nombres entiers et positifs de l'équation $x^m = y^n + 1$* , Nouv. Ann. Math., (2) 9 (1870), 469-471; 10 (1871), 204-206.
4. R. K. Guy and J. L. Selfridge, *What drives an aliquot sequence?*, Math. Comp., 29 (1975), 101-107.
5. P. Hagsis, Jr., *Every odd perfect number has at least eight prime factors* (preliminary report), Not. Amer. Math. Soc. 22 (1975), A-60.
6. W. McDaniel, *On odd multiply perfect numbers*, Boll. Un. Mat. Ital., (4) 3 (1970), 185-190.
7. W. Sierpiński, *Sur les nombres dont la somme des diviseurs est un puissance du nombre 2*, The Golden Jubilee Commemoration Volume (1958-9), Calcutta Math. Soc., 7-9.
8. D. Suryanarayana, *There is no odd super perfect number of the form p^{2a}* , Elem. Math., 28 (1973), 148-150.

Received February 7, 1975.

UNIVERSITY OF GEORGIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

| | |
|---|-----|
| Norman Larrabee Alling, <i>On Cauchy's theorem for real algebraic curves with boundary</i> | 315 |
| Daniel D. Anderson, <i>A remark on the lattice of ideals of a Prüfer domain</i> | 323 |
| Dennis Neal Barr and Peter D. Miletta, <i>A necessary and sufficient condition for uniqueness of solutions to two point boundary value problems</i> | 325 |
| Ladislav Beran, <i>On solvability of generalized orthomodular lattices</i> | 331 |
| L. Carlitz, <i>A three-term relation for some sums related to Dedekind sums</i> | 339 |
| Arthur Herbert Copeland, Jr. and Albert Oscar Shar, <i>Images and pre-images of localization maps</i> | 349 |
| G. G. Dandapat, John L. Hunsucker and Carl Pomerance, <i>Some new results on odd perfect numbers</i> | 359 |
| M. Edelstein and L. Keener, <i>Characterizations of infinite-dimensional and nonreflexive spaces</i> | 365 |
| Francis James Flanigan, <i>On Levi factors of derivation algebras and the radical embedding problem</i> | 371 |
| Harvey Friedman, <i>Provable equality in primitive recursive arithmetic with and without induction</i> | 379 |
| Joseph Braucher Fugate and Lee K. Mohler, <i>The fixed point property for tree-like continua with finitely many arc components</i> | 393 |
| John Norman Ginsburg and Victor Harold Saks, <i>Some applications of ultrafilters in topology</i> | 403 |
| Arjun K. Gupta, <i>Generalisation of a "square" functional equation</i> | 419 |
| Thomas Lee Hayden and Frank Jones Massey, <i>Nonlinear holomorphic semigroups</i> | 423 |
| V. Kannan and Thekkedath Thrivikraman, <i>Lattices of Hausdorff compactifications of a locally compact space</i> | 441 |
| J. E. Kerlin and Wilfred Dennis Pepe, <i>Norm decreasing homomorphisms between group algebras</i> | 445 |
| Young K. Kwon, <i>Behavior of Φ-bounded harmonic functions at the Wiener boundary</i> | 453 |
| Richard Arthur Levaro, <i>Projective quasi-coherent sheaves of modules</i> | 457 |
| Chung Lin, <i>Rearranging Fourier transforms on groups</i> | 463 |
| David Lowell Lovelady, <i>An asymptotic analysis of an odd order linear differential equation</i> .. | 475 |
| Jerry Malzan, <i>On groups with a single involution</i> | 481 |
| J. F. McClendon, <i>Metric families</i> | 491 |
| Carl Pomerance, <i>On multiply perfect numbers with a special property</i> | 511 |
| Mohan S. Putcha and Adil Mohamed Yaqub, <i>Polynomial constraints for finiteness of semisimple rings</i> | 519 |
| Calvin R. Putnam, <i>Hyponormal contractions and strong power convergence</i> | 531 |
| Douglas Conner Ravenel, <i>Multiplicative operations in BP^*BP</i> | 539 |
| Judith Roitman, <i>Attaining the spread at cardinals which are not strong limits</i> | 545 |
| Kazuyuki Saitô, <i>Groups of *-automorphisms and invariant maps of von Neumann algebras</i> | 553 |
| Brian Kirkwood Schmidt, <i>Homotopy invariance of contravariant functors acting on smooth manifolds</i> | 559 |
| Kenneth Barry Stolarsky, <i>The sum of the distances to N points on a sphere</i> | 563 |
| Mark Lawrence Teply, <i>Semiprime rings with the singular splitting property</i> | 575 |
| J. Pelham Thomas, <i>Maximal connected Hausdorff spaces</i> | 581 |
| Charles Thomas Tucker, II, <i>Concerning σ-homomorphisms of Riesz spaces</i> | 585 |
| Rangachari Venkataraman, <i>Compactness in abelian topological groups</i> | 591 |
| William Charles Waterhouse, <i>Basically bounded functors and flat sheaves</i> | 597 |
| David Westreich, <i>Bifurcation of operator equations with unbounded linearized part</i> | 611 |
| William Robin Zame, <i>Extendibility, boundedness and sequential convergence in spaces of holomorphic functions</i> | 619 |