

Pacific Journal of Mathematics

MULTIPLICATIVE OPERATIONS IN BP^*BP

DOUGLAS CONNER RAVENEL

MULTIPLICATIVE OPERATIONS IN BP^*BP

DOUGLAS C. RAVENEL

One of the present computational difficulties in complex cobordism theory is the lack of a known algebra splitting of BP^*BP , the algebra of stable cohomology operations for the Brown-Peterson cohomology theory, analogous to the splitting isomorphism

$$MU^*MU \approx MU^*(pt) \otimes S$$

where S is the Landweber-Novikov algebra. S has the added advantage of being a cocommutative Hopf algebra over Z .

This paper does not remove this difficulty, but we will show that the monoid of multiplicative operations in BP^*BP , (i.e. those operations which induce ring endomorphisms on BP^*X for any space X), which we will denote by $\Gamma(BP)$, has a submonoid analogous to the monoid of multiplicative operations in S .

The latter monoid is known (see Morava [3]) to be isomorphic to the group of formal power series $f(x)$ over Z such that $f(0) = 0$ and $f'(0) = 1$ and where the group operation is composition of power series.

For the basic properties of MU^*MU and BP^*BP , see Adams [1], especially §§ 11 and 16.

The main construction of this paper was inspired by the work of Honda ([3]) although none of his results are needed here. I am grateful to Jack Morava for bringing Honda's work to my attention.

Before stating our main result we must review the description of $\Gamma(BP)$ implicit in [1] § 16. An operation $\alpha \in \Gamma(BP)$ is characterized by its action on the canonical generator $z \in BP^*(CP^\infty) \cong \pi_*BP[[z]]$. It is shown that $\alpha(z)$ is a power series $f(z)$ over π_*BP where

$$(1) \quad f^{-1}(z) = z + {}_{\mu}t_1 z^p + {}_{\mu}t_2 z^{p^2} + {}_{\mu}t_3 z^{p^3} + {}_{\mu} \dots$$

where $+_{\mu}$ denotes the sum in the formal group defined over π_*BP and $t_i \in \pi_*BP$. (This formula appears on page 96 of [1].)

The action of α on π_*BP can be read off from (1). Let $l_n \in \pi_{2p^n-2}BP \otimes Q$ be defined by $\log^{BP} x = \sum_{n=0} l_n x^{p^n}$ (l_n is the m_{p^n-1} of [1]). Then we have

$$(2) \quad \alpha(l_n) = \sum_{0 \leq i \leq n} l_n t_{n-i}^{p^i}$$

and this formula also characterizes α .

In other words $\Gamma(BP)$ is an infinite dimensional affine space over

π_*BP with coordinates t_i with a composition law which can be read off from (2). The difficulty mentioned above is that the subset of elements with integer coordinates is not a submonoid.

The main object of this paper is to construct new coordinates s_i of $\Gamma(BP)$ such that the subset of elements with coordinates in \mathbb{Q}_p (the set of rational numbers with denominators prime to p) is a submonoid which will be denoted by $\gamma(BP)$. Moreover this submonoid is isomorphic to a direct product of countably many copies of \mathbb{Q}_p , although this isomorphism is somewhat accidental in a sense to be described below. The s_i are not unique as they depend upon a choice of generators of π_*BP . What's worse, they are not algebraic functions of the t_i , so they do not lead to a splitting of BP^*BP .

Let $\{v_j \in \pi_*BP\}$ be a set of generators. Define a ring endomorphism σ of π_*BP by $v_j^\sigma = v_j^p$, extending σ linearly over all of π_*BP . Then our main result is

THEOREM. *For every sequence of coordinates $\{t_n \in \pi_*BP\}$ there exists $\{s_n \in \pi_*BP\}$ such that*

$$(3) \quad \sum_{0 \leq i \leq n} l_i t_n^{p^i} = \sum_{0 \leq i \leq n} l_i s_n^{p^i}$$

for every $n > 0$, and for every $\{s_n\}$ there exist $\{t_n\}$ satisfying the same conditions.

COROLLARY. *Let $\alpha', \alpha'' \in \Gamma(BP)$ have coordinates $s'_n, s''_n \in \mathbb{Q}_p$ respectively. Then the coordinates s'''_n of $\alpha''' = \alpha' \cdot \alpha''$ are given by*

$$s'''_n = \sum_{i+j=n} s'_i s''_j$$

i.e. if we define power series $s'(x), s''(x), s'''(x)$ by $s'(x) = 1 + \sum_{n>0} s'_n x^n$, etc., then $s'''(x) = s'(x)s''(x)$.

The corollary follows from (3) by direct computation, remembering that σ fixes $\mathbb{Q}_p \subset \pi_*BP$.

To prove the theorem we need an analogue of (1) involving the new coordinates s_n , and some elementary properties of formal groups beginning with

LEMMA A. *For every $r \in \mathbb{Q}_p$ there is a power series $[r](z) \in \pi_*BP[[z]]$ such that*

- (a) $[1](z) = z$
- (b) $[r'](z) +_\mu [r''](z) = [r' + r''](z)$
- (c) $[r']([r''](z)) = [r'r''](z)$
- (d) $[r](z) \equiv rz \text{ modulo } (z^2)$.

Proof. If r is a positive integer define $[r](z)$ inductively by $[r](z) = z +_{\mu} [r-1](z)$; the existence of $[-1](z)$ is one of the formal group axioms; and if r is an integer prime to p , define $[1/r](z)$ to be the formal inverse of $[r](z)$. Then (c) enables us to define $[r](z)$ for any $r \in \mathbb{Q}_p$.

Now let $M = (m_1, m_2, m_3, \dots)$ be a sequence of nonnegative integers, of which all but a finite number are zero, and let

$$v^M = \prod_{j>0} v_j^{m_j} \in \pi_*BP.$$

Then the v^M 's form a \mathbb{Q}_p -basis of π_*BP and we can write

$$s_i = \sum s_{i,M} v^M$$

with $s_{i,M} \in \mathbb{Q}_p$ and the sum ranging over all M . With this notation we have

LEMMA B. *Formula (3) is equivalent to*

$$(4) \quad \sum_{n>0}^{\mu} t_n z^{p^n} = \sum_{n,M}^{\mu} [s_{n,M}] (v^M z^{p^n})$$

where \sum^{μ} denotes the formal group sum.

Proof. Multiplying both sides of (3) by z^{p^n} and summing over all positive n gives

$$\sum_{n>0} \log t_n z^{p^n} = \sum_{n,M} s_{n,M} \log v^M z^{p^n}$$

which is equivalent to (4).

LEMMA C. *For $r \in \mathbb{Q}_p$, there exist $r_n \in \pi_*BP$ with $r_0 = r$ such that for any $u \in BP^*CP^{\infty}$*

$$(5) \quad [r](u) = \sum_{n \geq 0}^{\mu} r_n u^{p^n}.$$

Proof. We will first show that there exist $r_{(k)} \in \pi_*BP$ such that

$$(6) \quad [r](u) = \sum_{k>0}^{\mu} r_{(k)} u^k$$

and then show that $r_{(k)} = 0$ unless $k = p^n$. The proof of the former statement is by induction on k . Let $r_{(1)} = r$ and suppose we have found $r_{(1)}, r_{(2)}, \dots, r_{(m)}$ such that

$$[r](u) = \sum_{k=1}^m r_{(k)} u^k \quad \text{modulo } (u^{m+1}).$$

Then we can take $r_{(m+1)}$ to be the coefficient of u^{m+1} in

$$[r](u) - \sum_{k=1}^{\mu} r_{(k)} u^k$$

and the first statement follows. Taking the log of both sides of (6) we have

$$r \log u = \sum_{k>0} \log r_{(k)} u^k$$

i.e.

$$r \sum_{n \geq 0} l_n u^{p^n} = \sum_{\substack{k>0 \\ j \geq 0}} l_j r_{(k)}^{p^j} u^{k p^j}$$

Equating coefficients of u^i yields the lemma.

LEMMA D. For $a', a'' \in \pi_*BP$, there exist $a_n \in \pi_*BP$ for $n \geq 0$ with $a_0 = a' + a''$ such that for any $u \in BP^*CP^\infty$

$$(7) \quad a'u + {}_{\mu} a''u = \sum_{n \geq 0} a_n u^{p^n}.$$

The proof of Lemma D is similar to that of Lemma C and is left to the reader.

We are now ready to prove the theorem via (4). Given $\{t_n\}$ we will construct $\{s_n\}$ by induction on n , beginning with $s_1 = t_1$. Suppose we have found s_1, s_2, \dots, s_m such that

$$(8) \quad \sum_{n>0} t_n z^{p^n} \equiv \sum_{\substack{0 < n \leq m \\ M}} [s_{n,M}] (v^M z^{p^n} \text{ modulo } (z^{1+p^n}))$$

By repeated application of Lemmas C and D we can find $w_n \in \pi_*BP$ such that the right hand side of (8) is equal to $\sum_{n>0} w_n z^{p^n}$ and we know $w_n = t_n$ for $n \leq m$. This allows us to set $s_{m+1} = t_{m+1} - w_{m+1}$, and the first half of the theorem is proved. Similarly if we are given $\{s_n\}$, let $t_1 = s_1$ and suppose we have found t_n for $n \leq m$. Then we can set $t_{m+1} = w_{m+1} + s_{m+1}$, and the theorem is proved.

Now I will describe the sense in which the commutativity of $\gamma(BP)$ is accidental. If this paper were being written for number theorists rather than topologists, we would replace π_*BP by $\pi_*BP \otimes_{\mathbb{Q}_p} W(k)$, where $W(k)$ is the Witt ring of a finite field k of characteristic p . Lemma A would go through only for $r \in W(\mathbb{F}_p)$, the p -adic integers. The ring endomorphism σ when restricted to $W(k)$ would be the lifting of the Frobenius automorphism on k . The corollary (with $s'_n, s''_n \in W(k)$) would then read

$$s'''_n = \sum_{i+j=n} s'_i (s''_j)^{\sigma^i}$$

The resulting group $\gamma_k(BP)$ would not be abelian if k has more than p elements.

REFERENCES

1. J. F. Adams, *Quillen's Work on Formal Groups and Complex Cobordism*, University of Chicago Lecture Notes, 1970.
2. T. Honda, *The theory of commutative formal groups*, J. Math. Soc. Japan, **22** (1970), 213-246.
3. J. Morava, *Structure Theorems for Cobordism Comodules*, (to appear).

Received December 4, 1974. Research partially supported by NSF Grant GP-32843.

COLUMBIA UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 57, No. 2

February, 1975

Norman Larrabee Alling, <i>On Cauchy's theorem for real algebraic curves with boundary</i>	315
Daniel D. Anderson, <i>A remark on the lattice of ideals of a Prüfer domain</i>	323
Dennis Neal Barr and Peter D. Miletta, <i>A necessary and sufficient condition for uniqueness of solutions to two point boundary value problems</i>	325
Ladislav Beran, <i>On solvability of generalized orthomodular lattices</i>	331
L. Carlitz, <i>A three-term relation for some sums related to Dedekind sums</i>	339
Arthur Herbert Copeland, Jr. and Albert Oscar Shar, <i>Images and pre-images of localization maps</i>	349
G. G. Dandapat, John L. Hunsucker and Carl Pomerance, <i>Some new results on odd perfect numbers</i>	359
M. Edelstein and L. Keener, <i>Characterizations of infinite-dimensional and nonreflexive spaces</i>	365
Francis James Flanigan, <i>On Levi factors of derivation algebras and the radical embedding problem</i>	371
Harvey Friedman, <i>Provable equality in primitive recursive arithmetic with and without induction</i>	379
Joseph Braucher Fugate and Lee K. Mohler, <i>The fixed point property for tree-like continua with finitely many arc components</i>	393
John Norman Ginsburg and Victor Harold Saks, <i>Some applications of ultrafilters in topology</i>	403
Arjun K. Gupta, <i>Generalisation of a "square" functional equation</i>	419
Thomas Lee Hayden and Frank Jones Massey, <i>Nonlinear holomorphic semigroups</i>	423
V. Kannan and Thekkedath Thrivikraman, <i>Lattices of Hausdorff compactifications of a locally compact space</i>	441
J. E. Kerlin and Wilfred Dennis Pepe, <i>Norm decreasing homomorphisms between group algebras</i>	445
Young K. Kwon, <i>Behavior of Φ-bounded harmonic functions at the Wiener boundary</i>	453
Richard Arthur Levaro, <i>Projective quasi-coherent sheaves of modules</i>	457
Chung Lin, <i>Rearranging Fourier transforms on groups</i>	463
David Lowell Lovelady, <i>An asymptotic analysis of an odd order linear differential equation</i>	475
Jerry Malzan, <i>On groups with a single involution</i>	481
J. F. McClendon, <i>Metric families</i>	491
Carl Pomerance, <i>On multiply perfect numbers with a special property</i>	511
Mohan S. Putcha and Adil Mohamed Yaqub, <i>Polynomial constraints for finiteness of semisimple rings</i>	519
Calvin R. Putnam, <i>Hyponormal contractions and strong power convergence</i>	531
Douglas Conner Ravenel, <i>Multiplicative operations in BP^*BP</i>	539
Judith Roitman, <i>Attaining the spread at cardinals which are not strong limits</i>	545
Kazuyuki Saitô, <i>Groups of $*$-automorphisms and invariant maps of von Neumann algebras</i>	553
Brian Kirkwood Schmidt, <i>Homotopy invariance of contravariant functors acting on smooth manifolds</i>	559
Kenneth Barry Stolarsky, <i>The sum of the distances to N points on a sphere</i>	563
Mark Lawrence Teply, <i>Semiprime rings with the singular splitting property</i>	575
J. Pelham Thomas, <i>Maximal connected Hausdorff spaces</i>	581
Charles Thomas Tucker, II, <i>Concerning σ-homomorphisms of Riesz spaces</i>	585
Rangachari Venkataraman, <i>Compactness in abelian topological groups</i>	591
William Charles Waterhouse, <i>Basically bounded functors and flat sheaves</i>	597
David Westreich, <i>Bifurcation of operator equations with unbounded linearized part</i>	611
William Robin Zame, <i>Extendibility, boundedness and sequential convergence in spaces of holomorphic functions</i>	619