

Pacific Journal of Mathematics

**EXTENSION FUNCTIONS FOR RANK 2, TORSION FREE
ABELIAN GROUPS**

ERIC FRIEDLANDER

EXTENSION FUNCTIONS FOR RANK 2, TORSION FREE ABELIAN GROUPS

ERIC M. FRIEDLANDER

The set of isomorphism classes of rank 2, torsion free abelian groups with a pure subgroup isomorphic to a given rank 1 group is shown to be in natural 1-1 correspondence with the set of pairs consisting of a quotient type and a type of an extension function. In terms of these invariants, necessary and sufficient conditions are determined for such a group to be homogeneous or to admit a pure cyclic subgroup. Moreover, this 1-1 correspondence has an explicit inverse, so that examples are readily obtained.

Our method is to combine the Korosh-Malcev-Derry matrix classification (see [2] for this and other well known aspects of the theory of abelian groups which we employ) with some elementary observations concerning abelian extensions of rank 1, torsion free abelian groups. We begin §1 by identifying $\text{Ext}_Z(X, Y)$ for rank 1, torsion free abelian groups X and Y in terms of "extension functions." A Korosh-Malcev-Derry matrix sequence for the total group of such an extension is readily given in terms of an extension function. Moreover, an extension function explicitly determines a subgroup of $Q \oplus Q$. We obtain an explicit necessary condition for two extension functions to determine isomorphic total groups, as well as express the Korosh-Malcev-Derry matrix conditions in terms of extension functions.

In §2, we explicate the 1-1 correspondence asserted above. We then "list" all homogeneous, rank 2, torsion free abelian groups of a given type. We also determine in terms of our invariants whether or not a rank 2, torsion free abelian group admits a pure cyclic subgroup.

We gratefully acknowledge many helpful conversations with C. Miller. Moreover, the referee's suggestion of generalizing an earlier version of this work proved most valuable. We refer the interested reader to ([1]) for a detailed study of rank 2, torsion free abelian groups up to quasi-isomorphism and to ([2]) for a survey of the literature on torsion free abelian groups.

1. Extension functions. A sequence (a_1, \dots, a_n, \dots) with each a_i an extended nonnegative integer, $0 \leq a_i \leq \infty$, is called a characteristic. Two characteristics, (a_1, \dots, a_n, \dots) and (b_1, \dots, b_n, \dots) , are said to have the same type if and only if $\sum_i (a_i - b_i)^2 < \infty$. If A is a torsion free abelian group, then the characteristic of any nonzero element x in A , $\text{char}(x) = (a_1, \dots, a_n, \dots)$, is defined by

$a_i = \text{char}_i(x) = \sup\{k: p_i^{-k}x \text{ in } A\}$, where p_i denotes the i th prime. Two torsion free, rank 1 abelian groups, X and Y , are isomorphic if and only if the type of $\text{char}(x)$ equals the type of $\text{char}(y)$ for any nonzero elements x and y of X and Y respectively.

We first explicitly compute $\text{Ext}_Z^1(X, Y)$ for any pair of rank 1, torsion free abelian groups X and Y .

PROPOSITION 1.1. *Let X and Y be rank 1, torsion free abelian groups and let x in X and y in Y be nonzero elements. Let $(a_1, \dots, a_n, \dots) = \text{char}(x)$ and $(b_1, \dots, b_n, \dots) = \text{char}(y)$. Then there exists an isomorphism*

$$\varphi(X, x; Y, y): \text{Ext}_Z(X, Y) \rightarrow \text{coker}(\theta_y: Y \rightarrow \prod' \mathbf{Z}/p_i^{a_i})$$

where \prod' denotes the product over all i with $b_i < \infty$, where $\mathbf{Z}/p_i^{a_i}$ denotes the p_i -adic integers \mathbf{Z}_{p_i} if $a_i = \infty$, and where $\theta_y(m/n \cdot y) = (m/n \cdot p_1^{b_1}, \dots, m/n \cdot p_n^{b_n}, \dots)$. Moreover, $\varphi(X, x; Y, y)$ is natural for maps $(X', x') \rightarrow (X, x)$ and maps $(Y', y') \rightarrow (Y, y)$.

Proof. Let $Y \rightarrow Q$ be defined by sending y to 1. Since Q and Q/Y are divisible abelian groups, $\text{Ext}_Z(X, Y)$ is naturally isomorphic to $\text{coker}(\text{Hom}(X, Q) \rightarrow \text{Hom}(X, Q/Y))$. Applying the serpent lemma to the following map of short exact sequences

$$\begin{array}{ccccccc} 0 \rightarrow & \text{Hom}(X/\mathbf{Z} \cdot x, Q) & \rightarrow & \text{Hom}(X, Q) & \rightarrow & \text{Hom}(\mathbf{Z} \cdot x, Q) & \rightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \\ 0 \rightarrow & \text{Hom}(X/\mathbf{Z} \cdot x, Q/Y) & \rightarrow & \text{Hom}(X, Q/Y) & \rightarrow & \text{Hom}(\mathbf{Z} \cdot x, Q/Y) & \rightarrow 0 \end{array}$$

we conclude that $\text{Ext}_Z(X, Y) \xrightarrow{\sim} \text{coker}(Y \rightarrow \text{Hom}(X/\mathbf{Z} \cdot x, Q/Y))$.

Let $\mu(p, a)$ denote the cyclic subgroup of Q/\mathbf{Z} generated by $1/p^a$ (let $\mu(p, \infty) = \lim_n \mu(p, n)$). Then

$$X/\mathbf{Z} \cdot x \xrightarrow{\sim} \bigoplus \mu(p_i, a_i) \quad \text{and} \quad Q/Y \xrightarrow{\sim} \bigoplus' \mu(p_i, \infty)$$

where the latter sum is taken over all i with $b_i < \infty$. Thus

$$\text{Hom}(X/\mathbf{Z} \cdot x, Q/Y) \xrightarrow{\sim} \text{Hom}(\bigoplus \mu(p_i, a_i), \bigoplus' \mu(p_i, \infty)) \xrightarrow{\sim} \prod' \mathbf{Z}/p_i^{a_i}.$$

The map $Y \rightarrow \text{Hom}(X/\mathbf{Z} \cdot x, Q/Y)$ is easily checked to send y to the sequence $\{p_i^{b_i}\}$ under these identifications.

To check the naturality of $\varphi(X, x; Y, y)$ for a map $f: (X', x') \rightarrow (X, x)$, one must verify the commutativity of the following square:

$$\begin{array}{ccc}
 \text{Ext}_{\mathbf{Z}}(X, Y) & \xrightarrow{\varphi(X,x;Y,y)} & \text{coker}(\theta_y: Y \rightarrow \Pi' \mathbf{Z}/p^a) \\
 \downarrow \text{Ext}_{\mathbf{Z}}(f, \mathbf{Z}) & & \downarrow \pi \\
 \text{Ext}_{\mathbf{Z}}(X', Y) & \xrightarrow{\varphi(\mathbf{Z}',x';Y,y)} & \text{coker}(\theta_y: Y \rightarrow \Pi' \mathbf{Z}/p^a)
 \end{array}$$

where π is induced by the factor by factor projection map. This verification, and the corresponding verification for naturality with respect to maps $g: (Y', y') \rightarrow (Y, y)$, are routine.

For a given (Y, y) , we call f in $\Pi' \mathbf{Z}/p^a$ an “extension function” for the corresponding extension $\varphi(X, x; Y, y)^{-1}(\bar{f})$ in $\text{Ext}_{\mathbf{Z}}(X, Y)$, where \bar{f} is the image of f in $\text{coker}(\theta_y)$ and (X, x) is any torsion free, rank 1 abelian group X with nonzero element x satisfying $\text{char}(x) = (a_1, \dots, a_n, \dots)$.

Observe that in the particular case that Y is free and y is a generator, Proposition 1.1 asserts that

$$\text{Ext}_{\mathbf{Z}}(X, \mathbf{Z}) \rightarrow (\Pi \mathbf{Z}/p^a)/\Delta(\mathbf{Z})$$

so that two extension functions determine isomorphic extensions if and only if they differ by a constant.

We recall that if A is a finite rank, torsion free abelian group, then $A \otimes \mathbf{Z}_p$ is a direct sum of \mathbf{Z}_p 's and Q_p 's for every prime p , since $\text{Ext}_{\mathbf{Z}}^1(Q_p, \mathbf{Z}_p) = 0$ (where \mathbf{Z}_p denotes the p -adic integers and Q_p denotes the field of p -adic numbers). We say that $\{x_1, \dots, x_n\}$ in $A \otimes \mathbf{Z}_p$ is a “basis” if $A \otimes \mathbf{Z}_p$ is the internal direct sum of the pure \mathbf{Z}_p modules generated by the x_i 's:

$$A \otimes \mathbf{Z}_p \xrightarrow{\sim} \mathbf{Z}_p \cdot x_{i_1} \oplus \dots \oplus \mathbf{Z}_p \cdot x_{i_k} \oplus Q_p \cdot x_{i_{k+1}} \oplus \dots \oplus Q_p \cdot x_{i_n}$$

A “matrix sequence” for a finite rank, torsion free abelian group A is a sequence of matrices $\{M_i\}$ expressing a given basis for $A \otimes Q$ in terms of bases for $A \otimes \mathbf{Z}_{p_i}$ thus, if A is rank 2, $\left\{ M_i \begin{pmatrix} 1 \\ 0 \end{pmatrix}, M_i \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}$ is the given basis for $A \otimes Q$ expressed in terms of a basis for $A \otimes \mathbf{Z}_{p_i}$. Such a matrix sequence determines the isomorphism class of A .

In the following proposition, we determine matrix sequences in terms of extension functions.

PROPOSITION 1.2. *Let X and Y be rank 1, torsion free abelian groups, let x in X and y in Y be nonzero elements, and let $\text{char}(x) =$*

(a_1, \dots, a_n, \dots) and $\text{char}(y) = (b_1, \dots, b_n, \dots)$. For a given extension function f in $\Pi' \mathbf{Z}/p_i^{a_i}$, let

$$0 \rightarrow Y \rightarrow A_f \rightarrow X \rightarrow 0$$

be the extension $\varphi(X, x; Y, y)^{-1}(\bar{f})$ in $\text{Ext}_{\mathbf{Z}}^1(X, Y)$. Then A_f has a matrix sequence

$$\left\{ \begin{pmatrix} p_i^{b_i} & -f_i \\ 0 & p_i^{a_i} \end{pmatrix} \right\}$$

where $0 \leq f_i < p_i^{a_i}$ with $f(i) = f_i$ in $\mathbf{Z}/p_i^{a_i}$ if $a_i, b_i < \infty$, where $f_i = 0$ and $p_i^{b_i}$ designates 1 if $b_i = \infty$, and where $f_i = f(i)$ in \mathbf{Z}_{p_i} and $p_i^{a_i}$ designates 1 if $a_i = \infty$.

Proof. The extension $\varphi(X, x; Y, y)^{-1}(\bar{f})$ is obtained by pull-back from $0 \rightarrow Y \rightarrow Q \rightarrow Q/Y \rightarrow 0$ via the composition $X \rightarrow X/\mathbf{Z} \cdot x \xrightarrow{\bar{f}} Q/Y$. For notational convenience, we view $A_f = Q \times_{Q/Y} X$ as a subgroup of $Q \oplus X$, containing $(1, 0) = y$ and $(0, x) = x$.

For each i with $b_i = \infty$, $\{y, p_i^{-a_i}y\}$ in A_f is a basis for $A_f \otimes \mathbf{Z}_{p_i}$. For each i with $b_i, a_i < \infty$, $\{p_i^{-b_i}y, f_i p_i^{-a_i-b_i}y + p_i^{-a_i}y\}$ in A_f is a basis for $A_f \otimes \mathbf{Z}_{p_i}$. For each i with $b_i < \infty$ and $a_i = \infty$, $\{f_{i,k} p_i^{-k-b_i}y + p_i^{-k}y\}$ is in A_f for all $k > 0$ where $f_i \equiv f_{i,k} \pmod{p_i^k}$, so that $\{p_i^{-b_i}y, f_i p_i^{-b_i}y + x\}$ is a basis for $A_f \otimes \mathbf{Z}_{p_i}$. Then $\begin{pmatrix} p_i^{b_i} & -f_i \\ 0 & p_i^{a_i} \end{pmatrix}$ is the matrix expressing the basis $\{y, x\}$ of $A_f \otimes Q$ in terms of the above basis for $A_f \otimes \mathbf{Z}_{p_i}$.

We next determine in terms of generators a standard model for the rank 2, torsion free abelian group A_f defined by the extension function f .

PROPOSITION 1.3. *Let X and Y be rank 1, torsion free abelian groups, let x in X and y in Y be nonzero elements, and let $\text{char}(x) = (a_1, \dots, a_n, \dots)$ and $\text{char}(y) = (b_1, \dots, b_n, \dots)$. For a given extension function f in $\Pi' \mathbf{Z}/p_i^{a_i}$, the map*

$$i(y, x): A_f = Q \times_{Q/Y} X \rightarrow Q \oplus Q$$

sending y to $(1, 0)$ and x to $(0, 1)$ is an isomorphism onto that subgroup $A(f)$ of $Q \oplus Q$ generated by $(p_i^{b_i}, 0)$, $(f_i p_i^{-a_i-b_i}, p_i^{-a_i})$ whenever $a_i, b_i < \infty$; $(p_i^{-k}, 0)$, $(0, p_i^{-a_i})$ for all $k > 0$, whenever $a_i < \infty$, $b_i = \infty$; $(p_i^{-k}, 0)$, $(0, p_i^{-k})$ for all $k > 0$, whenever $a_i = \infty = b_i$. $(p_i^{-b_i}, 0)$, $(f_{i,k} p_i^{-k-b_i}, p_i^{-k})$ where $f_{i,k}$ in \mathbf{Z} satisfies $f_{i,k} \equiv f_i \pmod{p_i^k}$ for all $k > 0$, whenever $a_i = \infty$, $b_i < \infty$.

Proof. Since A_f is torsion free and $\{y, x\}$ is a basis for $A_f \otimes Q$, $i(y, x)$ is an isomorphism of A_f onto its image $i(A_f)$. As checked in the

proof of Proposition 1.2, $A(f) \subset i(A_f)$. Moreover, one readily checks that $\text{char}(1, 0)$ in $A(f)$ equals (b_1, \dots, b_n, \dots) by determining the p -divisibility of $(1, 0)$ in $A(f) \otimes \mathbf{Z}_p$ for all primes p . Therefore, $A(f) \cap (Q \otimes 0) = i(Y)$. Furthermore, the inclusion $A(f) \subset i(A_f)$ determines $A(f)/A(f) \cap (Q \oplus 0) \subset i(X)$ with $(\overline{0, 1}) = i(y, x)(x)$. This is also an isomorphism, since $\text{char}(\overline{0, 1})$ in $A(f)/A(f) \cap (Q \oplus 0)$ equals $\text{char}(x)$ in X . Thus, $A(f)$ equals $i(A_f)$.

By employing the matrix classification for rank 2, torsion free abelian groups, we give below necessary and sufficient conditions for two extension functions f and g to determine isomorphic total groups A_f and A_g . Because of the very ineffective nature of the Korosh-Malcev-Derry matrix classification, the significance of Proposition 1.4 is the explicit necessary condition it provides.

PROPOSITION 1.4. *Let X and Y be rank 1, torsion free abelian groups, let x in X and y in Y be nonzero elements, and let $\text{char}(x) = (a_1, \dots, a_n, \dots)$ and $\text{char}(y) = (b_1, \dots, b_n, \dots)$. Two extension functions f, g in $\Pi' \mathbf{Z}/p_i^{a_i}$ determine isomorphic total groups A_f and A_g if and only if there exists a matrix $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ in $\text{GL}(Q \oplus Q)$ and matrices $\begin{pmatrix} \alpha_i & \beta_i \\ \gamma_i & \delta_i \end{pmatrix}$ in $\text{GL}(Y \otimes \mathbf{Z}_{p_i} \oplus X \otimes \mathbf{Z}_{p_i})$ satisfying*

$$(1.4.1)_i \quad \begin{pmatrix} p_i^{b_i} & -f_i \\ 0 & p_i^{a_i} \end{pmatrix} \cdot \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \alpha_i & \beta_i \\ \gamma_i & \delta_i \end{pmatrix} \cdot \begin{pmatrix} p_i^{b_i} & -g_i \\ 0 & p_i^{a_i} \end{pmatrix}$$

for all i . In particular, if A_f is isomorphic to A_g , then there exists $\alpha, \beta, \gamma, \delta$ in Q satisfying $\alpha\delta - \beta\gamma \neq 0$ such that f and g satisfy the following congruence condition for every i with $b_i < \infty$:

$$(1.4.2)_i \quad p_i^{b_i} \cdot \beta - f_i \cdot \delta \equiv g_i (f_i p_i^{-b_i} \gamma - \alpha) \pmod{p_i^{a_i}}$$

Proof. The conditions $(1.4.1)_i$ express the relationship between matrix sequences for bases $\{x, y\}$ and $\{x', y'\}$ of $A_f \otimes Q \cong Q \oplus Q \cong A_g \otimes Q$ related by $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ in $\text{GL}(Q \oplus Q)$ in terms of bases $\{x_i, y_i\}$ and $\{x'_i, y'_i\}$ of $Y \otimes \mathbf{Z}_{p_i} \oplus X \otimes \mathbf{Z}_{p_i}$ related by $\begin{pmatrix} \alpha_i & \beta_i \\ \gamma_i & \delta_i \end{pmatrix}$ in $\text{GL}(Y \otimes \mathbf{Z}_{p_i} \oplus X \otimes \mathbf{Z}_{p_i})$. The necessity of conditions $(1.4.1)_i$ is clear; their sufficiency is well known (see [2]). If $b_i < \infty$ and $\begin{pmatrix} \alpha_i & \beta_i \\ \gamma_i & \delta_i \end{pmatrix}$ is in $\text{GL}(Y \otimes \mathbf{Z}_{p_i} \oplus X \otimes \mathbf{Z}_{p_i})$ then β_i is in \mathbf{Z}_{p_i} . With this in mind, conditions $(1.4.2)_i$ quickly follow from $(1.4.1)_i$.

Since the 0 function, $g = 0$, determines $A_g = Y \oplus X$, conditions (1.4.2)_{*i*} determine useful necessary conditions an extension function f must satisfy for A_f to be isomorphic to $Y \oplus X$.

The following definition formulizes the relationship between extension functions f and g which determine isomorphic total groups A_f and A_g .

DEFINITION 1.5. Let characteristics (a_1, \dots, a_n, \dots) and (b_1, \dots, b_n, \dots) be given and let f, g be elements of $\Pi'Z/p_i^{a_i}$ where Π' is the product over all i with $b_i < \infty$. Then f and g are said to be of the same type with respect to (b_1, \dots, b_n, \dots) if there exists $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ in $GL(Q \oplus Q)$ and $\begin{pmatrix} \alpha_i & \beta_i \\ \gamma_i & \delta_i \end{pmatrix}$ in $GL(p_i^{-b_i}Z_{p_i} \oplus p_i^{-a_i}Z_{p_i})$ such that each of the equations (1.4.1)_{*i*} holds, where $p_i^{-m}Z_{p_i}$ is either Z_{p_i} or Q_{p_i} depending on whether $m < \infty$ or $m = \infty$.

2. Applications. As a first step toward classifying those rank 2, torsion free abelian groups A admitting a pure subgroup isomorphic to a given rank 1 group Y , we verify that the isomorphism class of A/H is independent of the choice of pure subgroup H isomorphic to Y . The proof of the following proposition was given to us by the referee.

PROPOSITION 2.1. *Let Y be a rank 1, torsion free abelian group, and let A be a rank 2, torsion free abelian group. For any two pure subgroups H, H' of A isomorphic to Y , A/H is isomorphic to A/H' .*

Proof. Let x and x' be nonzero elements of H and H' respectively. If there exists nonzero $m/n, m'/n'$ in Q with $m/n \cdot x + m'/n' \cdot x' = 0$ in A , then H and H' are equal; namely, each is the pure subgroup generated by $n'mx = -nm'x'$. Consequently, we may assume $\{x, x'\}$ span $A \otimes Q$.

Define $H_0 = \{r \text{ in } Q \mid rx + sx' \text{ is in } A \text{ for some } s \text{ in } Q\}$. Then the map $A \rightarrow H_0$ sending $rx + sx'$ to r is surjective with kernel H' . Similarly, if $H'_0 = \{s \text{ in } Q \mid rx + sx' \text{ is in } A \text{ for some } r \text{ in } Q\}$, then $A \rightarrow H'_0$ has kernel H . Therefore, it suffices to prove that H_0 is isomorphic to H'_0 .

Consider the inclusions $H \subset H_0$ sending rx to r , and $H' \subset H'_0$ sending sx' to s . Then $H_0 \rightarrow H'_0/H'$ sending r to the class of some s such that $rx + sx'$ is in A is well defined: if $rx + s'x'$ is also in A , then $(s - s')x'$ is in H' . Clearly, this map induces an isomorphism $H_0/H \rightarrow H'_0/H'$. Since H_0 and H'_0 are rank 1, torsion free with isomorphic subgroups H and H' respectively and isomorphic quotient groups H_0/H and H'_0/H' , H_0 is isomorphic to H'_0 .

The following theorem summarizes the discussion of §1 together with Proposition 2.1.

THEOREM 2.2. *Let Y be a rank 1, torsion free abelian group and let y in Y be a nonzero element. Then there is a natural 1-1 correspondence with explicit inverse*

$$\Phi(Y, y): E(Y) \rightarrow T(Y, y)$$

between the set $E(Y)$ of isomorphism classes of rank 2, torsion free abelian groups with a pure subgroup isomorphic to Y and the set $T(Y, y)$ of pairs consisting of

- (i) the type of some characteristic (a_1, \dots, a_n, \dots)
- (ii) the type with respect to $\text{char}(y)$ of some extension function f in $\Pi'Z/p_i^{a_i}$, which transforms to $\pm p_1^{c_1} \cdots p_n^{c_n} \cdots f$ in $\Pi'Z/p_i^{a_i+c_i}$ if all $c_i \geq 0$ and $\sum c_i < \infty$.

Proof. We define $\Phi(Y, y)(A)$ for A in $E(Y)$ as follows. Choose a rank 1, pure subgroup H of A with $O \neq h$ in H such that $\text{char}(h) = \text{char}(y) = (b_1, \dots, b_n, \dots)$. Let x in A/H be any nonzero element with $\text{char}(x) = (a_1, \dots, a_n, \dots)$. Let f in $\Pi'Z/p_i^{a_i}$ be an extension function representing $\varphi(A/H, x; H, h)(0 \rightarrow H \rightarrow A \rightarrow A/H \rightarrow 0)$ as in Proposition 1.1. Define $\Phi(Y, y)(A) = \{\text{char}(x), f\}$.

By Proposition 2.1, the type of $\text{char}(x)$ is well defined, independent of the choice of H or x . By Proposition 1.4, type of f depends at most upon choices for $A/H, x, H, h$, since A is isomorphic to A_p . If H' is another rank 1, pure subgroup of A with $O \neq h'$ in H' such that $\text{char}(h') = \text{char}(y)$, if x' in A/H' is chosen with $\text{char}(x') = \text{char}(x)$, and if f' in $\Pi'Z/p_i^{a_i}$ represents

$$\varphi(A/H', x'; H', h')(0 \rightarrow H' \rightarrow A \rightarrow A/H' \rightarrow 0),$$

then

$$\left\{ \begin{pmatrix} p_i^{b_i} & -f_i \\ 0 & p_i^{a_i} \end{pmatrix} \right\}, \quad \left\{ \begin{pmatrix} p_i^{b_i} & -f'_i \\ 0 & p_i^{a_i} \end{pmatrix} \right\}$$

are each matrix sequences for A by Proposition 1.2. Hence, by Proposition 1.4, f and f' have the same type. Finally, for a given rank 1, pure subgroup H of A with $O \neq h$ in H such that $\text{char}(h) = \text{char}(y)$, let $x' = mx$ be an integer nonzero multiple of x in A/H . As an extension function for $0 \rightarrow H \rightarrow A \rightarrow A/H \rightarrow 0$, f may be viewed as a classifying map $f: (A/H)/Z \cdot x \rightarrow Q/H$; this induces the classifying map $f': (A/H)/Z \cdot x' \rightarrow Q/H$. Under the identifications

$$\text{Hom}((A/H)/\mathbf{Z} \cdot x, Q/H) = \Pi' \mathbf{Z}/p_i^{a_i},$$

$\text{Hom}((A/H)/\mathbf{Z} \cdot x', Q/H) = \Pi' \mathbf{Z}/p_i^{a_i+c_i}$ with $c_i = \max\{c \mid p_i^c \text{ divides } m\}$, $f' = \pm p_i^{c_i} \cdots p_n^{c_n} \cdots f$ in $\Pi' \mathbf{Z}/p_i^{a_i}$.

The naturality of $\Phi(Y, y)$ with respect to Y and y follows from the naturality of extension functions as given in Proposition 1.1. The fact that $\Phi(Y, y)$ is injective follows from the fact that A is isomorphic to A_f if $\Phi(Y, y)(A) = \{(a_1, \dots, a_n, \dots), f\}$. The fact that $\Phi(Y, y)$ is surjective follows from Proposition 1.2, since $\Phi(Y, y)(A_f) = \{(a_1, \dots, a_n, \dots), f\}$. Finally, the explicit inverse of $\Phi(Y, y)$ is given in Proposition 1.3.

We say that a torsion free abelian group A is homogeneous of type (b_1, \dots, b_n, \dots) if for every nonzero x in A , the type of $\text{char}(x)$ equals the type of (b_1, \dots, b_n, \dots) . Using Theorem 2.2, we “list” all rank 2, torsion free abelian groups which are homogeneous of type

$$(b_1, \dots, b_n, \dots).$$

THEOREM 2.3. *Let (b_1, \dots, b_n, \dots) be a characteristic and let Y be a rank 1, torsion free abelian group with $O \neq y$ in Y such that $\text{char}(y) = (b_1, \dots, b_n, \dots)$. Then the set of all rank 2, torsion free abelian groups homogeneous of type (b_1, \dots, b_n, \dots) is the subset of $E(Y)$ consisting of those A in $E(Y)$ such that $\Phi(Y, y)(A) = \{(a_1, \dots, a_n, \dots), f \text{ in } \Pi' \mathbf{Z}/p_i^{a_i}\}$ satisfies*

- (i) if $b_i = \infty$, then $a_i = \infty$;
- (ii) if $b_i < \infty$ and $a_i = \infty$, then f_i in \mathbf{Z}_{p_i} is not rational;
- (iii) for all but finitely many i with $b_i < \infty$, $a_i \geq b_i$ and $p_i^{b_i} \mid f_i$;
- (iv) for every nonzero pair of integers $\{m, n\}$, there are only finitely many i with $a_i > b_i$, $p_i^{b_i} \mid f_i$, and $m - nf_i/p_i^{b_i}$ divisible by p_i .

Proof. The set of isomorphism classes of rank 2, torsion free abelian groups homogeneous of type (b_1, \dots, b_n, \dots) is clearly a subset of $E(Y)$. To prove the theorem, it suffices to prove the following for a given pair $\{(a_1, \dots, a_n, \dots), f \text{ in } \Pi' \mathbf{Z}/p_i^{a_i}\}$: the characteristic of every nonzero element (m, n) in $A(f) \cap \mathbf{Z} \oplus \mathbf{Z}$ as an element of $A(f)$ has the same type as (b_1, \dots, b_n, \dots) iff conditions (i)–(iv) are satisfied (where $A(f)$ is given in Proposition 1.3.). We write

$$(m, n) = (mp_i^{b_i} - nf_i)(p_i^{-b_i}, 0) + np_i^{a_i}(f_i p_i^{-a_i-b_i}, p_i^{-a_i})$$

in $A(f) \otimes \mathbf{Z}_{p_i}$, where $p_i^\infty = 1$ and $f_i = 0$ if $b_i = \infty$. We observe for any $k > 0$ that (m, n) is divisible by p_i^k in $A(f)$ iff (m, n) is divisible by p_i^k in $A(f) \otimes \mathbf{Z}_{p_i}$.

If $b_i = \infty$, then every (m, n) in $A(f) \cap \mathbf{Z} \oplus \mathbf{Z}$ is infinitely p_i divisible (as an element of $A(f)$) iff $a_i = \infty$. If $b_i < \infty$, then (m, n) in

$A(f) \cap \mathbf{Z} \oplus \mathbf{Z}$ is not infinitely p_i divisible iff either $a_i < \infty$ or $a_i = \infty$ and $mp_i^{b_i} - nf_i \neq 0$. Consequently, (i) and (ii) are equivalent to the condition that one nonzero element of $A(f)$ is infinitely p_i divisible iff all nonzero elements of $A(f)$ are.

(m, n) in $A(f) \cap \mathbf{Z} \oplus \mathbf{Z}$ is divisible by $p_i^{b_i}$ for all but finitely many i with $b_i < \infty$ iff $n = 0$ or for all but finitely many i with $b_i < \infty$, $a_i \geq b_i$ and $p_i^{b_i} | f_i$. (m, n) is not divisible by $p_i^{b_i+1}$ for all but finitely many i with $b_i < \infty$ iff for all but finitely many i with $a_i > b_i$, $mp_i^{b_i} - nf_i$ is not divisible by $p_i^{b_i+1}$. Consequently, (iii) and (iv) are equivalent to the condition that the characteristic of any nonzero element of $A(f)$ differs from (b_1, \dots, b_n, \dots) at only finitely many i with b_i .

To show how explicit Theorem 2.3 is, we provide the following simple example.

EXAMPLE 2.4. Let (b_1, \dots, b_n, \dots) be a characteristic such that $b_i < \infty$ for infinitely many i . Let Y be a rank 1, torsion free abelian group with nonzero element y with $\text{char}(y) = (b_1, \dots, b_n, \dots)$. Then for any positive integer k ,

$$\Phi(Y, y)^{-1}(\{b_1 + k, \dots, b_n + k, \dots\}, f = i \cdot p_i^{b_i} \text{ in } \Pi' \mathbf{Z}/p_i^{b_i+k})$$

is an indecomposable, rank 2, torsion free abelian group homogeneous of type (b_1, \dots, b_n, \dots) .

Proof. If A in $E(Y)$ is homogeneous and decomposable, then $A \approx Y \oplus Y$ so that the quotient type of A (i.e., the first invariant of Theorem 2.2) must be the type of (b_1, \dots, b_n, \dots) . Consequently, to check the example it suffices to verify conditions (i)–(iv) of Theorem 2.3 for $\{(b_1 + k, \dots, b_n + k, \dots), f = i \cdot p_i^{b_i}\}$. Conditions (i), (ii), and (iii) are immediate. To verify condition (iv), we observe that for sufficiently large i $0 < n \cdot i - m < p_i$ by the prime number theorem.

In particular, Example 2.4 gives examples of nonfree homogeneous groups A , with type $A = (0, \dots, 0, \dots)$, such that $A \otimes \mathbf{Z}_p$ is a free \mathbf{Z}_p module for all primes p . These groups are called locally free.

As another application of Theorem 2.2, we determine the subset $E(\mathbf{Z}) \cap E(Y)$ of $E(Y)$.

PROPOSITION 2.5. Let (b_1, \dots, b_n, \dots) be a characteristic and let Y be a rank 1, torsion free abelian group with $0 \neq y$ in Y such that $\text{char}(y) = (b_1, \dots, b_n, \dots)$. Then the subset $E(\mathbf{Z}) \cap E(Y)$ of $E(Y)$ consists of those A in $E(Y)$ such that $\Phi(Y, y)(A) = \{(a_1, \dots, a_n, \dots), f \text{ in } \Pi' \mathbf{Z}/p_i^{a_i}\}$ satisfies:

- (i) if $b_i = \infty$, then $a_i < \infty$;
- (ii) for all but finitely many i with $a_i, b_i > 0$, p_i does not divide f_i ;

(iii) *there exists a nonzero pair of integers $\{m, n\}$ such that $mp_i^{b_i} - nf_i \neq 0$ whenever $a_i = \infty$, and $m - nf_i$ is divisible by p_i for only finitely many i with $b_i = 0$ and $a_i > 0$.*

Proof. For Y cyclic, conditions (i), (ii), and (iii) are immediately satisfied for any pair of invariants $\{(a_1, \dots, a_n, \dots), f\}$. We may thus assume Y is not cyclic. As in the proof of Theorem 2.3, to prove the proposition it suffices to prove that (m, n) in $A(f) \cap \mathbf{Z} \oplus \mathbf{Z}$ generates a pure cyclic subgroup of $A(f)$ iff (i) and (ii) are satisfied and $\{m, n\}$ satisfies (iii). Since Y is assumed not to be cyclic, we need only consider (m, n) with $n \neq 0$. We write

$$(m, n) = (mp_i^{b_i} - nf_i)(p_i^{-b_i}, 0) + np_i^{a_i}(f_i p_i^{-a_i - b_i}, p_i^{-a_i}).$$

Now, (m, n) with $n \neq 0$ is not infinitely divisible in $A(f) \otimes \mathbf{Z}_{p_i}$ by p_i iff whenever $b_i = \infty$, then $a_i < \infty$; and whenever $a_i = \infty$, then $mp_i^{b_i} - nf_i \neq 0$. Furthermore, (m, n) with $n \neq 0$ is divisible by only finitely many p_i iff for only finitely many i with $b_i, a_i > 0$ f_i is divisible by p_i ; and for only finitely many i with $b_i = 0, a_i > 0$ $m - nf_i$ is divisible by p_i .

REFERENCES

1. R. A. Beaumont and R. S. Pierce, *Torsion free groups of rank two*, Mem. Amer. Math. Soc. Nr. 38.
2. L. Fuchs, *Infinite abelian groups*, Vol. II, Academic Press, New York and London, 1973.

Received January 23, 1974. Partially supported by the U.S. — France Exchange of Scientists Program.

PRINCETON UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$ 72.00 a year (6 Vols., 12 issues). Special rate: \$ 36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1975 Pacific Journal of Mathematics
All Rights Reserved

Zvi Artstein and John Allen Burns, <i>Integration of compact set-valued functions</i>	297
Mark Benard, <i>Characters and Schur indices of the unitary reflection group [321]³</i>	309
Simeon M. Berman, <i>A new characterization of characteristic functions of absolutely continuous distributions</i>	323
Monte Boisen and Philip B. Sheldon, <i>Pre-Prüfer rings</i>	331
Hans-Heinrich Brungs, <i>Three questions on duo rings</i>	345
Iracema M. Bund, <i>Birnbau-Orlicz spaces of functions on groups</i>	351
John D. Elwin and Donald R. Short, <i>Branched immersions between 2-manifolds of higher topological type</i>	361
Eric Friedlander, <i>Extension functions for rank 2, torsion free abelian groups</i>	371
Jon Froemke and Robert Willis Quackenbush, <i>The spectrum of an equational class of groupoids</i>	381
Barry J. Gardner, <i>Radicals of supplementary semilattice sums of associative rings</i>	387
Shmuel Glasner, <i>Relatively invariant measures</i>	393
George Rudolph Gordh, Jr. and Sibe Mardesic, <i>Characterizing local connectedness in inverse limits</i>	411
Siegfried Graf, <i>On the existence of strong liftings in second countable topological spaces</i>	419
Stanley P. Gudder and D. Strawther, <i>Orthogonally additive and orthogonally increasing functions on vector spaces</i>	427
Darald Joe Hartfiel and Carlton James Maxson, <i>A characterization of the maximal monoids and maximal groups in β_X</i>	437
Robert E. Hartwig and S. Brent Morris, <i>The universal flip matrix and the generalized faro-shuffle</i>	445
William Emery Haver, <i>Mappings between ANRs that are fine homotopy equivalences</i>	457
J. Bockett Hunter, <i>Moment sequences in l^p</i>	463
Barbara Jeffcott and William Thomas Spears, <i>Semimodularity in the completion of a poset</i>	467
Jerry Alan Johnson, <i>A note on Banach spaces of Lipschitz functions</i>	475
David W. Jonah and Bertram Manuel Schreiber, <i>Transitive affine transformations on groups</i>	483
Karsten Juul, <i>Some three-point subset properties connected with Menger's characterization of boundaries of plane convex sets</i>	511
Ronald Brian Kirk, <i>The Haar integral via non-standard analysis</i>	517
Justin Thomas Lloyd and William Smiley, <i>On the group of permutations with countable support</i>	529
Erwin Lutwak, <i>Dual mixed volumes</i>	531
Mark Mahowald, <i>The index of a tangent 2-field</i>	539
Keith Miller, <i>Logarithmic convexity results for holomorphic semigroups</i>	549
Paul Milnes, <i>Extension of continuous functions on topological semigroups</i>	553
Kenneth Clayton Pietz, <i>Cauchy transforms and characteristic functions</i>	563
James Ted Rogers Jr., <i>Whitney continua in the hyperspace $C(X)$</i>	569
Jean-Marie G. Rolin, <i>The inverse of a continuous additive functional</i>	585
William Henry Ruckle, <i>Absolutely divergent series and isomorphism of subspaces</i>	605
Rolf Schneider, <i>A measure of convexity for compact sets</i>	617
Alan Henry Schoenfeld, <i>Continuous measure-preserving maps onto Peano spaces</i>	627
V. Merriline Smith, <i>Strongly superficial elements</i>	643
Roger P. Ware, <i>A note on quadratic forms over Pythagorean fields</i>	651
Roger Allen Wiegand and Sylvia Wiegand, <i>Finitely generated modules over Bezout rings</i>	655
Martin Ziegler, <i>A counterexample in the theory of definable automorphisms</i>	665