

Pacific Journal of Mathematics

**ON THE EXISTENCE OF STRONG LIFTINGS IN SECOND
COUNTABLE TOPOLOGICAL SPACES**

SIEGFRIED GRAF

ON THE EXISTENCE OF STRONG LIFTINGS IN SECOND COUNTABLE TOPOLOGICAL SPACES

SIEGFRIED GRAF

Let X be a second countable topological space, \mathfrak{A} a σ -field of subsets of X containing all open sets and μ a finite positive measure on \mathfrak{A} , such that (X, \mathfrak{A}, μ) is a complete measure space and $\mu(U) > 0$ for every nonempty open $U \subset X$.

Then there exists a lifting $\phi : \mathfrak{A} \rightarrow \mathfrak{A}$ which satisfies $U \subset \phi(U)$ for every open subset $U \subset X$.

Basic notations and definitions. Throughout this paper \mathbf{N} denotes the nonnegative integers and \mathbf{R}_+ the nonnegative real numbers. Moreover

X is a second countable topological space ($\neq \emptyset$),
 \mathfrak{A} is a σ -field of subsets of X containing all
 open sets,
 $\mu : \mathfrak{A} \rightarrow \mathbf{R}_+$ is a countable additive measure,
 satisfying $\mu(U) > 0$ for every nonempty open
 subset $U \subset X$.

For $A, B \in \mathfrak{A}$ we denote by $A \subseteq B$ the fact that $\mu(A \setminus B) = 0$ and write $A \sim B$ if $A \subseteq B$ and $B \subseteq A$.

A subset $\mathcal{F} \subset \mathfrak{A}$ is called an \cap -system iff $\emptyset \in \mathcal{F}$, $X \in \mathcal{F}$ and $A \cap B \in \mathcal{F}$ for all $A, B \in \mathcal{F}$.

For an \cap -system $\mathcal{F} \subset \mathfrak{A}$ a mapping $\delta : \mathcal{F} \rightarrow \mathfrak{A}$ is a *partial (lower) density* iff it satisfies the following conditions:

- (i) $A \sim B$ implies $\delta(A) = \delta(B)$
- (ii) $A \sim \delta(A)$
- (iii) $\delta(\emptyset) = \emptyset$ and $\delta(X) = X$
- (iv) $\delta(A \cap B) = \delta(A) \cap \delta(B)$ for every $A, B \in \mathcal{F}$.

A mapping $\delta : \mathfrak{A} \rightarrow \mathfrak{A}$ with the above properties is called a (lower) *density* and if moreover

- (v) $\delta(X \setminus A) = X \setminus \delta(A)$ for every $A \in \mathfrak{A}$
- is fulfilled δ is a *lifting*.

A lifting or density $\delta : \mathfrak{A} \rightarrow \mathfrak{A}$ is called *strong* iff in addition

- (vi) $U \subset \delta(U)$ for every open $U \subset X$.

Let us restate one of the fundamental properties of partial densities. For an \cap -system $\mathcal{F} \subset \mathfrak{A}$, a partial density $\delta : \mathcal{F} \rightarrow \mathfrak{A}$, and $A, B \in \mathcal{F}$ with $A \subseteq B$ we have $\delta(A) \subseteq \delta(B)$. In particular δ preserves inclusions.

Existence of Strong Densities. The following two lemmas will provide us with the basic tools for the proof of the main theorem.

LEMMA 1. Let $\mathcal{F} \subset \mathcal{A}$ be an \cap -system, $\delta: \mathcal{F} \rightarrow \mathcal{A}$ a partial density, and $F_0 \in \mathcal{A}$ arbitrary. Suppose $A_0 \in \mathcal{A}$ is such that $A_0 \sim F_0$ and for F'', F', F in \mathcal{F} ,

$$F'' \subseteq F_0 \text{ implies } \delta(F'') \subset A_0,$$

and

$$F_0 \cap F \subseteq F' \text{ implies } A_0 \cap \delta(F) \subset \delta(F').$$

Then $\mathcal{F}_0 = \mathcal{F} \cup \{F \cap F_0: F \in \mathcal{F}\}$ is an \cap -system containing $\mathcal{F} \cup \{F_0\}$ and the mapping $\delta_0: \mathcal{F}_0 \rightarrow \mathcal{A}$ defined by

$$\delta_0(F) = \begin{cases} \delta(F), & \text{if } F \in \mathcal{F} \\ \delta(F') \cap A_0 & \text{if } F = F' \cap F_0 \text{ with } F' \in \mathcal{F} \end{cases}$$

is a partial density which extends δ .

Proof. It is obvious that \mathcal{F}_0 is an \cap -system containing $\mathcal{F} \cup \{F_0\}$. To show that δ_0 is well-defined let F, F' be elements of \mathcal{F} .

(1) Assume $F \sim F' \cap F_0$. This implies

$$F \subseteq F_0, \quad F \subseteq F' \text{ and } F' \cap F_0 \subseteq F.$$

From our assumptions concerning A_0 we may therefore conclude

$$(*) \quad \delta(F) \subset A_0 \text{ and } A_0 \cap \delta(F') \subset \delta(F).$$

Since δ is a partial density $F \subseteq F'$ implies $\delta(F) \subset \delta(F')$. This inclusion combined with (*) gives us $\delta(F) \subset A_0 \cap \delta(F') \subset \delta(F)$. Hence the equality $\delta(F) = A_0 \cap \delta(F')$ is established.

(2) Assume $F \cap F_0 \sim F' \cap F_0$. Then we have

$$F \cap F_0 \subseteq F' \text{ and } F' \cap F_0 \subseteq F.$$

Again it follows from our assumptions concerning A_0 that

$$A_0 \cap \delta(F) \subset \delta(F') \text{ and } A_0 \cap \delta(F') \subset \delta(F)$$

and hence that

$$A_0 \cap \delta(F') = A_0 \cap \delta(F).$$

(1) and (2) together prove that δ_0 is well-defined and that for $F, G \in \mathcal{F}_0$ the fact $F \sim G$ implies $\delta_0(F) = \delta_0(G)$. According to its definition δ_0 obviously satisfies the other conditions for a partial density. It is also clear that δ_0 extends δ .

LEMMA 2. *Let $\mathcal{F} \subset \mathfrak{A}$ be a countable \cap -system, $F_0 \in \mathfrak{A}$ arbitrary, $\mathcal{F}_0 = \mathcal{F} \cup \{F \cap F_0 : F \in \mathcal{F}\}$ and $\delta : \mathcal{F} \rightarrow \mathfrak{A}$ a partial density. Then there exists a partial density $\delta_0 : \mathcal{F}_0 \rightarrow \mathfrak{A}$ which extends δ .*

Proof. According to Lemma 1 the assertion of Lemma 2 is true if we can prove the existence of a set $A_0 \in \mathfrak{A}$ such that $F_0 \sim A_0$ and for F, F', F'' in \mathcal{F} ,

$$F'' \subseteq F_0 \text{ implies } \delta(F'') \subset A_0,$$

and

$$F_0 \cap F \subset F' \text{ implies } A_0 \cap \delta(F) \subset \delta(F').$$

As an easy calculation shows $A_0 \in \mathfrak{A}$ fulfills the last two conditions if and only if

$$\delta(F'') \subset A_0 \subset \delta(F') \cup C\delta(F)$$

whenever $F, F', F'' \in \mathcal{F}$ satisfy $F'' \subseteq F_0 \subseteq F' \cup CF$.

To establish the existence of such an A_0 let

$$\mathcal{G} := \{\delta(F'') : F'' \subset \mathcal{F} \text{ and } F'' \subseteq F_0\}$$

and

$$\mathcal{H} := \{\delta(F') \cup C\delta(F) : F, F' \in \mathcal{F} \text{ and } F_0 \subseteq F' \cup CF\}$$

Since \mathcal{F} is countable the sets \mathcal{G} and \mathcal{H} are also countable. Therefore $A := \cup \mathcal{G}$ and $B := \cap \mathcal{H}$ are in \mathfrak{A} . Property (ii) of partial densities implies

$$G \subseteq F_0 \text{ and } F_0 \subseteq H$$

for every $G \in \mathcal{G}$ and $H \in \mathcal{H}$. Due to the countable additivity of μ we therefore get:

$$(\dagger) \quad A \subseteq F_0 \text{ and } F_0 \subseteq B.$$

Let $G \in \mathcal{G}$ and $H \in \mathcal{H}$ be arbitrary. Then there are $F, F', F'' \in \mathcal{F}$ with $F'' \subseteq F_0 \subseteq F' \cup CF$, $G = \delta(F'')$, and $H = \delta(F') \cup C\delta(F)$. From $F'' \subseteq F' \cup CF$ we conclude

$$F'' \cap F \subseteq F' \text{ and hence } \delta(F'') \cap \delta(F) \subset \delta(F')$$

which implies $G = \delta(F'') \subset \delta(F') \cup C\delta(F) = H$. This gives us

$$(\ddagger) \quad A \subset B.$$

It is a consequence of (\dagger) and (\ddagger) that

$$A_0 := (F_0 \cup A) \cap B \sim F_0$$

and

$$G \subset A_0 \subset H$$

for all $G \in \mathcal{G}$, $H \in \mathcal{H}$.

Hence the statement of Lemma 2 is proved.

PROPOSITION 1. *Let \mathcal{B} be a countable base of the topology of X and \mathcal{F} the field generated by \mathcal{B} . Then there is a partial density $\delta: \mathcal{F} \rightarrow \mathfrak{A}$ satisfying $U \subset \delta(U)$ for every $U \in \mathcal{B}$.*

Proof. Without loss of generality we may assume that \mathcal{B} is stable under finite unions and intersections and contains \emptyset and X .

For $B \in \mathcal{B}$ let $\mathcal{B}(B) := \{A \in \mathcal{B} : A \sim B\}$ and define $\phi: \mathcal{B} \rightarrow \mathfrak{A}$ by $\phi(B) = \cup \mathcal{B}(B)$. Since $\cup \mathcal{B}(B)$ is open for every $B \in \mathcal{B}$ and since \mathfrak{A} contains all open sets ϕ is a well-defined map from \mathcal{B} to \mathfrak{A} . It is an immediate consequence of the definition of ϕ that for $A, B \in \mathcal{B}$ the equivalence $A \sim B$ implies $\phi(A) = \phi(B)$. Since $\mathcal{B}(B)$ is countable we have $B \sim \phi(B)$.

According to our assumptions every nonempty open set has strictly positive measure. Therefore $\mathcal{B}(\emptyset) = \{\emptyset\}$ and hence $\phi(\emptyset) = \emptyset$. Since $X \in \mathcal{B}(X)$ the condition $\phi(X) = X$ is fulfilled.

ϕ also preserves finite intersections. To show this let $A, B \in \mathcal{B}$ be arbitrary. Because \mathcal{B} is stable under finite intersections we have

$$\{A' \cap B' : A' \in \mathcal{B}(A), B' \in \mathcal{B}(B)\} \subset \mathcal{B}(A \cap B)$$

and consequently

$$\begin{aligned} \phi(A) \cap \phi(B) &= (\cup \mathcal{B}(A)) \cap (\cup \mathcal{B}(B)) \\ &= \cup \{A' \cap B' : A' \in \mathcal{B}(A), B' \in \mathcal{B}(B)\} \\ &\subset \cup \mathcal{B}(A \cap B) = \phi(A \cap B). \end{aligned}$$

To prove the inverse inclusion let $C \in \mathcal{B}(A \cap B)$ be arbitrary. Since \mathcal{B} is stable under finite unions we have $A \cup C \in \mathcal{B}(A)$ and hence $C \subset \phi(A)$. For the same reasons $C \subset \phi(B)$ is true and therefore the inclusion

$$\phi(A \cap B) \subset \phi(A) \cap \phi(B)$$

holds.

Using Lemma 2, the fact that \mathcal{F} is countable, and the fact that \mathcal{B} is an \cap -system contained in \mathcal{F} , we see by induction that ϕ can be extended to a partial density $\delta: \mathcal{F} \rightarrow \mathcal{A}$. From the definition of ϕ it follows immediately that $U \subset \phi(U) = \delta(U)$ for all $U \in \mathcal{B}$.

The following two propositions are stated without proofs. The proofs can be found in [4], although a slightly different notation is used there.

PROPOSITION 2. ([4], p. 64, Lemma 4.6). *Let $(\mathcal{A}_n)_{n \in \mathbb{N}}$ be a sequence of σ -fields of subsets of X with*

$$\mathcal{A}_n \subset \mathcal{A}_{n+1} \subset \mathcal{A}$$

and $\delta_n: \mathcal{A}_n \rightarrow \mathcal{A}$ a partial density with

$$\delta_{n+1}|_{\mathcal{A}_n} = \delta_n$$

for every $n \in \mathbb{N}$.

Denote by \mathcal{A}_∞ the σ -field generated by $\cup \{\mathcal{A}_n : n \in \mathbb{N}\}$. Then there exists a partial density $\delta_\infty: \mathcal{A}_\infty \rightarrow \mathcal{A}$ satisfying $\delta_\infty|_{\mathcal{A}_n} = \delta_n$ for all $n \in \mathbb{N}$.

Though the above proposition is not explicitly stated in his paper Traynor proved it independently in [8].

PROPOSITION 3. ([4], p. 61, Satz. 4.3 and Lemma 4.4). *Let \mathcal{L} be a σ -field, $A_0 \in \mathcal{A}$ arbitrary, and $\delta: \mathcal{L} \rightarrow \mathcal{A}$ a partial density.*

Then $\mathcal{L}_0 := \{(B \cap A_0) \cup (B' \setminus A_0) : B, B' \in \mathcal{L}\}$ is a σ -field containing $\mathcal{L} \cup \{A_0\}$ and there exists a partial density $\delta_0: \mathcal{L}_0 \rightarrow \mathcal{A}$ extending δ . δ_0 can be defined by

$$\begin{aligned} & \delta_0((B \cap A_0) \cup (B' \setminus A_0)) \\ &= (\delta((B \cap B_1) \cup (B' \setminus B_1)) \cap A_0) \cup (\delta((B \setminus B_2) \cup (B' \cap B_2)) \setminus A_0), \end{aligned}$$

where B, B' are elements of \mathcal{L} and where $B_1 \in \mathcal{L}$ satisfies $A_0 \subseteq B_1$ and $B_1 \subseteq A$ for every $A \in \mathcal{L}$ with $A_0 \subseteq A$ while $B_2 \in \mathcal{L}$ satisfies $CA_0 \subseteq B_2$ and $B_2 \subseteq C$ for every $C \in \mathcal{L}$ with $CA_0 \subseteq C$.

THEOREM 1. *There exists a strong lower density $\rho: \mathcal{A} \rightarrow \mathcal{A}$.*

Proof. Let \mathcal{B} be a countable base for the topology of X and \mathcal{F} the field generated by \mathcal{B} . According to Proposition 1 there is a partial density $\delta: \mathcal{F} \rightarrow \mathcal{A}$ such that $U \subset \delta(U)$ for every $U \in \mathcal{B}$.

Let $f: \mathbb{N} \rightarrow \mathcal{F}$ be a bijection and \mathfrak{A}_n the field generated by $\{f(m) : 0 \leq m \leq n\}$. It is clear that $\mathfrak{A}_n \subset \mathcal{F}$ is a σ -field. Define $\delta_n: \mathfrak{A}_n \rightarrow \mathfrak{A}$ by $\delta_n = \delta \upharpoonright \mathfrak{A}_n$. Then $(\mathfrak{A}_n, \delta_n)_{n \in \mathbb{N}}$ fulfills the assumptions of Proposition 2. Thus there is a partial density $\delta_\infty: \mathfrak{A}_\infty \rightarrow \mathfrak{A}$ extending δ , where \mathfrak{A}_∞ is the σ -field generated by $\mathcal{F} = \cup \{\mathfrak{A}_n : n \in \mathbb{N}\}$. Using Zorn's lemma and Propositions 2 and 3 we will show that δ_∞ can be extended to a density $\rho: \mathfrak{A} \rightarrow \mathfrak{A}$.

To this purpose define

$$\mathcal{T} := \{(\mathcal{L}, \psi) : \mathcal{L} \text{ } \sigma\text{-field containing } \mathfrak{A}_\infty \text{ and } \psi : \mathcal{L} \rightarrow \mathfrak{A} \\ \text{a partial density with } \psi \upharpoonright \mathfrak{A}_\infty = \delta_\infty\}.$$

For $(\mathcal{L}, \psi), (\mathcal{L}', \psi') \in \mathcal{T}$ let $(\mathcal{L}, \psi) \leq (\mathcal{L}', \psi')$ denote the fact that $\mathcal{L} \subset \mathcal{L}'$ and $\psi' \upharpoonright \mathcal{L} = \psi$. It is easy to check that \leq is an order-relation on \mathcal{T} .

Next we will prove that \mathcal{T} is inductively orderby by \leq .

Let $\mathcal{K} \subset \mathcal{L}$ be any totally ordered subset. The following two cases have to be considered.

(1) If for every sequence $(\mathcal{L}_n, \psi_n)_{n \in \mathbb{N}}$ in \mathcal{K} there is a $(\mathcal{L}, \psi) \in \mathcal{K}$ with $(\mathcal{L}_n, \psi_n) \leq (\mathcal{L}, \psi)$ for all $n \in \mathbb{N}$, then $\bar{\mathcal{L}} := \cup \{\mathcal{L} : (\mathcal{L}, \psi) \in \mathcal{K}\}$ is a σ -field containing \mathfrak{A}_∞ and $\bar{\psi}: \bar{\mathcal{L}} \rightarrow \mathfrak{A}$ defined by $\bar{\psi} \upharpoonright \mathcal{L} = \psi$ for every $(\mathcal{L}, \psi) \in \mathcal{K}$ is a partial density extending δ_∞ . Hence we have $(\bar{\mathcal{L}}, \bar{\psi}) \in \mathcal{T}$ and $(\mathcal{L}, \psi) \leq (\bar{\mathcal{L}}, \bar{\psi})$ for all $(\mathcal{L}, \psi) \in \mathcal{K}$.

(2) If there exists a sequence $(\mathcal{L}_n, \psi_n)_{n \in \mathbb{N}}$ in \mathcal{K} such that for each $(\mathcal{L}, \psi) \in \mathcal{K}$ there is an $n \in \mathbb{N}$ with $(\mathcal{L}, \psi) \leq (\mathcal{L}_n, \psi_n)$ we have

$$\cup \{\mathcal{L} : (\mathcal{L}, \psi) \in \mathcal{K}\} = \cup \{\mathcal{L}_n : n \in \mathbb{N}\} = \bar{\mathcal{L}}.$$

Let \mathcal{L}_∞ denote the σ -field generated by \mathcal{L} . From Proposition 2 we conclude the existence of a partial density $\psi_\infty: \mathcal{L}_\infty \rightarrow \mathfrak{A}$ which extends all ψ_n and hence we get $(\mathcal{L}, \psi) \leq (\mathcal{L}_\infty, \psi_\infty) \in \mathcal{T}$ for all $(\mathcal{L}, \psi) \in \mathcal{K}$.

Since $(\mathfrak{A}_\infty, \delta_\infty) \in \mathcal{T}$ and thus $\mathcal{T} \neq \emptyset$, Zorn's lemma gives us the existence of a maximal element $(\bar{\mathcal{F}}, \rho)$ in \mathcal{T} .

From Proposition 3 we conclude $\bar{\mathcal{F}} = \mathfrak{A}$ and hence ρ is a density. To prove the theorem it remains to check that ρ is strong. To this end let U be any open subset of X . Then

$$U = \cup \{B \in \mathcal{B} : B \subset U\}$$

for $B \in \mathcal{B}$ and $B \subset U$ we have

$$\rho(U) \supset \rho(B) = \delta_\infty(B) = \delta(B) = \delta(B) \supset B$$

and thus

$$U = \cup \{B \in \mathcal{B} : B \subset U\} \subset \rho(U).$$

Hence ρ is strong.

The main theorem. To verify the main theorem we need one more lemma which is an immediate consequence of a more general theorem of von Neumann and Stone ([6], p. 372, Th. 18). Easier direct proofs were given by Sion in [7], Gapaillard in [3], Traynor in [8] and the author in [4]. Let us restate this lemma but omit the proof.

LEMMA 3. *If (X, \mathfrak{A}, μ) is complete and $\delta : \mathfrak{A} \rightarrow \mathfrak{A}$ a density then there is a lifting $\phi : \mathfrak{A} \rightarrow \mathfrak{A}$ with $\delta(A) \subset \phi(A)$ for every $A \in \mathfrak{A}$.*

THEOREM 2. *If (X, \mathfrak{A}, μ) is complete then there exists a strong lifting $\phi : \mathfrak{A} \rightarrow \mathfrak{A}$.*

Proof. According to Theorem 1 there is a strong density $\delta : \mathfrak{A} \rightarrow \mathfrak{A}$. From Lemma 3 we get a lifting $\phi : \mathfrak{A} \rightarrow \mathfrak{A}$ satisfying $\delta(A) \subset \phi(A)$ for all $A \in \mathfrak{A}$. Since δ is strong we therefore conclude $U \subset \delta(U) \subset \phi(U)$ for every open set $U \subset X$.

Hence ϕ is a strong lifting.

Notes.

(1) All previous statements of this paper remain true for σ -finite measure spaces because for any σ -finite measure on a σ -field there exists a finite measure with the same nullsets.

(2) The methods of proofs show that for an arbitrary finite or σ -finite measure space (Y, \mathcal{L}, ν) and any countable $\mathcal{F} \subset \mathcal{L}$, such that for each finite $\mathcal{G} \subset \mathcal{F}$ with $\nu(\cap \mathcal{G}) = 0$ we have $\cap \mathcal{G} = \emptyset$, there is a density $\delta : \mathcal{L} \rightarrow \mathcal{L}$ satisfying $F \subset \delta(F)$ for all $F \in \mathcal{F}$.

If (Y, \mathcal{L}, ν) is furthermore complete then there is even a lifting $\phi : \mathcal{L} \rightarrow \mathcal{L}$ with these properties.

(3) Let X be a second countable topological space, \mathfrak{A} the Borel-field in X and \mathfrak{B} a σ -ideal in \mathfrak{A} containing no nonempty open sets. Then \mathfrak{A} has at most the power of the continuum. Assuming the continuum hypothesis and using the same methods as in the first part of this paper we get a strong density $\delta : \mathfrak{A} \rightarrow \mathfrak{A}$ with respect to \mathfrak{B} . From a lifting theorem of von Neumann-Stone (certificate [6] or [4], p. 47, Satz 3.3) we may conclude (using the continuum hypothesis another time) that there is even a strong lifting $\phi : \mathfrak{A} \rightarrow \mathfrak{A}$.

(4) The existence of a strong lifting was well-known for separable metric spaces with finite outer regular measures strictly positive on non-empty open sets (see Ionescu-Tulcea [5] and Sion [7]) and for locally compact metrizable spaces with (not necessarily σ -finite) Radon measures whose support is the whole space (see Ionescu-Tulcea [5]). The first fact is a special case of the main theorem of this paper while the second fact can be derived from this theorem using Proposition 2, p. 108 in Ionescu-Tulcea [5], and observing that any metrizable compact space is separable.

(5) Bichteler proves in [1] a strong-lifting-theorem implying the existence of a strong lifting for locally compact metrizable spaces with

Radon measures whose support is the whole space. But the result of this paper are no (immediate) consequences of Bichteler's theorem.

(6) Eifrig [2] gives a proof of the existence of a strong lifting for the interval $[0, 1]$ using methods of nonstandard-analysis.

(7) For applications of strong-lifting-theorems to integral representations and disintegration of measures see Ionescu-Tulcea [5] and Sion [7].

REFERENCES

1. K. Bichteler, *Integration Theory* (with Special Attention to Vector Measures), Lect. Notes in Mathematics **315** (1973), Springer Verlag, Berlin-Heidelberg-New York.
2. B. Eifrig, *Ein nicht-standard Beweis für die Existenz eines starken Liftings in \mathcal{L}_∞ ($[0,1]$)*, Contributions to Nonstandard Analysis, Edited by W.A.J. Luxemburg and A. Robinson, North Holland Publ. Co., Amsterdam 1972.
3. J. Gapaillard, *Relevements, Bases de derivation ensemblistes et fonctionnelles*, These, Nantes 1972.
4. S. Graf, *Schnitte Boolescher Korrespondenzen und ihre Dualisierungen*, Dissertation, Erlangen 1973.
5. A. Ionescu Tulcea and C. Ionescu Tulcea, *Topics in the Theory of Lifting*, Springer Verlag, Berlin-Heidelberg — New York 1969.
6. J. von Neumann and M. H. Stone, *The determination of representative elements in the Residual classes of a Boolean algebra*, Fund. Math., **25** (1935), 353–376.
7. M. Sion, *A theory of Semigroup Valued Measure*, Lect. Notes in Mathematics, **355** (1973), Springer Verlag, Berlin-Heidelberg — New York.
8. T. Traynor, *An Elementary Proof of the Lifting Theorem*, Pacific J. Math., **53** (1974), 267–272.

Received May 13, 1974.

UNIVERSITY OF BRITISH COLUMBIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1975 Pacific Journal of Mathematics
All Rights Reserved

Pacific Journal of Mathematics

Vol. 58, No. 2

April, 1975

Zvi Artstein and John Allen Burns, <i>Integration of compact set-valued functions</i>	297
Mark Benard, <i>Characters and Schur indices of the unitary reflection group [321]³</i>	309
Simeon M. Berman, <i>A new characterization of characteristic functions of absolutely continuous distributions</i>	323
Monte Boisen and Philip B. Sheldon, <i>Pre-Prüfer rings</i>	331
Hans-Heinrich Brungs, <i>Three questions on duo rings</i>	345
Iracema M. Bund, <i>Birnbaum-Orlicz spaces of functions on groups</i>	351
John D. Elwin and Donald R. Short, <i>Branched immersions between 2-manifolds of higher topological type</i>	361
Eric Friedlander, <i>Extension functions for rank 2, torsion free abelian groups</i>	371
Jon Froemke and Robert Willis Quackenbush, <i>The spectrum of an equational class of groupoids</i>	381
Barry J. Gardner, <i>Radicals of supplementary semilattice sums of associative rings</i>	387
Shmuel Glasner, <i>Relatively invariant measures</i>	393
George Rudolph Gordh, Jr. and Sibe Mardesic, <i>Characterizing local connectedness in inverse limits</i>	411
Siegfried Graf, <i>On the existence of strong liftings in second countable topological spaces</i>	419
Stanley P. Gudder and D. Strawther, <i>Orthogonally additive and orthogonally increasing functions on vector spaces</i>	427
Darald Joe Hartfiel and Carlton James Maxson, <i>A characterization of the maximal monoids and maximal groups in β_X</i>	437
Robert E. Hartwig and S. Brent Morris, <i>The universal flip matrix and the generalized faro-shuffle</i>	445
William Emery Haver, <i>Mappings between ANRs that are fine homotopy equivalences</i>	457
J. Bockett Hunter, <i>Moment sequences in l^p</i>	463
Barbara Jeffcott and William Thomas Spears, <i>Semimodularity in the completion of a poset</i>	467
Jerry Alan Johnson, <i>A note on Banach spaces of Lipschitz functions</i>	475
David W. Jonah and Bertram Manuel Schreiber, <i>Transitive affine transformations on groups</i>	483
Karsten Juul, <i>Some three-point subset properties connected with Menger's characterization of boundaries of plane convex sets</i>	511
Ronald Brian Kirk, <i>The Haar integral via non-standard analysis</i>	517
Justin Thomas Lloyd and William Smiley, <i>On the group of permutations with countable support</i>	529
Erwin Lutwak, <i>Dual mixed volumes</i>	531
Mark Mahowald, <i>The index of a tangent 2-field</i>	539
Keith Miller, <i>Logarithmic convexity results for holomorphic semigroups</i>	549
Paul Milnes, <i>Extension of continuous functions on topological semigroups</i>	553
Kenneth Clayton Pietz, <i>Cauchy transforms and characteristic functions</i>	563
James Ted Rogers Jr., <i>Whitney continua in the hyperspace $C(X)$</i>	569
Jean-Marie G. Rolin, <i>The inverse of a continuous additive functional</i>	585
William Henry Ruckle, <i>Absolutely divergent series and isomorphism of subspaces</i>	605
Rolf Schneider, <i>A measure of convexity for compact sets</i>	617
Alan Henry Schoenfeld, <i>Continuous measure-preserving maps onto Peano spaces</i>	627
V. Merriline Smith, <i>Strongly superficial elements</i>	643
Roger P. Ware, <i>A note on quadratic forms over Pythagorean fields</i>	651
Roger Allen Wiegand and Sylvia Wiegand, <i>Finitely generated modules over Bezout rings</i>	655
Martin Ziegler, <i>A counterexample in the theory of definable automorphisms</i>	665