

Pacific Journal of Mathematics

ON M -PROJECTIVE AND M -INJECTIVE MODULES

GORO AZUMAYA, F. MBUNTUM AND KALATHOOR VARADARAJAN

ON M -PROJECTIVE AND M -INJECTIVE MODULES

G. AZUMAYYA, F. MBUNTUM AND K. VARADARAJAN

In this paper necessary and sufficient conditions are obtained for a direct sum $\bigoplus_{\alpha \in J} A_\alpha$ of R -modules to be M -injective in the sense of Azumaya. Using this result it is shown that if $\{A_\alpha\}_{\alpha \in J}$ is a family of R -modules with the property that $\bigoplus_{\alpha \in K} A_\alpha$ is M -injective for every countable subset K of J then $\bigoplus_{\alpha \in J} A_\alpha$ is itself M -injective. Also we prove that arbitrary direct sums of M -injective modules are M -injective if and only if M is locally noetherian, in the sense that every cyclic submodule of M is noetherian. We also obtain some structure theorems about Z -projective modules in the sense of Azumaya, where Z denotes the ring of integers. Writing any abelian group A as $D \oplus H$ with D divisible and H reduced, we show that if A is Z -projective then H is torsion free and every pure subgroup of finite rank of H is a free direct summand of H .

Most of these results were motivated by the results of B. Sarath and K. Varadarajan regarding injectivity of direct sums.

1. M -projective and M -injective modules. Throughout this paper R denotes a ring with $1 \neq 0$ and all the modules considered are left unitary modules over R . By an ideal in R we mean a left ideal in R . M denotes a fixed R -module. We first recall the notions of M -projective and M -injective modules originally introduced by one of the authors [1].

DEFINITION 1.1. An R -module H is called M -projective, if given a diagram

$$\begin{array}{ccc} & H & \\ & \downarrow f & \\ M \xrightarrow{\varphi} & N & \longrightarrow 0 \end{array}$$

of maps of R -modules with the horizontal sequence exact, \exists a map $h: H \rightarrow M$ such that $\varphi \circ h = f$.

The notion of an M -injective module is defined dually.

REMARK 1.2. Regarding R as a left module over itself in the usual way it turns out that R -injective modules are the same as the injective modules over R . However R -projective modules are not the same as projective modules over R .

LEMMA 1.3. *Every divisible abelian group D is Z -projective.*

Proof. Trivial consequence of the fact $\text{Hom}(D, Z) = 0 = \text{Hom}(D, Z_k)$ whenever D is divisible.

REMARK 1.4. We know that projective modules over Z are free. Hence no divisible abelian group $D \neq 0$ is projective over Z .

LEMMA 1.5. *Suppose H is a torsion free abelian group with the property that every pure subgroup of rank 1 of H is a free direct summand of H . Then every pure subgroup of finite rank of H is also a free direct summand of H .*

Proof. By induction on the rank of the subgroup. Let S be a pure subgroup of H of rank k with $k > 1$. We can pick a pure subgroup B of S of rank 1. Then B is also pure in H and hence by assumption B is free abelian and $H = C \oplus B$ for some C . Since $S \supset B$ we get $S = (S \cap C) \oplus B$. Now $S \cap C$ is of rank $(k - 1)$ and pure in S and hence pure in H . By the inductive hypothesis $S \cap C$ is free abelian and $H = (S \cap C) \oplus L$ for some L . From $C \supset S \cap C$ we now $C = (S \cap C) \oplus (L \cap C)$. Thus $S = (S \cap C) \oplus B$ is free abelian and

$$\begin{aligned} H &= C \oplus B = (S \cap C) \oplus (L \cap C) \oplus B \\ &= (S \cap C) \oplus B \oplus (L \cap C) = S \oplus (L \cap C). \end{aligned}$$

DEFINITION 1.6. We say that a torsion free abelian group H has property (P) if every pure subgroup of finite rank of H is free and a direct summand of H .

Given any abelian group A we can write A as $D \oplus H$ where D is the maximal divisible subgroup of A and H is reduced. Also $H \cong A/D$ is well-determined up to an isomorphism. We will refer to any group isomorphic to H as the reduced part of A .

THEOREM 1.7. *Suppose H is reduced abelian group which is Z -projective. Then H is torsion-free with property (P).*

Proof. It is well-known that a reduced abelian group which is not torsion-free admits of a nonzero finite cyclic direct summand [3, Th 9, p. 21]. Clearly the identity map $Z_m \rightarrow Z_m$ (for $m \geq 1$) can not be lifted to a map $Z_m \rightarrow Z$. This proves that Z_m is not Z -projective. Hence if a reduced abelian group H is Z -projective it has to be torsion free.

For any $a \neq 0$ in H let $S_a = \{x \in H \mid x \text{ and } a \text{ are linearly dependent over } Z\}$. Then it is trivial to see that S_a is a pure subgroup of

rank 1 in H . Moreover S_a is reduced since H is. Hence \exists a prime p such that $S_a \neq pS_a$. Let $c \in S_a$ be such that $c \notin pS_a$. Since S_a is a pure subgroup of H we see that $c \notin pH$. Hence $\eta(c) \neq 0$ where $\eta: H \rightarrow H/pH$ denotes the canonical quotient map. Regarding H/pH as a vector space over Z_p we can complete $\eta(c)$ to a basis $\{\eta(c)\} \cup \{u_j\}_{j \in J}$ of H/pH over Z_p . Let $\theta: H/pH \rightarrow Z_p$ be the Z_p -linear map determined by $\theta(\eta(c)) = 1 \in Z_p$ and $\theta(u_j) = 0$ for all $j \in J$. The Z -projectivity of H now yields a map $h: H \rightarrow Z$ with

$$\begin{array}{ccc}
 & H & \\
 & \downarrow \theta \circ \eta & \\
 Z & \xrightarrow{\varphi} & Z_p \longrightarrow 0 \\
 & \nwarrow h & \\
 & &
 \end{array}$$

commutative, where $\varphi: Z \rightarrow Z_p$ is the canonical quotient map. From $\varphi h(c) = \theta \circ \eta(c) = 1 \in Z_p$ it follows that $\varphi h(c) \neq 0$. Hence $g = h|_{S_a}: S_a \rightarrow Z$ is a non-zero homomorphism. It follows that $\text{Im } g = kZ$ for some integer $k \geq 1$. Composing g with the obvious isomorphism $kZ \cong Z$ we get an epimorphism $g': S_a \rightarrow Z$. Since Z is free the sequence $S_a \xrightarrow{g'} Z \rightarrow 0$ splits. S_a being a torsion-free group of rank 1 it now follows that $S_a \xrightarrow{g'} Z$ is an isomorphism. Thus for $\alpha \neq 0$ in H the subgroup S_α is isomorphic to Z .

Our next step is to show that S_a is a direct summand of H . Let c be a generator for $S_a \cong Z$ and $V = \{\alpha \in \text{Hom}(H, Z) \mid \alpha(c) \neq 0\}$. From what we have seen already V is a nonempty set. Let $l = \min_{\alpha \in V} |\alpha(c)|$. We will show that $l = 1$. Suppose on the contrary $l > 1$. There definitely exists an element $\alpha \in V$ such that $\alpha(c) = l$. Let p be a prime divisor of l and $l = kp$. Now $c \notin pS_a$. The argument used already yields a map $h: H \rightarrow Z$ such that $\varphi h(c) = 1 \in Z_p$. This means $h(c) = np + 1$ for some $n \in Z$. Writing $n = kd + r$ with $d \in Z$ and r an integer satisfying $0 \leq r < k$ consider the element $h - d\alpha \in \text{Hom}(H, Z)$. Now, $\{h - d\alpha\}(c) = np + 1 - dl = np + 1 - dkp = rp + 1$. Clearly, $0 < rp + 1 < rp + p = (r + 1)p \leq kp = l$. Thus $\beta = h - d\alpha$ is in V and $|\beta(c)| = rp + 1 < l$, contradicting the definition of l . This contradiction proves that $h = 1$. It now follows that \exists an $\alpha: H \rightarrow Z$ with $\alpha(c) = 1$, in which case \exists a splitting $\mu: Z \rightarrow H$ for α with $\mu(1) = c$. Hence $S_a = \mu(Z)$ is a direct summand of H .

It is clear that every pure subgroup of rank 1 of H is of the form S_α for some $\alpha \neq 0$ in H . Now appealing to Lemma 1.5 we immediately see that H has property (P).

COROLLARY 1.8. *Let $A = D \oplus H$ with D the maximal divisible subgroup of A . If A is Z -projective then H is torsion-free and*

has property (P).

COROLLARY 1.9. *A finitely generated abelian group A is Z -projective $\Leftrightarrow A$ is free of finite rank.*

COROLLARY 1.10 *Suppose H is a reduced decomposable torsion-free abelian group. (i.e., H is the direct sum of rank 1 torsion-free abelian groups). Then H is Z -projective $\Leftrightarrow H$ is free.*

PROPOSITION 1.11. *Let p be a prime. An abelian group A is Z_{p^∞} -injective if and only if $A \cong (\bigoplus_{\alpha \in J} Z_{p^\infty}) \oplus B$, a direct sum of copies of Z_{p^∞} with an abelian group B having no p -torsion.*

Proof. Suppose $A \cong (\bigoplus_{\alpha \in J} Z_{p^\infty}) \oplus B$ with B having no p -torsion. Since $\bigoplus_{\alpha \in J} Z_{p^\infty}$ is divisible, it is injective over Z and hence Z_{p^∞} -injective as well. The only subgroups of Z_{p^∞} are Z_{p^∞} and Z_{p^k} for some integer $k \geq 1$. When B has no p -torsion $\text{Hom}(Z_{p^k}, B) = 0 = \text{Hom}(Z_{p^\infty}, B)$. This proves that B is Z_{p^∞} -injective.

Conversely, assume A to be Z_{p^∞} -injective. Let $\alpha \in A$ be an element in the p -primary torsion of A . Suppose the order of α is p^k . Then \exists a homomorphism $Z_{p^k} \xrightarrow{f} A$ carrying the element 1 of Z_{p^k} to α . Since A is Z_{p^∞} -injective \exists an extension $g: Z_{p^\infty} \rightarrow A$ of f . Then $\text{Im } g$ is divisible, $\alpha \in \text{Im } g$ and $\text{Im } g$ is in the p -primary torsion of A . This proves that the p -primary torsion of A is divisible. Since any divisible subgroup of A is a direct summand of A and since any divisible p -primary abelian group is a direct sum of copies of Z_{p^∞} it follows that $A \cong (\bigoplus_{\alpha \in J} Z_{p^\infty}) \oplus B$ with B having no p -torsion.

We now recall the definitions of an M -epimorphism and an M -monomorphism due to one of the authors [1], and state two results due to him.

DEFINITION 1.12. (i) Let A, B be R -modules and $\theta: A \rightarrow B$ an epimorphism. θ is said to be an M -epimorphism if \exists a map $\psi: A \rightarrow M$ such that $\text{Ker } \theta \cap \text{Ker } \psi = 0$.

(ii) Let $\alpha: A \rightarrow B$ be a monomorphism. α is called an M -monomorphism if \exists a map $\beta: M \rightarrow B$ such that $\text{Im } \alpha$ and $\text{Im } \beta$ together generate B .

PROPOSITION 1.13 [1], [5]. *The following conditions on an R -module H are equivalent.*

- (1) H is M -projective
- (2) Given any M -epimorphism $\theta: A \rightarrow B$ and any $f: H \rightarrow B \in$ a map $h: H \rightarrow A$ such that $\theta \circ h = f$
- (3) Every M -epimorphism $\theta: C \rightarrow H$ splits.

PROPOSITION 1.14. *Dual of Proposition 1.13.*

DEFINITION 1.15. For any module H let $C^p(H)$ (respy $C^i(H)$) = the class of all modules M such that H is M -projective (respy M -injective). For any module M let $C_p(M)$ (respy $C_i(M)$) denote the class of M -projective (respy M -injective) modules.

PROPOSITION 1.16 [1], [5].

(1) $C^p(H)$ is closed under submodules, homomorphic images and the formation of finite direct sums.

(2) $C^i(H)$ is closed under submodules, homomorphic images and arbitrary direct sums.

(3) $C_p(H)$ (respy $C_i(H)$) is closed under direct sums (respy direct products) and direct summands (respy direct factors)

REMARKS.

1.17. In general $C^p(H)$ is not closed under formation of arbitrary direct sums. For instance let $R = Z$ and $H = Q$ the additive group of the rationals. From Lemma 1.3 we see that Q is Z -projective. Thus $Z \in C^p(Q)$. Let J be an *infinite* set and for each $\alpha \in J$ let $M_\alpha = Z$. Then each $M_\alpha \in C^p(Q)$. Clearly Q is a quotient of $\bigoplus_{\alpha \in J} M_\alpha$ and the identity map of Q can not be lifted to a map of Q into $\bigoplus_{\alpha \in J} M_\alpha$. This means $\bigoplus_{\alpha \in J} M_\alpha \notin C^p(Q)$.

1.17'. Since $C^p(H)$ is closed under submodules from 1.17 it follows that $C^p(H)$ in general is not closed under formation of arbitrary direct products.

1.18. In general $C^i(H)$ is not closed under formation of arbitrary direct products. Let $R = Z$ and $H = Z$. From Proposition 1.11 we have $Z_{p^\infty} \in C^i(Z)$. Let $M = \prod_p Z_{p^\infty}$, the direct product taken over all primes. It is known and quite easy to see that \exists a subgroup of M which is isomorphic to Q . If $M \in C^i(Z)$ from (2) of Proposition 1.16 it would that $Q \in C^i(Z)$. Since the identity map of Z can not be extended to a map of Q into Z it follows that Z is not Q -injective. In other words $Q \notin C^i(Z)$. This in turn implies $M \notin C^i(Z)$.

2. M -injectivity of direct sums. For any module A and any $x \in A$ we denote the left annihilator $\{\lambda \in R \mid \lambda x = 0\}$ of x by L_x .

DEFINITION 2.1. An element $x \in A$ is said to be dominated by M if $L_x \supset L_m$ for some $m \in M$.

Given a family $\{A\}_{\alpha \in J}$ of modules let x be the element of $\prod_{\alpha \in J} A_\alpha$ whose α -component is x_α . Let $I_x = \{\lambda \in R \mid \lambda x \in \bigoplus_{\alpha \in J} A_\alpha\}$.

DEFINITION 2.2. We call $x \in \prod_{\alpha \in J} A_\alpha$ a special element if $I_x x_\alpha =$

0 for almost all α . In other words \exists a finite subset F of J such that $\lambda x_\alpha = 0$ for all $\lambda \in I_x$ and for all $\alpha \notin F$.

PROPOSITION 2.3. *A is M -injective $\Leftrightarrow A$ is Rm -injective for all $m \in M$.*

Proof. This is an easy consequence of 1.16 (2). The implication \Rightarrow follows from the closedness of $C^i(A)$ under submodules. As for \Leftarrow , by the closedness of $C^i(A)$ under direct sums it follows that A is $\bigoplus_{m \in M} Rm$ -injective. Since M is a homomorphic image of $\bigoplus_{m \in M} Rm$ and since $C^i(A)$ is closed under homomorphic images, it follows that A is M -injective.

THEOREM 2.4. *$\bigoplus_{\alpha \in J} A_\alpha$ is M -injective \Leftrightarrow each A_α is M -injective and every element of $\prod_{\alpha \in J} A_\alpha$ dominated by M is special.*

Proof. \Rightarrow : Let $x \in \pi A_\alpha$ be dominated by M , that is, there is an $m \in M$ such that $L_m \subset L_x$. This implies that the mapping $\lambda m \rightarrow \lambda x (\lambda \in R)$ is well defined and gives a homomorphism $f: Rm \rightarrow \pi A_\alpha$. The image of the submodule $I_x m$ by f is clearly $I_x x (\subset \bigoplus A_\alpha)$. Thus the restriction of f to $I_x m$ is regarded as a homomorphism $I_x m \rightarrow \bigoplus A_\alpha$. Since $\bigoplus A_\alpha$ is Rm -injective, this homomorphism can be extended to a homomorphism $Rm \rightarrow \bigoplus A_\alpha$ which means that there exists an $u \in \bigoplus A_\alpha$ such that $\lambda x = \lambda u$ for all $\lambda \in I_x$. It follows then that $I_x x \alpha = I_x u \alpha$ for all $\alpha \in J$. But since $u \alpha = 0$ for almost all α , it follows that $I_x x \alpha = 0$ for almost all α too, i.e., x is special.

\Leftarrow : Let $m \in M$ and consider the cyclic submodule Rm of M . Let I be a left ideal of R . Then IM is a submodule of Rm . (Conversely every submodule of Rm is of the form Im with a suitable left ideal I). Let there be given a homomorphism $h: Im \rightarrow \bigoplus A_\alpha$. Then since $\bigoplus A_\alpha \subset \pi A_\alpha$ and πA_α is M -whence Rm -injective, h can be extended to a homomorphism $Rm \rightarrow \pi A_\alpha$. Let $x \in \pi A_\alpha$ be the image of m . Then the homomorphism is given by $\lambda m \rightarrow \lambda x (\lambda \in R)$. Therefore it follows that $Ix = h(IM) \subset \bigoplus A_\alpha$ whence $I \subset I_x$. On the other hand, since clearly $L_m \subset L_x$, x is dominated by M and thus x is special by assumption, i.e., $I_x x_\alpha = 0$ whence $Ix_\alpha = 0$ for almost all α . Let u be the element of $\bigoplus A_\alpha$ whose α -component is x_α or 0 according as $Ix_\alpha \neq 0$ or $Ix_\alpha = 0$. Then it is clear that $\lambda u = \lambda x$ for all $\lambda \in I$. Further, it is also clear that $L_m \subset L_x \subset L_u$ and therefore the mapping $\lambda m \rightarrow \lambda u (\lambda \in R)$ is well defined. This mapping gives a homomorphism $f: Rm \rightarrow \bigoplus A_\alpha$ which is an extension of h , because $f(\lambda m) = \lambda u = \lambda x$ for all $\lambda \in I$. This implies that $\bigoplus A_\alpha$ is Rm -injective and so is M -injective (by Proposition 2.3).

THEOREM 2.5. *The direct sum of any family of M -injective modules is M -injective \Leftrightarrow every cyclic submodule of M is noetherian.*

Proof. \Leftarrow . Let $\{A_\alpha\}$ be a family of M -injective modules. Let x be an element of πA_α dominated by M ; thus there is an $m \in M$ such that $L_m \subset L_x$. Consider $I_x m$. Since clearly $L_x \subset I_x$ whence $L_m \subset I_x$, it follows that $I_x/L_m \cong I_x m$. On the other hand, $I_x m$ is a submodule of the Noetherian module Rm . Hence I_x/L_m is finitely generated, i.e., there exist a finite number of elements $\lambda_1, \lambda_2, \dots, \lambda_n$ of I_x such that

$$I_x = R\lambda_1 + R\lambda_2 + \dots + R\lambda_n + L_m.$$

It follows therefore $I_x x_\alpha = R\lambda_1 x_\alpha + R\lambda_2 x_\alpha + \dots + R\lambda_n x_\alpha$ for all components x_α . Since, however, for each i , $\lambda_i x_\alpha = 0$ for almost all α , it follows that $I_x x_\alpha = 0$ for almost all α , that is, x is special. Thus $\bigoplus A_\alpha$ is M -injective by Theorem 2.4.

\Rightarrow . Let Rm , $m \in M$ be any cyclic submodule of M . Then $R/L_m \cong Rm$, and there is a (1-1) correspondence between the left ideals of R containing L_m and submodules of Rm . Thus in order to show that Rm is noetherian it is sufficient to prove that there is no properly ascending infinite sequence of ideals of R containing L_m . Suppose there exists an infinite sequence $L_m \subset I_1 \subset I_2 \subset I_3 \subset \dots$ of ideals I_j with $I_j \neq I_{j+1}$ for every $j \geq 1$. Let $B_j = R/I_j$, $\eta_j: R \rightarrow B_j$ the canonical projection. Let A_j be the injective hull of B_j . Then each A_j is M -injective also. By assumption \exists an $m \in M$ s.t. $I_1 \supset L_m$. The element $x = (x_j)_{j \geq 1}$ of $\prod_{j \geq 1} A_j$ where $x_j = \eta_j(1)$ is clearly dominated by M . For any $\lambda \in I_j$ we have $\lambda x_k = 0$ for $k \geq j$. Hence $I_j \subset I_x$ for all $j \geq 1$. Let λ_j be any element of I_{j+1} which is not in I_j . Then $\lambda_j x_j \neq 0$ and $\lambda_j \in I_x$. This proves that $I_x x_j \neq 0$ for every $j \geq 1$. This means x is not a special element and hence by theorem 2.4, $\bigoplus_{j \geq 1} A_j$ is not M -injective. This proves the implication \Rightarrow .

REMARK 2.6. A result of H. Bass [2] asserts that arbitrary direct sums of injective modules over R are injective $\Leftrightarrow R$ is noetherian. Theorem 2.5 is a generalization of this result of H. Bass. When $M = R$ we get the result of Bass.

THEOREM 2.7. *Suppose $\{A_\alpha\}_{\alpha \in J}$ is a family of R -modules such that for every countable subset K of J , $\bigoplus_{\alpha \in K} A_\alpha$ is M -injective. Then $\bigoplus_{\alpha \in J} A_\alpha$ is itself M -injective.*

Proof. Assume that $\bigoplus_{\alpha \in J} A_\alpha$ is not M -injective. Then, by Theorem 2.4, there exists an $x \in \prod_{\alpha \in J} A_\alpha$ which is dominated by M but is not special, i.e., $I_x x_\alpha \neq 0$ for infinitely many $\alpha \in J$. Let K be

an infinite countable subset of the infinite set $\{\alpha \in J \mid I_x x_\alpha \neq 0\}$. Let \mathbf{y} be element of $\prod_{\alpha \in K} A_\alpha$ whose α -component y_α is equal to x_α for all $\alpha \in K$. Then clearly $I_x \subset I_y$, so that it follows that \mathbf{y} is dominated by M and $I_y y_\alpha = I_y x_\alpha \neq 0$ for all $\alpha \in K$. This implies again by Theorem 2.4 that $\bigoplus_{\alpha \in K} A_\alpha$ is not M -injective (because each A_α is M -injective by the assumption of our theorem). This is a contradiction, and so the proof is completed.

REFERENCES

1. G. Azumaya, *M-projective and M-injective modules* (unpublished).
2. H. Bass, *Injective dimension in Noetherian rings*, Trans. Amer. Math. Soc., **102** (1962), 18-29.
3. I. Kaplansky, *Infinite Abelian Groups*, University of Michigan Press (1969).
4. B. Sarath and K. Varadarajan, *Injectivity of direct sums*; Communications in Algebra, **1** (6), (1974), 517-530.
5. M. S. Shrikhande, *On hereditary and cohereditary modules*, Canad. J. Math., **XXV** (1973), 892-896.

Received June 19, 1974. Research done while the last mentioned author was partially supported by NRC grant A 8225.

INDIANA UNIVERSITY

AND

THE UNIVERSITY OF CALGARY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Shashi Prabha Arya and M. K. Singal, <i>More sum theorems for topological spaces</i>	1
Goro Azumaya, F. Mbuntum and Kalathoor Varadarajan, <i>On M-projective and M-injective modules</i>	9
Kong Ming Chong, <i>Spectral inequalities involving the infima and suprema of functions</i>	17
Alan Hetherington Durfee, <i>The characteristic polynomial of the monodromy</i>	21
Emilio Gagliardo and Clifford Alfons Kottman, <i>Fixed points for orientation preserving homeomorphisms of the plane which interchange two points</i>	27
Raymond F. Gittings, <i>Finite-to-one open maps of generalized metric spaces</i>	33
Andrew M. W. Glass, W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>a^*-closures of completely distributive lattice-ordered groups</i>	43
Matthew Gould, <i>Endomorphism and automorphism structure of direct squares of universal algebras</i>	69
R. E. Harrell and Les Andrew Karlovitz, <i>On tree structures in Banach spaces</i>	85
Julien O. Hennefeld, <i>Finding a maximal subalgebra on which the two Arens products agree</i>	93
William Francis Keigher, <i>Adjunctions and comonads in differential algebra</i>	99
Robert Bernard Kelman, <i>A Dirichlet-Jordan theorem for dual trigonometric series</i>	113
Allan Morton Krall, <i>Stieltjes differential-boundary operators. III. Multivalued operators—linear relations</i>	125
Hui-Hsiung Kuo, <i>On Gross differentiation on Banach spaces</i>	135
Tom Louton, <i>A theorem on simultaneous observability</i>	147
Kenneth Mandelberg, <i>Amitsur cohomology for certain extensions of rings of algebraic integers</i>	161
Coy Lewis May, <i>Automorphisms of compact Klein surfaces with boundary</i>	199
Peter A. McCoy, <i>Generalized axisymmetric elliptic functions</i>	211
Muril Lynn Robertson, <i>Concerning Siu's method for solving $y'(t) = F(t, y(g(t)))$</i>	223
Richard Lewis Roth, <i>On restricting irreducible characters to normal subgroups</i>	229
Albert Oscar Shar, <i>P-primary decomposition of maps into an H-space</i>	237
Kenneth Barry Stolarsky, <i>The sum of the distances to certain pointsets on the unit circle</i>	241
Bert Alan Taylor, <i>Components of zero sets of analytic functions in C^2 in the unit ball or polydisc</i>	253
Michel Valadier, <i>Convex integrands on Souslin locally convex spaces</i>	267
Januario Varela, <i>Fields of automorphisms and derivations of C^*-algebras</i>	277
Arnold Lewis Villone, <i>A class of symmetric differential operators with deficiency indices $(1, 1)$</i>	295
Manfred Wollenberg, <i>The invariance principle for wave operators</i>	303