

Pacific Journal of Mathematics

**CONCERNING SIU'S METHOD FOR SOLVING $y'(t) = F(t,$
 $y(g(t)))$**

MURIL LYNN ROBERTSON

CONCERNING SIU'S METHOD FOR SOLVING

$$y'(t) = F(t, y(g(t)))$$

MURIL L. ROBERTSON

A procedure is given, which is parameterized by a certain set of real-valued functions, that yields sufficient conditions on each of g and F to guarantee a solution to $y'(t) = F(t, y(g(t)))$.

The following is the main result.

THEOREM 1. *Suppose that f is a real-valued continuous function with connected domain J of real numbers so that*

- (1) $0 \in J$ and $f(0) > 0$
- (2) f is increasing on $J \cap [0, +\infty)$
- (3) f is decreasing on $J \cap (-\infty, 0]$
- (4) $0 < k < 1$, if the range of f is unbounded; and $k = 1$ if the range of f is bounded
- (5) B is a Banach space, $F: J \times B \rightarrow B$, and $N: J \rightarrow \mathbb{R}$
- (6) F is continuous and there is a constant C so that

$$\left| \int_0^x \|F(s, 0)\| ds \right| \leq C f(x), \text{ for all } x \text{ in } J.$$

- (7) $\|F(t, x) - F(t, y)\| \leq N(t) \|x - y\|$, for $t \in J$ and $x, y \in B$
- (8) N is positive and Lebesgue integrable or subintervals of J
- (9) g is any continuous function from J into J so that $g(x) \in f^{-1}[f(0), k|f'^{\text{sign}x}(x)|/N(x)]$ for all $x \in J$. ($f'^{\text{sign}x}$ denotes the right-hand derivative if $x > 0$ and it denotes the left-hand derivative if $x < 0$.)
- (10) $q \in B$.

Then, there is a unique function $y: J \rightarrow B$ so that $y'(t) = F(t, y(g(t)))$, $y(0) = q$ and $\|y(t)\| \leq \text{Constant} \cdot f(t)$, for all t in J .

LEMMA. *If f satisfies conditions (1), (2), and (3) in the statement of Theorem 1, then $\int_0^x f'^{\text{sign}x}(s) ds$ exists in the Lebesgue sense and is less than or equal to $f(x) - f(0)$, for each $x \in J$.*

Proof of Lemma. Suppose $x > 0$. Let $f_n(s) = [f(s + 1/n) - f(s)]n$. Then $f'^+(s) = \lim f_n(s)$ for almost all $s > 0$. Clearly each f_n is summable, because each f_n is continuous. Also, for each n

$$\begin{aligned} \int_0^x f_n(s) ds &= \int_0^x n \left[f\left(s + \frac{1}{n}\right) - f(s) \right] ds \\ &= n \int_{1/n}^{x+1/n} f(s) ds - n \int_0^x f(s) ds \end{aligned}$$

$$\begin{aligned}
&= n \int_x^{x+1/n} f(s) ds - n \int_0^{1/n} f(s) ds \quad (\text{which approaches} \\
&\hspace{15em} f(x) - f(0)) \\
&\leq 2 \cdot \{\text{sup of } f \text{ on } [0, x + 1]\}
\end{aligned}$$

Thus, by Fatou's lemma [see 3, page 39], f'^+ is summable on $[0, x]$ for all $x > 0$ and $x \in J$ and

$$\int_0^x f'^+(s) ds \leq \liminf \int_0^x f_n(s) ds \leq f(x) - f(0).$$

The proof is similar for $x < 0$.

Proof of Theorem 1. Let $\|\cdot\|$ be the norm of B and define $|\cdot|$ to be the norm defined by $|z| = \sup\{\|z(x)\|/f(x) : x \in J\}$ for each z continuous from J into B such that this supremum exists. Let Y denote the Banach space of all such z , with norm $|\cdot|$. For each $z \in Y$ and $x \in J$, let $(Tz)(x) = q + \int_0^x F(s, z(g(s))) ds$. Suppose $z, w \in Y$. Then

$$\begin{aligned}
\|(Tz)(x) - (Tw)(x)\| &= \left\| \int_0^x [F(s, z(g(s))) - F(s, w(g(s)))] ds \right\| \\
&\leq \left| \int_0^x N(s) \|z(g(s)) - w(g(s))\| ds \right| \\
&\leq \left| \int_0^x N(s) f(g(s)) ds \right| |z - w|.
\end{aligned}$$

Thus, $|Tz - Tw| \leq \sup \left\{ \left| \int_0^x N(s) f(g(s)) ds \right| / f(x) \right\} |z - w|$. The following shows that T is a contraction.

Case 1. Suppose $x > 0$. Then if $0 \leq s \leq x$, $f(g(s)) \in [f(0), kf'^+(s)/N(s)]$. Thus, $\left| \int_0^x N(s) f(g(s)) ds \right| = \int_0^x N(s) f(g(s)) ds \leq \int_0^x kf'^+(s) ds \leq k(f(x) - f(0))$, by the lemma.

Case 2. Suppose $x < 0$. Then if

$$x \leq s \leq 0, f(g(s)) \in [f(0), -kf'^-(s)/N(s)].$$

Thus,

$$\left| \int_0^x N(s) f(g(s)) ds \right| = \int_x^0 N(s) f(g(s)) ds \leq s k \int_0^x f'^-(s) ds \leq k(f(x) - f(0)),$$

by the lemma.

Thus, in either case $c = \sup \left\{ \left| \int_0^x N(x) f(g(s)) ds \right| / f(x) \right\} \leq \sup \{k(1 - f(0)/f(x))\}$. So, if the range of f is unbounded, $c \leq k$ and if the range of f is bounded by L and $k = 1$, then $c \leq 1 - f(0)/L$. So T

has contraction constant c . The zero function Z is in Y , because $(TZ)(x) = q + \int_0^x F(s, 0)ds$ and $\|(TZ)(x)\|/f(x) \leq [\|q\| + Cf(x)]/f(x) \leq \|q\|/f(0) + C$. Now if $w \in Y$, $\|(Tw)(x)\|/f(x) \leq |Tw - TZ| + |TZ|$. Thus $Tw \in Y$. So, by Banach's contraction mapping principle, T has a unique fixed point in Y . This proves Theorem 1.

REMARKS. (1) Given any g one may find an appropriate N and apply Theorem 1, by requiring $N(x) \leq k|f'^{\text{sign}x}(x)|/f(g(x))$.

(2) At any particular x , there is an f so that $f'^{\text{sign}x}(x) = \infty$ and so that t does not have infinite derivative in a deleted neighborhood of x . For this type f , $g(x)$ could be any number.

In [5], Siu essentially uses the method of Theorem 1 with $f(x) = \exp(|x|/e)$ to obtain:

THEOREM 2 (Siu). *If $|g(x)| \leq |x| + c$, where $0 < c < 1/e$, for all real numbers x , then $y' = y(g)$, $y(0) = q$ has unique solution subject to $\|y(x)\| \leq \text{constant} \cdot \exp(|x|/e)$*

In [4], the author proves:

THEOREM 3. *If $\{I(i)\}$ is a sequence of intervals so that $I(0) = \{0\} \subseteq I(i) \subseteq I(i + 1)$, $I(i) = [a(i), b(i)]$, and $\max\{a(i - 1) - a(i), b(i) - b(i - 1)\} < 1$ for each positive integer i ; then $y' = y(g)$, $y(0) = q$ has unique solution on $\cup\{I(i)\}$, whenever g is continuous and $g(I(i)) \subseteq I(i)$ for each positive integer i .*

The following theorem is comparable to each of Theorem 2 and Theorem 3.

THEOREM 4. *Suppose the hypothesis of Theorem 3 holds and k is in $(0, 1)$ such that $\max\{a(i - 1) - a(i), b(i) - b(i - 1)\} < k$ for each positive integer i . Then, for each positive integer i , there exists $\delta(i) > 0$ such that if g is a continuous function from $\cup\{I(i)\}$ into $\cup\{I(i)\}$ such that $g(I(i)) \subseteq [a(i) - \delta(i), b(i) + \delta(i)]$, then $y'(t) = F(t)$, $y(g(t))$, $y(0) = q$ has a solution on $\cup\{I(i)\}$ for any F such that $N = 1$, where F and N satisfy the conditions listed for them in the hypothesis of Theorem 1.*

Proof of Theorem 4. Let f be a positive continuous piecewise linear function with domain $\cup\{I(i)\}$ such that f has slope $M(i)$ or $(b(i - 1), b(i))$ and slope $-M(i)$ on $(a(i), a(i - 1))$ where the sequence $\{M(i)\}$ is chosen such that for each nonnegative integer n ,

$M(n+1)$

$$> \max \left\{ \left[f(0) + \sum_{i=1}^n M(i)(a(i-1) - a(i)) \right] / [k - (a(n) - a(n+1))], \right. \\ \left. \left[f(0) + \sum_{i=1}^n M(i)(b(i) - b(i-1)) \right] / [k - (b(n+1) - b(n))] \right\}.$$

Let

$$\delta(n+1) = \min \left\{ a(n+1) - a(n+2), b(n+2) - b(n+1), \right. \\ \left. \left[kM(n+1) - \left[f(0) + \sum_{i=1}^{n+1} M(i)(a(i-1) - a(i)) \right] \right] / M(n+2), \right. \\ \left. \left[kM(n+1) - \left[f(0) + \sum_{i=1}^{n+1} M(i)(b(i) - b(i-1)) \right] \right] / M(n+2) \right\}.$$

It follows that the hypotheses of Theorem 1 hold.

REMARK. The solution in Theorem 4 is unique in the Banach space Y of Theorem 1, which depends on f .

The following is a straightforward application of Theorem 1.

THEOREM 5. Suppose

$$(1) \quad -1 < -k < a < 0 < b < k < 1$$

$$(2) \quad m \geq \max \{1/(k+a), 1/(k-b)\}$$

$$(3) \quad nk \geq \max \{(1-ma)/(1+mb), (1+mb)(1-ma)\}$$

$$(4) \quad l(x) = \begin{cases} x - (\log nk)/n & , \quad x \leq a \\ a - (\log km)/(1-ma)/n & , \quad a < x < b \\ -x + a + b - (\log kn(mb+1))/(1-ma)/n & , \quad b \leq x \end{cases}$$

and

$$u(x) = \begin{cases} -x + a + b + (\log nk)/(1-ma)/(1+mb)/n & , \quad x \leq a \\ b + (\log km)/(1+mb)/n & , \quad a < x < b \\ x + (\log nk)/n & , \quad b \leq x \end{cases}$$

$$(5) \quad l \leq g \leq u \text{ and } g \text{ is continuous.}$$

Then, there is a unique solution to $y'(t) = F(t, y(g(t)))$, $y(0) = q$ where F and N satisfy the hypotheses of Theorem 1, $N=1$ and, $\|y(x)\| \leq \text{constant} \cdot f(x)$, for all real numbers x , where

$$f(x) = \begin{cases} (1-ma) \exp(-n(x-a)) & , \quad x \leq a \\ 1-mx & , \quad a \leq x \leq 0 \\ 1+mx & , \quad 0 \leq x \leq b \\ (mb+1) \exp(n(x-b)) & , \quad b \leq x \end{cases}$$

REMARK. If $a = -b$, then $u = -l$.

The following is a generalization of a theorem by D. R. Anderson [1].

COROLLARY TO THEOREM 5. Suppose $0 < \beta < 1$, $\varepsilon > 0$, and

$$|g(x)| \leq \begin{cases} -x + \frac{1}{e} - \varepsilon & x \leq -\beta \\ \beta + \left(\log \frac{1}{\beta}\right) / e - \varepsilon & -\beta < x < \beta \\ x + \frac{1}{e} - \varepsilon & \beta \leq x. \end{cases}$$

Then, there is a solution to $y'(t) = F(t, y(g(t)))$, $y(0) = q$ for $N = 1$ any y subject to $\|y(x)\| \leq \text{constant} \cdot f(x)$ for an appropriate f .

Proof. Straightforward.

REMARK. As β approaches 0, g is allowed to become indefinitely large at 0.

The author wishes to thank Professor Bob Dorroh for his help with the lemma to Theorem 1.

REFERENCES

1. D. R. Anderson, *An existence theorem for a solution of $f'(x) = F(x, f(g(x)))$* , SIAM Review, **8** (1966), 359-362.
2. R. J. Oberg, *On the local existence of solutions of certain functional-differential equations*, Trans. Amer. Math. Soc., (1969), 295-302.
3. F. Riesz, and B. Sz. Nagy, *Functional Analysis*, Ungar, New York, 1955.
4. M. L. Robertson, *The equation $y'(t) = F(t, y(g(t)))$* , Pacific J. Math., **43** (1972), 483-491.
5. Y. T. Siu, *On the solution of the equation $f'(x) = \lambda f(g(x))$* , Math. Z., **90** (1965) 391, -392.

Received April 16, 1973 and in revised form March 6, 1975.

AUBURN UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 59, No. 1

May, 1975

Shashi Prabha Arya and M. K. Singal, <i>More sum theorems for topological spaces</i>	1
Goro Azumaya, F. Mbuntum and Kalathoor Varadarajan, <i>On M-projective and M-injective modules</i>	9
Kong Ming Chong, <i>Spectral inequalities involving the infima and suprema of functions</i>	17
Alan Hetherington Durfee, <i>The characteristic polynomial of the monodromy</i>	21
Emilio Gagliardo and Clifford Alfons Kottman, <i>Fixed points for orientation preserving homeomorphisms of the plane which interchange two points</i>	27
Raymond F. Gittings, <i>Finite-to-one open maps of generalized metric spaces</i>	33
Andrew M. W. Glass, W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>a^*-closures of completely distributive lattice-ordered groups</i>	43
Matthew Gould, <i>Endomorphism and automorphism structure of direct squares of universal algebras</i>	69
R. E. Harrell and Les Andrew Karlovitz, <i>On tree structures in Banach spaces</i>	85
Julien O. Hennefeld, <i>Finding a maximal subalgebra on which the two Arens products agree</i>	93
William Francis Keigher, <i>Adjunctions and comonads in differential algebra</i>	99
Robert Bernard Kelman, <i>A Dirichlet-Jordan theorem for dual trigonometric series</i>	113
Allan Morton Krall, <i>Stieltjes differential-boundary operators. III. Multivalued operators—linear relations</i>	125
Hui-Hsiung Kuo, <i>On Gross differentiation on Banach spaces</i>	135
Tom Louton, <i>A theorem on simultaneous observability</i>	147
Kenneth Mandelberg, <i>Amitsur cohomology for certain extensions of rings of algebraic integers</i>	161
Coy Lewis May, <i>Automorphisms of compact Klein surfaces with boundary</i>	199
Peter A. McCoy, <i>Generalized axisymmetric elliptic functions</i>	211
Muril Lynn Robertson, <i>Concerning Siu's method for solving $y'(t) = F(t, y(g(t)))$</i>	223
Richard Lewis Roth, <i>On restricting irreducible characters to normal subgroups</i>	229
Albert Oscar Shar, <i>P-primary decomposition of maps into an H-space</i>	237
Kenneth Barry Stolarsky, <i>The sum of the distances to certain pointsets on the unit circle</i>	241
Bert Alan Taylor, <i>Components of zero sets of analytic functions in C^2 in the unit ball or polydisc</i>	253
Michel Valadier, <i>Convex integrands on Souslin locally convex spaces</i>	267
Januario Varela, <i>Fields of automorphisms and derivations of C^*-algebras</i>	277
Arnold Lewis Villone, <i>A class of symmetric differential operators with deficiency indices $(1, 1)$</i>	295
Manfred Wollenberg, <i>The invariance principle for wave operators</i>	303