

# Pacific Journal of Mathematics

## **NORMS OF RANDOM MATRICES**

**GRAHAME BENNETT, VICTOR WAYNE GOODMAN  
AND CHARLES MICHAEL NEWMAN**

## NORMS OF RANDOM MATRICES

G. BENNETT, V. GOODMAN<sup>1</sup>, AND C. M. NEWMAN<sup>2</sup>

Under rather general conditions on the matrix entries, we obtain estimates for the probability distribution of the norm of a random matrix transformation from  $\ell_n^2$  to  $\ell_m^q$  ( $2 \leq q < \infty$ ). Asymptotically, the expected norm is remarkably small and this enables us to produce an interesting class of bounded linear operators from  $\ell^2$  to  $\ell^q$ . As an application, we complete the characterization of  $(p, q)$ -absolutely summing operators on Hilbert space, thereby answering a question left open by several previous authors.

1. Introduction. Many questions in the theory of  $\ell^p$  spaces require, for their solution, the existence of finite matrices with  $\pm 1$  entries whose norms satisfy prescribed conditions. In several cases the required matrices have been given explicitly: the simplest examples stem from the orthogonality of the Walsh functions (see, for example, [10] where non-complemented subspaces of  $\ell^p$  are constructed) or from the Rademacher functions via Khintchine's inequality (see, for example, [6] where the  $p$ -absolutely summing operators on Hilbert space are characterized). In many problems, however, the construction of suitable matrices leads to formidable combinatorial difficulties. We consider in this paper one such problem for which no constructive method is available. The appropriate matrix is obtained here probabilistically by showing that "most" matrices satisfy the prescribed norm inequalities.

Specifically, the problem we consider is that of characterizing the ideal,  $\Pi_{p,q}$ , of  $(p, q)$ -absolutely summing operators on Hilbert space. Recall that a bounded linear operator  $T$  on  $\ell^2$  is  $(p, q)$ -absolutely summing ( $1 \leq q \leq p \leq \infty$ ) if  $(\|Tx_n\|)_{n=1}^\infty \in \ell^p$  whenever  $(x_n)_{n=1}^\infty$  is a sequence of elements of  $\ell^2$  with the property that  $(\langle x_n, y \rangle)_{n=1}^\infty \in \ell^q$  for each  $y \in \ell^2$ . This problem has received a good deal of attention in recent years and the known results are described below. We denote by  $\mathfrak{S}_r$  ( $1 \leq r < \infty$ ) the Schatten  $r$ -class of all compact linear operators  $T$  on  $\ell^2$  for which  $\sum_{n=1}^\infty |\lambda_n|^r < \infty$  where  $\{\lambda_n\}_{n=1}^\infty$  are the eigenvalues of  $(T^*T)^{1/2}$ , counted according to multiplicities (and arranged in order of decreasing modulus); for convenience the class of all bounded linear operators on  $\ell^2$  is denoted by  $\mathfrak{S}_\infty$ .

We then have:

- (a) if  $p = q < \infty$ ,  $\Pi_{p,q} = \mathfrak{S}_2$ ;
- (b) if  $p = \infty$  or  $(1/q) - (1/p) \geq 1/2$ ,  $\Pi_{p,q} = \mathfrak{S}_\infty$ ;

- (c) if  $(1/q) - (1/p) < 1/2$  and  $q \leq 2$ ,  $\Pi_{p,q} = \mathfrak{S}_r$  with  $1/r = (1/p) - (1/q) + (1/2)$ ;
- (d) if  $2 < q < p < \infty$ ,  $\mathfrak{S}_{2p/q} \subseteq \Pi_{p,q} \subseteq \mathfrak{S}_p$ .

(a) is due to Pełczyński [6]; (b) to Kwapien [4]; (c) and the first part of (d) to Mitiagin [4]; and the second part of (d) to Pietsch-Triebel [8]. Special cases were discovered earlier by Grothendieck ((a) for  $p = 1$ ) [3]; Pietsch ((a) for  $1 \leq p \leq 2$ ) [7]; and Orlicz ((b) for  $p = 2$ ,  $q = 1$ ) [5].

The only outstanding case is thus  $2 < q < p < \infty$ . This case is more subtle than the others in that a new ideal,  $\mathfrak{S}_{r,s}$ , generated by the Lorentz sequence space  $\ell^{r,s}$  is involved.  $\mathfrak{S}_{r,s}$  ( $1 < r, s < \infty$ ) is the set of all compact linear operators  $T$  on  $\ell^2$  for which  $\sum_{n=1}^{\infty} n^{(s/r)-1} |\lambda_n|^s < \infty$  where the  $\lambda_n$ 's are defined as above. In [1], it is shown that

- (e) if  $2 < q < p < \infty$ ,  $\mathfrak{S}_{2p/q,p} \subseteq \Pi_{p,q}$ , with equality when  $q$  is an even integer;

we here remove the restriction that  $q$  be an even integer, thereby completing the description of  $\Pi_{p,q}$ . This is done in Section 3 of the paper by using the following result whose proof is given in Section 2.

**THEOREM 1.** *Let  $A = (a_{ij})$  be an  $m \times n$  matrix whose entries are independent, mean-zero random variables with  $|a_{ij}| \leq 1$  for all  $i, j$ . For  $2 \leq q < \infty$ , there is a constant  $K$ , depending only on  $q$ , such that*

$$E(\|A\|_{2,q}) \leq K \max(m^{1/q}, n^{1/2})$$

where  $\|A\|_{2,q}$  denotes the operator norm of  $A: \ell_n^2 \rightarrow \ell_m^q$ .

It should be noted that the estimate of Theorem 1 is best possible (up to the choice of  $K$ ) in the sense that every  $m \times n$  matrix  $A' = (a'_{ij})$  with  $a'_{ij} = \pm 1$  has  $\|A'\|_{2,q} \geq \max(m^{1/q}, n^{1/2})$ .

**2. Random matrices.** Theorem 1 is proved below using the following lemmas. Our techniques were suggested by methods used in deriving limit theorems for large deviations of sums of random variables ([2] and [9]). We begin with a standard result whose proof is included for completeness.

**LEMMA 1.** *Let  $(X_j)_{j=1}^n$  be independent, mean-zero random variables with  $|X_j| \leq 1$  for all  $j$ ; then for any  $\lambda > 0$  and real  $b_1, \dots, b_n$ ,*

$$P\left(\left|\sum_{j=1}^n b_j X_j\right| \geq \lambda\right) \leq 2 \exp\left(-\lambda^2/4 \sum_{j=1}^n b_j^2\right).$$

*Proof.* Using the elementary inequality,  $\exp(x) - x \leq \exp(x^2)$ , it follows that for all real  $\mu$ ,

$$E[\exp(\mu X_j)] = E[\exp(\mu X_j) - \mu X_j] \leq E[\exp(\mu^2 X_j^2)] \leq \exp(\mu^2).$$

The independence of  $(X_j)_{j=1}^n$  thus yields

$$E\left[\exp\left(\mu \sum_{j=1}^n b_j X_j\right)\right] \leq \exp\left(\mu^2 \sum_{j=1}^n b_j^2\right).$$

Applying Chebyshev's inequality, we obtain, for  $\mu > 0$ ,

$$P\left(\sum_{j=1}^n b_j X_j \geq \lambda\right) \leq \exp\left(\mu^2 \sum_{j=1}^n b_j^2 - \lambda\mu\right).$$

Taking  $\mu = \lambda/2 \sum_{j=1}^n b_j^2$  gives the desired result.

The next lemma is really the technical key to our results (at least for  $q \neq 2$ ).

**LEMMA 2.** *For each  $q \geq 2$ , there is a constant  $C$ , depending only on  $q$ , so that if  $(X_j)_{j=1}^n$  satisfy the hypotheses of Lemma 1,*

$$E\left[\exp\left(\mu \left|\sum_{j=1}^n b_j X_j\right|^q\right)\right] \leq 1 + C\mu \left(\sum_{j=1}^n b_j^2\right)^{q/2}$$

for  $0 \leq \mu \leq (\sum_{j=1}^n b_j^2)^{-q/2} n^{1-q/2}/8$ .

*Proof.* Without loss of generality we may take  $\sum_{j=1}^n b_j^2 = 1$ .

$$\begin{aligned} E\left[\exp\left(\mu \left|\sum_{j=1}^n b_j X_j\right|^q\right)\right] &= \int_0^\infty \exp(\mu\lambda^q) dP\left(\left|\sum_{j=1}^n b_j X_j\right| \leq \lambda\right) \\ &= 1 + \int_0^\infty q\mu\lambda^{q-1} \exp(\mu\lambda^q) P\left(\left|\sum_{j=1}^n b_j X_j\right| > \lambda\right) d\lambda \end{aligned}$$

via integration by parts. Since  $|\sum_{j=1}^n b_j X_j| \leq (\sum_{j=1}^n b_j^2)^{1/2} n^{1/2}$ , we have  $P(|\sum_{j=1}^n b_j X_j| > n^{1/2}) = 0$ . We apply the estimate from Lemma 1 to obtain

$$\begin{aligned} E\left[\exp\left(\mu \left|\sum_{j=1}^n b_j X_j\right|^q\right)\right] &\leq 1 + 2q\mu \int_0^{n^{1/2}} \lambda^{q-1} \exp(\mu\lambda^q - \lambda^2/4) d\lambda \\ &\leq 1 + 2q\mu \int_0^\infty \lambda^{q-1} \exp(-\lambda^2/8) d\lambda \end{aligned}$$

since  $0 \leq \mu \leq n^{1-q/2}/8$ . This gives the desired result with

$$C = 2q \int_0^\infty \lambda^{q-1} \exp(-\lambda^2/8) d\lambda = 8^{q/2} q \Gamma(q/2).$$

We now assume that  $q \geq 2$  is fixed and that  $\{a_{ij}\}$  are independent random variables satisfying the hypotheses of Lemma 1. Given real  $x_j$  we obtain estimates (independent of  $\{x_j\}$ ) on the probability distribution for the random variable

$$Y = \sum_{i=1}^m \left| \sum_{j=1}^n x_j a_{ij} \right|^q / \left( \sum_{j=1}^n x_j^2 \right)^{q/2} .$$

LEMMA 3. For any real  $\lambda$  and positive integers  $m, n$ ,

$$P(Y \geq Cm + 8\lambda n^{q/2}) \leq \exp(-\lambda n) ,$$

where  $C$  is the constant appearing in Lemma 2.

*Proof.* For real  $\mu$ , we set  $K(\mu) = \log E[\exp(\mu Y)]$  so that  $E[\exp(\mu Y - K(\mu))] = 1$ . It follows from Chebyshev's inequality that for any real  $\nu$ ,

$$P(\mu Y \geq K(\mu) + \nu) \leq \exp(-\nu) .$$

On the other hand, we have from Lemma 2 that for  $0 \leq \mu \leq n^{1-q/2}/8$ ,

$$\begin{aligned} E\left[\exp(\mu Y)\right] &= \prod_{i=1}^m \left[ E\left[\exp\left(\mu \left|\sum_{j=1}^n b_j a_{ij}\right|^q\right)\right] \right] \\ &\leq \prod_{i=1}^m (1 + C\mu) \leq \exp(mC\mu) , \end{aligned}$$

where  $b_j = x_j / (\sum_{j=1}^n x_j^2)^{1/2}$  so that  $K(\mu) \leq mC\mu$ . Setting  $\nu = \lambda n$  and  $\mu = n^{1-q/2}/8$ , we obtain the desired result.

We now consider the random  $m \times n$  matrix  $A$  with entries  $a_{ij}$  and denote the norm of  $A: \ell_n^2 \rightarrow \ell_m^q$  by

$$\|A\|_{2,q} = \sup_{\sum_{j=1}^n x_j^2 = 1} \left( \sum_{i=1}^m \left| \sum_{j=1}^n a_{ij} x_j \right|^q \right)^{1/q} = \sup_{\|x\|_2 = 1} (\|Ax\|_q) .$$

LEMMA 4. There exist constants  $c_1, c_2$  (depending at most on  $q$ ) such that for all  $\lambda > 0$ ,

$$P(\|A\|_{2,q} \geq c_1(m + \lambda n^{q/2})^{1/q}) \leq \exp(-(\lambda - c_2)n) .$$

*Proof.* Let  $0 < \varepsilon < 1$  be fixed; by an  $\varepsilon$ -net for the unit sphere  $S = \{x: \|x\|_2 = 1\}$  in  $\mathbf{R}^n$ , we mean a finite subset,  $N$ , of  $S$  satisfying

$$\sup_{x \in S} \min_{y \in N} \|x - y\|_2 < \varepsilon .$$

Any matrix  $A$  effectively attains its norm on such a set  $N$ : more precisely, we have

$$\|A\|_{2,q} = \sup_{x \in S} \|Ax\|_q \leq \max_{y \in N} \|Ay\|_q + \sup_{x \in S} \min_{y \in N} \|A(x - y)\|_q,$$

so that

$$\|A\|_{2,q} \leq \frac{1}{1 - \varepsilon} \max_{y \in N} \|Ay\|_q.$$

For such an  $N$ , we have, putting  $c_1 = (\max(C, 8))^{1/q}/(1 - \varepsilon)$ ,

$$\begin{aligned} P(\|A\|_{2,q} \geq c_1(m + \lambda n^{q/2})^{1/q}) &\leq P\left(\max_{y \in N} \|Ay\|_q \geq (Cm + 8\lambda n^{q/2})^{1/q}\right) \\ &\leq \sum_{y \in N} \exp(-\lambda n) = |N| \exp(-\lambda n) \end{aligned}$$

by Lemma 3, where  $|N|$  denotes the number of elements in  $N$ . Moreover, using elementary geometrical arguments, it is straightforward to show that  $N$  may be chosen with  $|N| \leq \exp(c_2 n)$  with  $c_2$  a constant depending only on  $\varepsilon$ ; this completes the proof.

We can now proceed to prove Theorem 1.

*Proof of Theorem 1.* We let  $\mu_1 = c_1^q(m + c_2 n^{q/2})$ ; then by Jensen's inequality,

$$\begin{aligned} (E[\|A\|_{2,q}])^q &\leq E[(\|A\|_{2,q})^q] \\ &= \int_0^\infty P[(\|A\|_{2,q})^q \geq \mu] d\mu \\ &\leq \mu_1 + \int_{\mu_1}^\infty P[\|A\|_{2,q} \geq \mu^{1/q}] d\mu \\ &\leq \mu_1 + \int_{\mu_1}^\infty \exp\{-n^{1-q/2}(-m + \mu/c_1^q) + c_2 n\} d\mu \\ &= c_1^q(m + c_2 n^{q/2} + n^{(q/2)-1}), \end{aligned}$$

from which the desired result immediately follows.

Before concluding this section we apply Lemma 4 in a somewhat different manner to obtain the following result.

**THEOREM 2.** For  $2 \leq q < \infty$ , there is a constant  $K'$ , depending only on  $q$ , so that if (for each  $m, n = 1, 2, \dots$ )  $A_{m,n}$  is an  $m \times n$  random matrix satisfying the hypotheses of Theorem 1, then with probability one,

$$\limsup_{\max(m,n) \rightarrow \infty} \|A_{m,n}\|_{2,q} / \max(m^{1/q}, n^{1/2}) \leq K'.$$

*Proof.* Using the estimate of Lemma 4 with  $\lambda = c_2 + \max(1, m/n^{q/2})$  gives

$$P(\|A_{m,n}\|_{2,q}/\max(m^{1/q}, n^{1/2}) \geq K') \leq \exp(-\max(n, mn^{1-q/2}))$$

with  $K' = c_1(2 + c_2)^{1/q}$ . By the Borel-Cantelli lemma,

$$P(\|A_{m,n}\|_{2,q}/\max(m^{1/q}, n^{1/2}) \geq K' \text{ for infinitely many } m, n) = 0$$

if  $\sum_{m,n} P(\|A_{m,n}\|_{2,q}/\max(m^{1/q}, n^{1/2}) \geq K') < \infty$ . Hence it suffices to show that

$$\sum_{m,n=1}^{\infty} \exp(-\max(n, mn^{1-q/2})) < \infty,$$

which is easily verified.

REMARK. Defining for an  $m \times n$  matrix  $A$ ,

$$L_{2,q}(A) = \inf_{\|x\|_2=1} \|Ax\|_q,$$

we have of course that  $0 \leq L_{2,q}(A) \leq \|A\|_{2,q}$  with  $L_{2,q}(A) = 0$  for  $m < n$ . Estimates similar to those obtained above can be used to derive asymptotic lower bounds for  $L_{2,q}(A)$  (at least with  $m$  much larger than  $n$ ). It can be shown, for example, that if  $\inf_{i,j} E[(a_{ij})^2] > 0$ , then for  $2 < q < \infty$  and any  $\delta > 0$ , there exists a constant  $d > 0$  so that with probability one,

$$\liminf_{\substack{m \rightarrow \infty \\ m \geq \delta n^{q/2}}} L_{2,q}(A_{m,n})/m^{1/q} \geq d.$$

3. Absolutely summing operators. In this section we complete the description of the  $(p, q)$ -absolutely summing operators on Hilbert space, thereby answering a question left open in [4], [8] and [1].

THEOREM 3. If  $2 < q < p < \infty$ , then  $\Pi_{p,q} = \mathfrak{S}_{2p/q,p}$ .

*Proof.* The inclusion  $\Pi_{p,q} \supseteq \mathfrak{S}_{2p/q,p}$  has already been established in Theorem I of [1]. For the converse, we apply Theorem 1 (or Lemma 4 or Theorem 2), choosing the matrix entries  $(a_{ij})$  independently with  $P(a_{ij} = +1) = P(a_{ij} = -1) = 1/2$ . It follows that there exists, for each positive integer  $n$ , at least one matrix of order  $[n^{q/2}] \times n$  with all  $\pm 1$  entries satisfying  $\|A\|_{2,q} \leq Kn^{1/2}$ , where  $K$  is a constant depending only on  $q$ . This generalizes Proposition 2 of [1], and the argument used to prove Theorem II of that paper shows that  $\Pi_{p,q} \subseteq \mathfrak{S}_{2p/q,p}$ .

*Added in proof.* For further applications of these results the reader should consult the forthcoming paper "On uncomplemented subspaces  $L^p$ ,  $1 < p < 2$ , being prepared jointly with L. E. Dor and W. B. Johnson.

For extensions to matrix transformations of  $\ell^p$  into  $\ell^q$ ,  $1 \leq p$ ,  $q \leq \infty$ , consult "Hadamard multipliers," being prepared by G. Bennett.

## REFERENCES

1. G. Bennett, Some ideals of operators on Hilbert space, to appear in *Studia Math.*
2. S. N. Bernstein, *The Theory of Probability*, Gostekhizdat, 1946.
3. A. Grothendieck, Résumé de la théorie métrique des produits tensoriels topologiques, *Boletim Soc. Mat. Sao Paulo* **8** (1956), 1-79.
4. S. Kwapien, Some remarks on  $(p, q)$ -absolutely summing operators in  $\ell_p$  spaces, *Studia Math.* **29** (1968), 327-337.
5. W. Orlicz, Über unbedingte Konvergenz in Funktionenräumen, II. *Studia Math.* **4** (1933), 41-47.
6. A. Pełczyński, A characterization of Hilbert-Schmidt operators, *Studia Math.* **28** (1967), 355-360.
7. A. Pietsch, Absolut  $p$ -summierende Abbildungen in normierten Räumen, *Studia Math.* **28** (1967), 333-353.
8. A. Pietsch und H. Triebel, Interpolationstheorie für Banachideale von beschränkten linearen Operatoren, *Studia Math.* **31** (1968), 95-109.
9. W. Richter, Refinement of an inequality of S. N. Bernstein, *Vestnik Leningrad Univ.* **1** (1959), 24-29.
10. A. Sobczyk, Projections in Minkowski and Banach spaces, *Duke Math. J.* **8** (1941), 78-106.

Received February 20, 1975. The second author was supported in part by NSF grant MPS-73-08715, and the third author was supported in part by NSF grant MPS-74-04870.

INDIANA UNIVERSITY





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)

University of California  
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

# Pacific Journal of Mathematics

Vol. 59, No. 2

June, 1975

Aharon Atzmon, <i>A moment problem for positive measures on the unit disc</i> . . . . .	317
Peter W. Bates and Grant Bernard Gustafson, <i>Green's function inequalities for two-point boundary value problems</i> . . . . .	327
Howard Edwin Bell, <i>Infinite subrings of infinite rings and near-rings</i> . . . . .	345
Grahame Bennett, Victor Wayne Goodman and Charles Michael Newman, <i>Norms of random matrices</i> . . . . .	359
Beverly L. Brechner, <i>Almost periodic homeomorphisms of <math>E^2</math> are periodic</i> . . . . .	367
Beverly L. Brechner and R. Daniel Mauldin, <i>Homeomorphisms of the plane</i> . . . . .	375
Jia-Arng Chao, <i>Lusin area functions on local fields</i> . . . . .	383
Frank Rimi DeMeyer, <i>The Brauer group of polynomial rings</i> . . . . .	391
M. V. Deshpande, <i>Collectively compact sets and the ergodic theory of semi-groups</i> . . . . .	399
Raymond Frank Dickman and Jack Ray Porter, <i><math>\theta</math>-closed subsets of Hausdorff spaces</i> . . . . .	407
Charles P. Downey, <i>Classification of singular integrals over a local field</i> . . . . .	417
Daniel Reuven Farkas, <i>Miscellany on Bieberbach group algebras</i> . . . . .	427
Peter A. Fowler, <i>Infimum and domination principles in vector lattices</i> . . . . .	437
Barry J. Gardner, <i>Some aspects of <math>T</math>-nilpotence. II: Lifting properties over <math>T</math>-nilpotent ideals</i> . . . . .	445
Gary Fred Gruenhage and Phillip Lee Zenor, <i>Metrization of spaces with countable large basis dimension</i> . . . . .	455
J. L. Hickman, <i>Reducing series of ordinals</i> . . . . .	461
Hugh M. Hilden, <i>Generators for two groups related to the braid group</i> . . . . .	475
Tom (Roy Thomas Jr.) Jacob, <i>Some matrix transformations on analytic sequence spaces</i> . . . . .	487
Elyahu Katz, <i>Free products in the category of <math>k_w</math>-groups</i> . . . . .	493
Tsang Hai Kuo, <i>On conjugate Banach spaces with the Radon-Nikodým property</i> . . . . .	497
Norman Eugene Liden, <i><math>K</math>-spaces, their antispaces and related mappings</i> . . . . .	505
Clinton M. Petty, <i>Radon partitions in real linear spaces</i> . . . . .	515
Alan Saleski, <i>A conditional entropy for the space of pseudo-Menger maps</i> . . . . .	525
Michael Singer, <i>Elementary solutions of differential equations</i> . . . . .	535
Eugene Spiegel and Allan Trojan, <i>On semi-simple group algebras. I</i> . . . . .	549
Charles Madison Stanton, <i>Bounded analytic functions on a class of open Riemann surfaces</i> . . . . .	557
Sherman K. Stein, <i>Transversals of Latin squares and their generalizations</i> . . . . .	567
Ivan Ernest Stux, <i>Distribution of squarefree integers in non-linear sequences</i> . . . . .	577
Lowell G. Sweet, <i>On homogeneous algebras</i> . . . . .	585
Lowell G. Sweet, <i>On doubly homogeneous algebras</i> . . . . .	595
Florian Vasilescu, <i>The closed range modulus of operators</i> . . . . .	599
Arthur Anthony Yanushka, <i>A characterization of the symplectic groups <math>PSp(2m, q)</math> as rank 3 permutation groups</i> . . . . .	611
James Juei-Chin Yeh, <i>Inversion of conditional Wiener integrals</i> . . . . .	623