# Pacific Journal of Mathematics

SOME ASPECTS OF *T*-NILPOTENCE. II: LIFTING PROPERTIES OVER *T*-NILPOTENT IDEALS

BARRY J. GARDNER

Vol. 59, No. 2

June 1975

# SOME ASPECTS OF *T*-NILPOTENCE II: LIFTING PROPERTIES OVER *T*-NILPOTENT IDEALS

## B. J. GARDNER

It has been shown by Năstăsescu and Popescu that every nonzero (left, unital) module over a ring R has a simple submodule if and only if the Jacobson radical J of R is right T-nilpotent and every nonzero R/J-module has a simple submodule. The work presented here arose largely from an attempt to find a general framework for results like this.

In §2 it is shown that if R has a right T-nilpotent ideal I, then a bijection from the torsion classes of R/I-modules to those of R-Modules can be obtained by associating with each  $\mathscr{T} \subseteq Mod(R/I)$  the lower radical class it defines as a class of R-modules. §3 contains applications involving the lifting of torsion properties and in §4 it is shown that if R has a right T-nilpotent ideal I such that R/I is the direct sum of its torsion and divisible ideals, then R has this property also.

In the present paper we continue the general investigation of T-nilpotence begun in [8]. The emphasis here is on T-nilpotent ideals, whereas previously we looked at whole rings, but there are some methodological similarities between the two papers.

1. Preliminaries. Except where we state otherwise, all rings considered are associative and have identities (though some results hold more generally) and modules are always left, unital modules. An ideal I (one or two-sided) is left T-nilpotent if for every sequence  $a_1, a_2, \cdots$  of its elements,  $a_1a_2 \cdots a_n = 0$  for some *n*. Right T-nilpotence is defined analogously. The terms are due to Bass [2], though the concepts originated with Levitzki [19]. The terminology relating to the radical theory of modules varies somewhat; we shall describe things in the following way: An ordered pair  $(\mathcal{R}, \mathcal{S})$  of classes of modules over a ring will be called a radical theory if (i)  $\mathscr{R} \cap \mathscr{S} = \{0\}$ , (ii)  $\mathscr{R}$  is homomorphically closed, (iii)  $\mathscr{S}$  is hereditary (i.e. closed under taking submodules), (iv) every module M has a largest submodule  $\mathscr{R}(M)$  from  $\mathscr{R}$  and (v)  $M/\mathscr{R}(M) \in \mathscr{S}$  for all M. In this situation,  $\mathscr{R}$  will be called a radical class,  $\mathscr{S}$  a semi-simple class. If also  $\mathcal{R}$  is hereditary, it will be called a torsion class, while if  $\mathcal{S}$  is homomorphically closed it will be called a TTF class.  $L_{\mathbb{R}}(\mathcal{M})$  is the lower radical class defined by a class  $\mathcal{M}$  of R-modules. With these conventions established, we refer the reader to [4], [5] and [14] for further details. Mod(R) is the class of all modules

over a ring R. (0: M) is the annihilator of a module M.

2. Torsion from iterated annihilators. Until further notice, I is a right ideal of a ring R.

For any *R*-module *M*, we define submodules  $M^{(\alpha)}$  for all ordinals  $\alpha$ , as follows:

$$egin{aligned} M^{\scriptscriptstyle(0)} &= 0 ext{; } M^{\scriptscriptstyle(lpha+1)} &= \{m \in M \,|\, Im \sqsubseteq M^{\scriptscriptstyle(lpha)}\} ext{ ; } \ M^{\scriptscriptstyle(eta)} &= igcup_{lpha < eta} M^{\scriptscriptstyle(lpha)} ext{, if } eta ext{ is a limit .} \end{aligned}$$

Eventually,  $M^{(\mu+1)} = M^{(\mu)}$ . If  $M^{(\mu)} = M$ , then

$$0=M^{\scriptscriptstyle(0)}\subseteq M^{\scriptscriptstyle(1)}\subseteq\cdots\subseteq M^{\scriptscriptstyle(lpha)}\subseteq M^{\scriptscriptstyle(lpha+1)}\subseteq\cdots\subseteq M^{\scriptscriptstyle(\mu)}=M$$

is called the upper ascending *I*-series of *M*. More generally, a transfinite chain

$$0 = M_{0} \subseteq M_{1} \subseteq \cdots \subseteq M_{\alpha} \subseteq M_{\alpha+1} \subseteq \cdots \subseteq M_{\mu} = M$$

of submodules is called an ascending I-series for M if

 $IM_{lpha+1} \subseteq M_{lpha} ext{ for each } lpha ext{ and } \bigcup_{lpha < eta} M_{lpha} = M_{eta} ext{ for all limits } eta ext{ .}$ 

Finally,  $\mathcal{M}_I = \{M \in \text{Mod } (R) | IM = 0\}.$ 

THEOREM 2.1. The following conditions are equivalent for a module M:

(i) The upper ascending I-series of M exists.

(ii) M has an ascending I-series.

(iii) Every nonzero homomorphic image of M has a nonzero submodule in  $\mathcal{M}_{I}$ .

(iv) For any  $x \in M$  and  $a_1, a_2, \dots \in I$ , there exists n such that  $a_n a_{n-1} \cdots a_2 a_1 x = 0$ .

 $(\mathbf{v}) \quad M \in L_R(\mathcal{M}_I).$ 

Proof. (i)  $\Rightarrow$  (ii) is clear. (ii)  $\Rightarrow$  (iii): If  $N \subseteq M$ , let

$$\beta = \min \left\{ \alpha \, | \, M_{\alpha} \not\subseteq N \right\}$$

for some ascending *I*-series

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_{\alpha} \subseteq M_{\alpha+1} \subseteq \cdots \subseteq M_{\mu} = M$$
.

Then  $\beta \neq 0$  and  $\beta$  is not a limit, so  $(M_{\beta} + N)/N$  is a nonzero submodule of M/N with  $I(M_{\beta} + N)/N \subseteq (M_{\beta-1} + N)N = 0$ .

(ii)  $\Rightarrow$  (iv): With the same notation, for  $x \in M$  and  $a_1, a_2, \dots \in I$ 

446

$$\gamma = \min \left\{ lpha \, | \, M_lpha \, ext{ contains some } a_n a_{n-1} \cdots a_2 a_1 x 
ight\}$$
 ,

and let  $a_m a_{m-1} \cdots a_2 a_1 x$  be in  $M_{\gamma}$ . Then  $\gamma$  is not a limit, and if  $\gamma \neq 0$ , then  $a_{m+1}(a_m a_{m-1} \cdots a_2 a_1 x) \in IM_{\gamma} \subseteq M_{\gamma-1}$ , contrary to the assumed minimality of  $\gamma$ . Hence  $\gamma = 0$ , so  $a_m a_{m-1} \cdots a_2 a_1 x = 0$ .

(iv)  $\Rightarrow$  (iii): Let  $0 \neq x \in M$ . If  $Ix \neq 0$ , let  $a_1x \neq 0$ , where  $a_1 \in I$ . If  $Ia_1x \neq 0$ , let  $a_2a_1x \neq 0$ , where  $a_2 \in I$ . Proceeding thus, we obtain an element  $b = a_na_{n-1} \cdots a_2a_1x \neq 0$  where  $a_1, a_2, \cdots, a_n \in I$  and Ib = 0. Thus M has a submodule which is annihilated by I. Since property (iv) is preserved in homomorphic images, the result follows.

(iii)  $\Rightarrow$  (i): Let  $M^{(\mu)} = M^{(\mu+1)}$ . Then  $[M/M^{(\mu)}]^{(1)} = M^{(\mu+1)}/M^{(\mu)} = 0$ , so  $M/M^{(\mu)} = 0$ .

The equivalence of (iii) and (v) is well-known.

The proof just given is patterned after some proofs in [8]. Kashu [15], working with modules over a ring R not necessarily having an identity, has shown that  $L_R(\mathscr{Z})$ , where  $\mathscr{Z}$  is the class of modules with trivial multiplication, consists precisely of those modules M such that for every sequence  $a_1, a_2, \cdots$  in R and for every  $x \in M$ ,  $a_n a_{n-1} \cdots a_2 a_1 x = 0$  eventually. Kellett [16], in the same setting, has shown that  $L_R(\mathscr{Z})$  is the class of modules with an upper ascending R-series.

Since  $L_R(\mathcal{M}_I) = Mod(R)$  if and only if  $R \in L_R(\mathcal{M}_I)$  we have the following consequence of Theorem 2.1.

COROLLARY 2.2.  $L_R(\mathcal{M}_I) = Mod(R)$  if and only if I is right T-nilpotent.

PROPOSITION 2.3.  $L_{\mathbb{R}}(\mathscr{M}_{I})$  is a torsion class. Moreover, if  $\mathscr{R}$  is a torsion subclass of  $L_{\mathbb{R}}(\mathscr{M}_{I})$ , then  $\mathscr{R} = L_{\mathbb{R}}(\mathscr{R} \cap \mathscr{M}_{I})$ .

*Proof.* Since  $\mathscr{M}_I$  is hereditary, the first assertion is clear. If  $\mathscr{R}$  is a torsion subclass of  $L_{\mathbb{R}}(\mathscr{M}_I)$ , then clearly  $L_{\mathbb{R}}(\mathscr{R} \cap \mathscr{M}_I) \subseteq \mathscr{R}$ . On the other hand, if  $M \in \mathscr{R} \subseteq L_{\mathbb{R}}(\mathscr{M}_I)$ , then every nonzero homomorphic image of M has a non zero submodule  $N \in \mathscr{M}_I$ . But any such N is in  $\mathscr{R}$ , as the latter is a torsion class. Thus each nonzero homomorphic image of M has a nonzero submodule in  $\mathscr{R} \cap \mathscr{M}_I$ , so  $M \in L_{\mathbb{R}}(\mathscr{R} \cap \mathscr{M}_I)$ .

For the remainder of this section, I is a *two-sided* ideal.

The proof of the following result is straightforward.

PROPOSITION 2.4. With the notation of Proposition 2.3,  $\mathscr{R} \cap \mathscr{M}_I$  is a torsion class in Mod (R/I).

THEOREM 2.5. The correspondence  $\mathscr{T} \mapsto L_{\mathbb{R}}(\mathscr{T})$  defines a bijection from the set of torsion classes in Mod  $(\mathbb{R}/I)$  to the set of torsion subclasses of  $L_{\mathbb{R}}(\mathscr{M}_I)$  in Mod  $(\mathbb{R})$ .

*Proof.* By Propositions 2.3 and 2.4, the correspondence is surjective. If  $\mathscr{T}, \mathscr{U}$  are torsion classes in Mod (R/I) such that  $L_{\mathbb{R}}(\mathscr{T}) = L_{\mathbb{R}}(\mathscr{U})$ , then every nonzero R/I-module in  $\mathscr{U} \subseteq L_{\mathbb{R}}(\mathscr{T})$  has a nonzero R-(and thus R/I-) submodule in  $\mathscr{T}$ , whence  $\mathscr{U} \subseteq \mathscr{T}$ , since  $\mathscr{U}$  is homomorphically closed. Injectivity of the correspondence follows.

Using Proposition 2.2, we obtain.

COROLLARY 2.6. If I is right T-nilpotent, the correspondence  $\mathscr{T} \mapsto L_{\mathbb{R}}(\mathscr{T})$  defines a bijection from the torsion classes in Mod  $(\mathbb{R}/I)$  to those in Mod  $(\mathbb{R})$ .

3. Lifting some torsion properties. We now apply the results of the previous section to obtain some connections between the kinds of torsion classes a ring R and a factor ring R/I can have when I is right T-nilpotent.

Suppose we have canonically associated with each ring R a class  $\mathscr{X}_{R}$  of R-modules. Let

$$X = X(R) = \bigcap \{ (0: M) \mid M \in \mathscr{X}_R \} .$$

Suppose further that  $\mathscr{X}_{R/X} = \mathscr{X}_R$  for all R.

THEOREM 3.1. Every nonzero R-module has a nonzero submodule in  $\mathscr{X}_{\mathbb{R}}$  if and only if X is right T-nilpotent and every nonzero R/X-module has a nonzero submodule in  $\mathscr{X}_{\mathbb{R}}$ .

*Proof.* If X is right T-nilpotent, then by Proposition 2.2, every nonzero R-module has a nonzero submodule in  $\mathcal{M}_X = \text{Mod}(R/X)$ . If in addition every nonzero R/X-module has a nonzero submodule in  $\mathcal{R}_R$ , the same must therefore be true of all nonzero R-modules. If, on the other hand, nonzero R-modules all have nonzero submodules in  $\mathcal{R}_R$ , then

$$\operatorname{Mod}(R) = L_{\mathbb{R}}(\mathscr{X}_{\mathbb{R}}) \subseteq L_{\mathbb{R}}(\mathscr{M}_{\mathbb{X}})$$
,

so by Proposition 2.2, X is right T-nilpotent. Also, if  $0 \neq M \in \mathcal{M}_X = Mod(R/X)$ , then M has a nonzero submodule in  $\mathcal{H}_R$ .

An example of a class satisfying the requirements of Theorem 3.1 is the class of simple R-modules, together with 0. A ring R is

448

(left) semi-artinian if every nonzero R-module has a simple submodule. The following result was obtained in a different manner by Năstăsescu and Popescu [21].

COROLLARY 3.2. A ring R is semi-artinian if and only if its Jacobson radical J is right T-nilpotent and R/J is semi-artinian.

We call a nonzero module homogeneous (the terminology varies) if it is a rational extension of every nonzero submodule. (For the relevant details concerning rational extensions, see [23].) Hudry [9], [10], [11] calls a ring R (left) locally homogeneous (briefly, L. H.) if every nonzero R-module has a homogeneous submodule.

For any ring R, let

 $H = H(R) = \bigcap \{(0: M) | M \text{ is a homogeneous } R \text{-module} \}.$ 

COROLLARY 3.3. A ring R is L. H. if and only if H is right T-nilpotent and R/H is L.H.

Examples of L.H. rings are left noetherian ([9], [10], [11]) and semi-artinian rings.

Nonzero submodules of homogeneous modules are homogeneous (see, for example, [23], p. 625) and thus a ring R is L.H. if and only if every nonzero R-module has a cyclic homogeneous submodule. When R is commutative, a cyclic R-module R/P is homogeneous precisely when P is prime (see, for example [23], pp. 625-626). Thus for commutative rings R, H(R) is the prime radical B(R) and a commutative L.H. ring is one for which every nonzero module has an associated prime. The latter have been studied (under the name D-rings) by by Nguyen-Trong-Kham [17], [18].

COROLLARY 3.4. A commutative ring R is a D-ring if and only if B(R) is T-nilpotent and R/B(R) is a D-ring.

A nonzero module is co-irreducible if it is an essential extension of every nonzero submodule. (Such modules are also called *uniform*.) Năstăsescu and Popescu [20], [21] call a ring R (*left*) locally co-irreducible (briefly, L.C.) if every nonzero R-module has a co-irreducible submodule.

COROLLARY 3.5. A ring R is L.C. if and only if

 $C = C(R) = \bigcap \{(0: M) | M \text{ is a co-irreducible } R \text{-module} \}$ 

is right T-nilpotent and R/C is L.C.

Let  $\mathscr{X}_R$  be a nonempty class of *R*-modules and  $\widetilde{\mathscr{X}}_R$  be the class of *R*-modules, every nonzero homomorphic image of which has a nonzero submodule in  $\mathscr{X}_R$ . For the purposes of further applications of Theorem 2.5, it may be worthwhile to note that if  $\mathscr{X}_R$  is not homomorphically closed, one can have  $\widetilde{\mathscr{X}}_R \subseteq L_R(\mathscr{X}_R)$ . For instance, let *R* be the ring *Z* of integers and  $\mathscr{X}_Z$  the class of unbounded *p*groups, together with 0. Then  $\widetilde{\mathscr{X}}_Z$  is the class of divisible *p*-groups, while  $L_Z(\mathscr{X}_Z)$  is the class of all *p*-groups. When  $\mathscr{X}_R$  is the class of homogeneous modules, Hudry [12], [13] has shown that  $\widetilde{\mathscr{X}}_R = L_R(\mathscr{X}_R)$ for all rings *R*. A parallel argument establishes the corresponding equality for the co-irreducible modules.

We next consider the lifting of a property possessed by the ring of integers.

THEOREM 3.6. Let R be a ring with a right T-nilpotent ideal I. If every nontrivial torsion class in Mod (R/I) is determined by simple modules, the same is true in Mod (R).

*Proof.* The bijection of Corollary 2.6 clearly takes Mod (R/I) to Mod (R) and  $\{0\}$  to  $\{0\}$ . Thus if  $\mathscr{R}$  is a nontrivial torsion class in Mod (R), then  $\mathscr{R} = L_R(L_{R/I}(\mathscr{C}))$  for a set  $\mathscr{C}$  of simple modules. It follows that  $\mathscr{R} = L_R(\mathscr{C})$ .

The last result is eminently generalizable.

If a ring has the form  $A \oplus B$ , where A and B are ideals, the class of modules M such that AM = 0 is a torsion class. (The corresponding radical theory is said to be *centrally splitting* [3].)

THEOREM 3.7. Let R be a ring a with right T-nilpotent ideal I such that  $R/I = A \bigoplus B$  (ring direct sum). Then

$$L_{\scriptscriptstyle R}(\{M\in {
m Mod}\ (R)\,|\, IM=0 \ {
m and} \ AM=0\}$$

is a TTF class.

*Proof.* The identity of A lifts to an idempotent  $e \in R$ , since I is nil. Since ReR is an idempotent ideal,

$$\mathscr{R} = \{ M \in \text{Mod} (R) | ReRM = 0 \}$$
$$= \{ M \in \text{Mod} (R) | eM = 0 \}$$

is a TTF class [14]. Let

$$\mathscr{T} = L_{\scriptscriptstyle R}(\{M \in {\rm Mod}\ (R) \,|\, IM = 0 \ {\rm and} \ AM = 0\})$$
.

We show that  $\mathscr{R} = \mathscr{T}$ .

If  $N \in \mathscr{T} \subseteq L_{\mathbb{R}}(\mathscr{M}_{I})$ , it has an ascending *I*-series

 $0 = N_{\scriptscriptstyle 0} \subseteq N_{\scriptscriptstyle 1} \subseteq \cdots \subseteq N_{\scriptscriptstyle lpha} \subseteq N_{\scriptscriptstyle lpha+1} \subseteq \cdots \subseteq N_{\scriptscriptstyle \mu} = N$  ,

each term of which belongs to T. It follows from Corollary 2.6 that

$$\mathcal{T} \cap \mathcal{M}_I = \{M \in \operatorname{Mod}(R) \mid IM = 0 \text{ and } AM = 0\}.$$

Thus  $AN_1 = 0$ , so  $eN_1 \subseteq AN_1 = 0$  and  $N_1 \in \mathscr{R}$ . Each  $N_{\alpha+1}/N_{\alpha} \in \mathscr{T}$ , so in the same way,  $e(N_{\alpha+1}/N_{\alpha}) = 0$ , i.e.  $N_{\alpha+1}/N_{\alpha} \in \mathscr{R}$ . Clearly also  $N_{\beta} \in \mathscr{R}$  for a limit  $\beta$  if  $N_{\alpha} \in \mathscr{R}$  for every  $\alpha < \beta$ . Hence  $\mathscr{R}$  contains all  $N_{\alpha}$ , and in particular,  $N \in \mathscr{R}$ . This proves that  $\mathscr{T} \subseteq \mathscr{R}$ .

On the other hand, if  $K \in \mathscr{R}$  and IK = 0, then AK = 0, so that

$$\mathscr{R} = L_{\scriptscriptstyle R}(\{M \in \mathscr{R} \, | \, IM = 0\}) \ \subseteq L_{\scriptscriptstyle R}(\{M \in \operatorname{Mod}\,(R) \, | \, IM = 0 \, \, ext{and} \, \, AM = 0\} = \mathscr{T}.$$

As a consequence, we have the following frequently-proved result [1], [6], [22].

COROLLARY 3.8. If R is right perfect, it has finitely many torsion classes, all of which are TTF classes and all but  $\{0\}$  of which are determined by simple modules.

*Proof.* The Jacobson radical J of R is right T-nilpotent and any torsion class in Mod (R/J) has the form  $\{M \in \text{Mod } (R/J) | AM = 0\}$ for a ring direct summand A of R/J. By Corollary 2.6, the torsion classes in Mod (R) have the form

$$L_{\mathbb{R}}(\{M \in \operatorname{Mod}(\mathbb{R}) | JM = 0 \text{ and } AM = 0\})$$

and so, by Theorem 3.7, are TTF classes. The rest follows from Corollary 2.6 by an argument like that used for Theorem 3.6.

4. A splitting theorem. Throughout this section  $G_t$  and  $G_a$  denote, respectively, the maximum torsion and divisible subgroups of an abelian group G. Things such as torsion modules and divisible rings are those modules and rings whose additive groups have the stated properties.

THEOREM 4.1. Let R be a ring with a right T-nilpotent ideal I such that R/I is the direct sum of a torsion (necessarily bounded) ideal and a torsion-free divisible ideal. Then R has the same property.

*Proof.* Let  $A = (R/I)_t$  and  $B = (R/I)_d$ . Our assumption implies

that  $R/I = A \bigoplus B$  (ring direct sum). Thus A, B have identities  $e_A$ ,  $e_B$  respectively, and any R/I-module M has the form  $M_A \bigoplus M_B$  where is an A-module with  $BM_A = 0$  and  $M_B$  is a B-module with  $AM_B = 0$ . In particular,  $M_A$  is bounded and  $M_B$  is divisible. If  $x \in M_B$  and nx = 0 for some n > 0, then  $x = e_B x = n(e_B/n)x = (e_B/n)nx = 0$ , so  $M_B$  is torsion-free.

 $\operatorname{Let}$ 

$$\mathscr{R}_t = L_R(\{M \in \operatorname{Mod}(R) | IM = 0 \text{ and } BM = 0\}$$

If N is a torsion R-module, then every nonzero homomorphic image of N has a nonzero submodule  $M \in \mathscr{M}_I$  (Proposition 2.2). But M is a torsion R/I-module, so BM = 0. It follows that  $N \in \mathscr{R}_t$  and thus  $\mathscr{R}_t$  is the class of all torsion R-modules. By Theorem 3.7,  $\mathscr{R}_t$  is a TTF class so  $L = \prod_{i=1}^{\infty} N_i \in \mathscr{R}_t$ , where each  $N_i \cong N$ . But  $L_t$ , as a fully invariant subgroup, is a submodule, so  $\hat{L} = L/L_t \in \mathscr{R}_t$  and, if nonzero, has a nonzero submodule T such that IT = 0 and BT = 0 (so that, in particular, T is a torsion module). But  $\hat{L}$  is torsion-free, so  $\hat{L} = 0$ ; but then N must be bounded. Thus all torsion R-modules are bounded.

Let K be a nonzero torsion-free R-module. Since I is right Tnilpotent, K has a nonzero submodule P such that IP = 0 (Proposition 2.2). Clearly AP = 0, so P = BP is divisible. Thus  $K_d \neq 0$ . But  $K/K_d$  is again a torsion-free R-module, and so, if nonzero, must, by the same argument, have a nonzero divisible submodule. Hence  $K = K_d$ .

Thus for any *R*-module *V*,  $V_t$  is bounded and hence a direct summand, while  $V/V_t$  is divisible. It follows that  $V = V_t \bigoplus V_d$ . In particular,  $R = R_t \bigoplus R_d$  (module direct sum). But  $R_t$  and  $R_d$  are ideals, so the theorem is proved.

We conclude this section with a few remarks which give some indication of the range of applicability of the foregoing theorem. If a ring R has no nonzero nilpotent ideals then neither does  $R_t$ (whether it has an identity or not) so that  $R_t$  is bounded (elementary, in fact) [7], and hence a group direct summand. If in addition  $R/R_t$ is divisible, then  $R = R_t \bigoplus R_d$  (ring direct sum). Simple rings are either torsion-free divisible or elementary p-rings; torsion-free regular rings are divisible.

The author thanks the referee for suggestions which led to some shorter proofs and some improvements in the layout of this paper.

#### References

<sup>1.</sup> J. S. Alin and E. P. Armendariz, *TTF-classes over perfect rings*, J. Austral. Math. Soc., **11** (1970), 499-503.

2. H. Bass, Finitistic dimension and a homological generalization of semi-primary rings, Trans. Amer. Math. Soc., **95** (1960), 466-488.

3. R. L. Bernhardt, Splitting hereditary torsion theories over semiperfect rings, Proc. Amer. Math. Soc., **22** (1969), 681-687.

4. S. E. Dickson, A torsion theory for abelian categories, Trans. Amer. Math. Soc., 121 (1966), 223-235.

5. \_\_\_\_\_, Direct decompositions of radicals, pp. 366-374 of Proceedings of the Conference on Categorical Algebra, La Jolla, 1965. Springer-Verlag, Berlin-Heidelberg-New York, 1966.

6. V. Dlab, A characterization of perfect rings, Pacific J. Math., 33 (1970), 79-88.

7. L. Fuchs, Ringe und ihre additive Gruppe, Publ. Math. Debrecen, 4 (1956), 488-508.

8. B. J. Gardner, Some aspects of T-nilpotence, Pacific J. Math., 53 (1974), 117-130.

9. A. Hudry, Applications de la théorie de la localisation aux anneaux et aux modules, Thesis, Université de Lyon, 1970.

10. \_\_\_\_\_, Sur la localisation dans une catégorie de modules, C. R. Acad. Sci. Paris Sér. A-B **270** (1970), A 925-A 928.

11. \_\_\_\_, Sur les anneaux localement homogènes, C. R. Acad. Sci. Paris Sée. A-B **271** (1970), A1214-A1217.

12. \_\_\_\_\_, Une remarque sur la décomposition primaire de O. Goldman, Publ. Dép. Math. (Lyon) 8-1 (1971), 63-68.

13. \_\_\_\_\_, Sur les modules primaires de O. Goldman, C. R. Acad. Sci. Paris Sér. A-B **274** (1972), A1772-A1774.

14. J. P. Jans, Some aspects of torsion, Pacific J. Math., 15 (1965), 1249-1259.

15. A. I. Kashu, Radikaly v kategorii A-modulei, opredelennye s pomoshchyu elementov, Bul. Akad. Štiince RSS Moldoven, (1969), no. 1, 24-30.

16. J. M. Kellett, A torsion theory for modules over rings without identities, Thesis, University of Florida, 1968.

17. Nguyen-Trong-Kham, *D-anneaux*. I, Rev. Roumaine Math. Pures Appl., **17** (1972), 1671-1680.

18. \_\_\_\_\_, Théorie de la décomposition primaire dans les D-anneaux. II, Rev. Roumaine Math. Pures Appl., **18** (1973), 65-76.

19. J. Levitzki, Contributions to the theory of nilrings, (Hebrew, with English summary), Riveon Lematematika, 7 (1953), 50-70.

20. C. Năstăsescu and N. Popescu, Sur la structure des objets de certaines catégories abéliennes, C. R. Acad. Sci. Paris Sér. A-B **262** (1966), A1295-A1297.

21. \_\_\_\_, Anneaux semi-artiniens, Bull. Soc. Math. France, 96 (1968), 357-368.

22. E. A. Rutter, Jr., Torsion theories over semiperfect rings, Proc. Amer. Math. Soc., 34 (1972), 389-395.

23. H. H. Storrer, On Goldman's primary decomposition, pp. 621-661 of Lectures on Rings and Modules (Lecture Notes in Mathematics. Vol. 246) Springer-Verlag, Berlin-Heidelberg-New York, 1972.

Received March 19, 1975.

UNIVERSITY OF TASMANIA, HOBART, AUSTRALIA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor) University of California Los Angeles, California 90024 J. DUGUNDJI

Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

#### ASSOCIATE EDITORS

E.F. BECKENBACH

R. A. BEAUMONT

University of Washington

Seattle, Washington 98105

### B. H. NEUMANN

F. Wolf

K. Yoshida

### SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON \* \* \* AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The Pacific Journal of Mathematics is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

> Copyright © 1975 by Pacific Journal of Mathematics Manufactured and first issued in Japan

# Pacific Journal of Mathematics Vol. 59, No. 2 June, 1975

Aharon Atzmon, A moment problem for positive measures on the unit disc	317
Peter W. Bates and Grant Bernard Gustafson, Green's function inequalities for	
two-point boundary value problems	327
Howard Edwin Bell, Infinite subrings of infinite rings and near-rings	345
Grahame Bennett, Victor Wayne Goodman and Charles Michael Newman, Norms of	250
ranaom matrices $E^2$	339
Beveriy L. Brechner, Almost periodic homeomorphisms of E <sup>-</sup> are periodic	367
Beverly L. Brechner and R. Daniel Mauldin, <i>Homeomorphisms of the plane</i>	375
Jia-Arng Chao, Lusin area functions on local fields	383
Frank Rimi DeMeyer, The Brauer group of polynomial rings	391
M. V. Deshpande, Collectively compact sets and the ergodic theory of	•
semi-groups	399
Raymond Frank Dickman and Jack Ray Porter, $\theta$ -closed subsets of Hausdorff	407
spaces	407
Charles P. Downey, <i>Classification of singular integrals over a local field</i>	417
Daniel Reuven Farkas, <i>Miscellany on Bieberbach group algebras</i>	427
Peter A. Fowler, <i>Infimum and domination principles in vector lattices</i>	437
Barry J. Gardner, Some aspects of T-nilpotence. II: Lifting properties over	
<i>T</i> -nilpotent ideals	445
Gary Fred Gruenhage and Phillip Lee Zenor, Metrization of spaces with countable	
large basis dimension	455
J. L. Hickman, <i>Reducing series of ordinals</i>	461
Hugh M. Hilden, Generators for two groups related to the braid group	475
Tom (Roy Thomas Jr.) Jacob, Some matrix transformations on analytic sequence	
spaces	487
Elyahu Katz, <i>Free products in the category of</i> $k_w$ <i>-groups</i>	493
Tsang Hai Kuo, On conjugate Banach spaces with the Radon-Nikodým property	497
Norman Eugene Liden, <i>K</i> -spaces, their antispaces and related mappings	505
Clinton M. Petty, Radon partitions in real linear spaces	515
Alan Saleski, A conditional entropy for the space of pseudo-Menger maps	525
Michael Singer, Elementary solutions of differential equations	535
Eugene Spiegel and Allan Trojan, On semi-simple group algebras. I	549
Charles Madison Stanton, Bounded analytic functions on a class of open Riemann	557
Sherman K. Stain. Transversals of Latin squares and their generalizations	567
Juan Ernest Stuy. Distribution of squarefree integers in non-linear sequences	507
I owell C. Sweet. On homeoencous glochnas	505
Lowell G. Sweet, On homogeneous algebras	585
Lowell G. Sweet, On doubly homogeneous algebras	595
Florian Vasilescu, The closed range modulus of operators	599
Arthur Anthony Yanushka, A characterization of the symplectic groups $PSp(2m, q)$ as rank 3 permutation groups	611
James Juei-Chin Yeh, Inversion of conditional Wiener integrals	623