

# Pacific Journal of Mathematics

**FREE PRODUCTS IN THE CATEGORY OF  $k_w$ -GROUPS**

ELYAHU KATZ

# FREE PRODUCTS IN THE CATEGORY OF $k_w$ -GROUPS

ELYAHU KATZ

**This paper provides a short proof of the existence of free-products in the category  $k_w$ -groups.**

1. Introduction. Graev defined free products in the category of Hausdorff topological groups, and gave a lengthy proof of their existence using norms [2]. Hulanicki showed that free products exist in the category of compact groups [3]. S. A. Morris pointed out that the later result extends to the category of almost periodic groups [5]. The proof in the above mentioned paper is correct, except that it should rely on Graev's proof for the existence of free products in the category of Hausdorff topological groups and should not on the erroneous proof in that paper.

The method of proof of this paper has been adopted by Ordman to prove the existence of free products in the category of  $k$ -groups [6].

## 2. Definitions and statement of the main result.

DEFINITION 1. a  $k_w$ -space is a Hausdorff topological space  $X$  with compact subsets  $X_n$  such that: (i)  $X = \bigcup_{n=1}^{\infty} X_n$ ; (ii)  $X_{n+1} \supset X_n$  for all  $n$ ; (iii) a subset  $A$  of  $X$  is closed in  $X$ , if and only if  $A \cap X_n$  is compact for all  $n$ .

By a  $k_w$ -decomposition  $X = \bigcup X_n$ , we mean that  $X_n$  have properties (i), (ii) and (iii).

A  $k_w$ -group  $G$ , is a topological group which is also a  $k_w$ -space. We denote by  $KG$  the category of  $k_w$ -groups.

DEFINITION 2. Let  $G_1$  and  $G_2$  be two  $k_w$ -groups. Then,  $G_1 * G_2$  is their free product in the category of  $k_w$ -groups if the following axioms hold: (i) the underlying group of  $G_1 * G_2$  is their free product as groups. (ii)  $G_1$  and  $G_2$  are topological subgroups of  $G_1 * G_2$ . (iii) If  $\gamma_i: G_i \rightarrow H$ ,  $i = 1, 2$ , are continuous homomorphisms into the  $k_w$ -group  $H$ , then they extend uniquely to a continuous homomorphism  $\Gamma: G_1 * G_2 \rightarrow H$ .

We will refer to the following concept frequently.

DEFINITION 3 (Graev [1]). The Hausdorff topological group  $F(X)$  is a free topological group with basis  $(X, e)$  a pointed topological space, if the following axioms hold: (i)  $X$  is a subspace of  $F(X)$ .

(ii)  $X - e$  generates  $F(X)$  freely as a group. (iii) For any continuous map  $\phi: X \rightarrow G$ , where  $G$  is a Hausdorff topological group, and  $\phi(e)$  equals the identity of  $G$ , then  $\phi$  extends uniquely to a continuous homomorphism  $\Phi: F(X) \rightarrow G$ .

*The main result.* Free products exist in the category of  $k_w$ -groups.

**3. Proof of the main result.** Let  $\psi: |F(G_1 \vee G_2)| \rightarrow |G_1| * |G_2|$  be the homomorphism induced by the inclusion map of  $G_1 \vee G_2$  into  $|G_1| * |G_2|$ , where  $G_1, G_2 \in KG$ ,  $\vee$  denotes the disjoint union identifying distinguished points,  $| \cdot |$  is the forgetful functor into the category of groups and  $*$  denotes the free product in the category of groups, in addition to its use in Definition 2. Denote by  $K$  the kernel of  $\psi$  with the subspace topology of  $F(G_1 \vee G_2)$ . Our first object is to prove that  $K$  is a closed subset. For then,  $|G_1| * |G_2|$  with the identification topology of  $F(G_1 \vee G_2)$  via  $\psi$ , is a  $k_w$ -group, which we denote by  $G_1 *' G_2$ .

To carry out the proof of the just mentioned fact we need some further notation. The map  $h: (G_1 \vee G_1^{-1} \vee G_2 \vee G_2^{-1})^n \rightarrow F(G_1 \vee G_2)$  is defined to send words in  $G_1 \vee G_2$  whose length is at most  $n$ , to their reduced form. This map is obviously continuous, since it is induced by the multiplication map of  $F(G_1 \vee G_2)$ . We also use the following notation:  $Z^n = Y_1^n \times \dots \times Y_n^n$ , where  $Y_i^n = (L_{q_i}^n)^{p_i}$ ,  $p_i = \pm 1$ ,  $q_i = 1, 2$  and  $G_i = \bigcup_{j=1}^\infty L_i^j$  are  $k_w$  decompositions,  $i = 1, 2$ .

The sets,  $h[(L_1^n \vee (L_1^n)^{-1} \vee L_2^n \vee (L_2^n)^{-1})^n]$  which consist of all reduced words in  $L_1^n \vee L_2^n$ , whose length is at most  $n$ , with the subspace topology of  $F(G_1 \vee G_2)$ , form a  $k_w$ -decomposition of  $F(G_1 \vee G_2)$  [4]. Thus in order to show that  $K$  is closed, it suffices to show that  $K \cap h(Z^n)$  is compact for any  $n$ .

Suppose  $Z^n$  is of the form  $(L_{i_0}^n)^{l_1} \times \dots \times (L_{i_0}^n)^{l_n}$ ,  $i_0$  fixed, 1 or 2. Denote by  $\mu: \times G_{i_0} \rightarrow G_{i_0}$  the map induced by the multiplication in  $G_{i_0}$ , and denote by  $\phi: Z^n \rightarrow \times G_{i_0}$  the map induced by the inclusion of  $L_{i_0}^n$  into  $G_{i_0}$ . By the continuity of the maps  $h, \phi, \mu$ , the set  $K \cap h(Z^n) = h(\ker \mu\phi)$  is compact. If  $Z^n$  is not of the above form,  $K \cap h(Z^n)$  will be, by an induction argument, a product of compact sets.

Next we prove the  $G_1 *' G_2$  is a free product. Let  $t': F(G_1 \vee G_2) \rightarrow H$  be the extension of  $t_i: G_i \rightarrow H$ ,  $i = 1, 2$ , two continuous homomorphisms into the  $k_w$ -group  $H$ . The homomorphism  $\psi$  factors through  $t'$ , i.e., the following diagram commutes:

$$\begin{array}{ccc}
 F(G_1 \vee G_2) & \xrightarrow{\psi} & G_1 *' G_2 \\
 t' \downarrow & \swarrow t & \\
 H & & 
 \end{array}$$

Since  $\psi$  is an identification map,  $t$  is continuous. The uniqueness of  $t$  follows from the algebraic structure.

We still have to show that  $G_1 *' G_2$  induces the original topology on  $|G_i|$ . Let  $g: G_1 *' G_2$  be induced by  $l_{G_1}$  and the constant map on  $G_2$ . Since  $g^{-1}(U) \cap G_1 = U$ , if  $U$  is open in  $G_1$ ,  $U$  is also open in  $G_1'$ , where  $G_1'$  is  $|G_1|$  with the subspace topology of  $G_1 *' G_2$ . Suppose that  $V'$  is open in  $G_1 *' G_2$  and let  $V = V' \cap G_1'$ . Denote by  $\chi: F(G_1) \rightarrow F(G_1 \vee G_2)$  the map induced by the inclusion, and  $p: F(G_1) \rightarrow G_1$  the map which sends a word to its reduced form. Graev proved in [1] that  $p$  is an open map. Since  $V = p(\psi\chi)^{-1}(V')$ ,  $V$  is open in  $G_1$ . This proves that  $G_1 *' G_2$  induces the original topology on  $|G_1|$ . We conclude that  $G_1 *' G_2$  is the free product of  $G_1$  and  $G_2$  in  $KG$ .

4. **Remarks.** In our proof that  $G_1 *' G_2$  satisfies property (iii) of Definition 2 we used only the fact that  $H$  is a Hausdorff topological group. Thus  $G_1 *' G_2$  is also the free product of  $G_1$  and  $G_2$  in the category of Hausdorff topological groups.

We point out that the free product in  $KG$  is actually the co-product in  $KG$ . This follows from the uniqueness property of co-products.

#### REFERENCES

1. M. I. Graev, *Free topological groups*, Izv. Akad. Nauk SSSR. Ser. Mat., **12** (1948), 279-324, (Russian). English Transl., Amer. Math. Soc. Transl. no. 35, (1951).
2. ———, *On free products of topological groups*, (Russian), Izv. Akad. Nauk SSSR. Ser. Mat., **14** (1950), 343-354.
3. Hulanicki, *Isomorphic embeddings of free products of compact groups*, Colloq. Math., **16** (1967), 235-241.
4. J. Mack, S. A. Morris, E. T. Ordman, *Free topological groups and the projective dimension of a locally compact abelian group*, Proc. Amer. Math. Soc., (to appear).
5. S. A. Morris, *Free products of topological groups*, Bull. Austral. Math. Soc., **4** (1971), 17-29.
6. E. T. Ordman, *Free Product in the Category of  $k$ -Groups*, (to appear).

Received April 26, 1974 and in revised form May 29, 1975.

TEL-AVIV UNIVERSITY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

|   |     |
|---|-----|
| Aharon Atzmon, <i>A moment problem for positive measures on the unit disc</i> .....   | 317 |
| Peter W. Bates and Grant Bernard Gustafson, <i>Green's function inequalities for two-point boundary value problems</i> .....                  | 327 |
| Howard Edwin Bell, <i>Infinite subrings of infinite rings and near-rings</i> .....  | 345 |
| Grahame Bennett, Victor Wayne Goodman and Charles Michael Newman, <i>Norms of random matrices</i> .....                                       | 359 |
| Beverly L. Brechner, <i>Almost periodic homeomorphisms of <math>E^2</math> are periodic</i> .....   | 367 |
| Beverly L. Brechner and R. Daniel Mauldin, <i>Homeomorphisms of the plane</i> .....   | 375 |
| Jia-Arng Chao, <i>Lusin area functions on local fields</i> .....  | 383 |
| Frank Rimi DeMeyer, <i>The Brauer group of polynomial rings</i> .....   | 391 |
| M. V. Deshpande, <i>Collectively compact sets and the ergodic theory of semi-groups</i> .....   | 399 |
| Raymond Frank Dickman and Jack Ray Porter, <i><math>\theta</math>-closed subsets of Hausdorff spaces</i> .....                                | 407 |
| Charles P. Downey, <i>Classification of singular integrals over a local field</i> .....   | 417 |
| Daniel Reuven Farkas, <i>Miscellany on Bieberbach group algebras</i> .....  | 427 |
| Peter A. Fowler, <i>Infimum and domination principles in vector lattices</i> .....  | 437 |
| Barry J. Gardner, <i>Some aspects of T-nilpotence. II: Lifting properties over T-nilpotent ideals</i> .....                                   | 445 |
| Gary Fred Gruenhagen and Phillip Lee Zenor, <i>Metritzation of spaces with countable large basis dimension</i> .....                          | 455 |
| J. L. Hickman, <i>Reducing series of ordinals</i> .....   | 461 |
| Hugh M. Hilden, <i>Generators for two groups related to the braid group</i> .....   | 475 |
| Tom (Roy Thomas Jr.) Jacob, <i>Some matrix transformations on analytic sequence spaces</i> .....  | 487 |
| Elyahu Katz, <i>Free products in the category of <math>k_w</math>-groups</i> .....  | 493 |
| Tsang Hai Kuo, <i>On conjugate Banach spaces with the Radon-Nikodým property</i> .....  | 497 |
| Norman Eugene Liden, <i>K-spaces, their antispace and related mappings</i> .....  | 505 |
| Clinton M. Petty, <i>Radon partitions in real linear spaces</i> .....   | 515 |
| Alan Saleski, <i>A conditional entropy for the space of pseudo-Menger maps</i> .....  | 525 |
| Michael Singer, <i>Elementary solutions of differential equations</i> .....   | 535 |
| Eugene Spiegel and Allan Trojan, <i>On semi-simple group algebras. I</i> .....  | 549 |
| Charles Madison Stanton, <i>Bounded analytic functions on a class of open Riemann surfaces</i> .....  | 557 |
| Sherman K. Stein, <i>Transversals of Latin squares and their generalizations</i> .....  | 567 |
| Ivan Ernest Stux, <i>Distribution of squarefree integers in non-linear sequences</i> .....  | 577 |
| Lowell G. Sweet, <i>On homogeneous algebras</i> .....   | 585 |
| Lowell G. Sweet, <i>On doubly homogeneous algebras</i> .....  | 595 |
| Florian Vasilescu, <i>The closed range modulus of operators</i> .....   | 599 |
| Arthur Anthony Yanushka, <i>A characterization of the symplectic groups <math>P\text{Sp}(2m, q)</math> as rank 3 permutation groups</i> ..... | 611 |
| James Juei-Chin Yeh, <i>Inversion of conditional Wiener integrals</i> .....   | 623 |