

# Pacific Journal of Mathematics

**INVERSION OF CONDITIONAL WIENER INTEGRALS**

JAMES JUEI-CHIN YEH

# INVERSION OF CONDITIONAL WIENER INTEGRALS

*Dedicated to Professor Robert H. Cameron*

J. YEH

**Given two Wiener measurable functionals  $X$  and  $Y$  on the Wiener space  $C[0, t]$ , of which the latter is Wiener integrable, the conditional Wiener integral of  $Y$  given  $X$  is defined as the conditional expectation  $E^w(Y|X)$  given as a function on the value space of  $X$ . Several Fourier inversion formulae for retrieving the conditional Wiener integral  $E^w(Y|X)$  in which  $X[x] = x(t)$  for  $x \in C[0, t]$  are derived. Examples of evaluation of  $E^w(Y|X)$  are given. It is shown that the Kac-Feynman formula can be derived by applying an inversion formula to  $E^w(Y|X)$  where**

$$Y[x] = \exp \left\{ - \int_0^t V[x(s)] ds \right\} .$$

1. Introduction. We have recently derived an inversion formula for conditional expectations. (See [4].) In the present paper we report on some inversion formulae for conditional Wiener integrals. Here the probability space is the Wiener measure space on the Wiener space  $C[0, t]$  of the real valued continuous functions  $x$  on  $[0, t]$  with  $x(0) = 0$  for fixed  $t \in (0, \infty)$ . By a conditional Wiener integral we mean the conditional expectation  $E^w(Y|X)$  of a real or complex valued Wiener integrable functional  $Y$  conditioned by a Wiener measurable functional  $X$  on  $C[0, t]$  which is given as a function on the value space of  $X$ . We shall be concerned exclusively with  $X$  given by  $X[x] = x(t)$  for  $x \in C[0, t]$ . Thus  $E^w(Y|X)$  will be a real or complex valued function on  $R^1$ . A precise definition of conditional Wiener integral as well as a brief discussion of the Wiener measure space are given in §2. Three inversion formulae for conditional Wiener integrals are proved in §3: a (C. 1) summability type inversion formula (Theorem 1), a Lévy type inversion formula (Theorem 2) and an inversion formula under the assumption of the Lebesgue integrability of  $E^w[e^{t u X} Y]$ ,  $u \in R^1$  (Theorem 3). Examples of evaluation of conditional Wiener integrals are given in §4. Below we relate conditional Wiener integrals to the Kac-Feynman formula.

Consider the Wiener integrable functional

$$\exp \left\{ - \int_0^t V[x(s)] ds \right\} \quad \text{for } x \in C[0, t]$$

where  $V$  is a nonnegative continuous function on  $R^1$ . Under the

additional condition that  $V$  is bounded, M. Kac [2], showed that a real valued function  $U$  on  $R^1 \times (0, \infty)$  defined by

$$U(\xi, t) = \sum_{k=0}^{\infty} (-1)^k U_k(\xi, t) \quad \text{for } (\xi, t) \in R^1 \times (0, \infty)$$

where the sequence  $\{U_k, k = 0, 1, 2, \dots\}$  is defined inductively by

$$U_0(\xi, t) = \frac{1}{\sqrt{2\pi t}} \exp \left\{ -\frac{\xi^2}{2t} \right\}$$

$$U_{k+1}(\xi, t) = \frac{1}{\sqrt{2\pi}} \int_0^t \left\{ \int_{-\infty}^{\infty} \frac{1}{\sqrt{t-s}} \exp \left\{ -\frac{1}{2} \frac{(\xi - \eta)^2}{t-s} \right\} V(\eta) U_k(\eta, s) d\eta \right\} ds$$

for  $k = 0, 1, 2, \dots$

satisfies the integral equation

$$(1.1) \quad U(\xi, t) = \frac{1}{\sqrt{2\pi t}} \exp \left\{ -\frac{\xi^2}{2t} \right\} - \frac{1}{\sqrt{2\pi}} \int_0^t \times \left\{ \int_{-\infty}^{\infty} \frac{1}{\sqrt{t-s}} \exp \left\{ -\frac{1}{2} \frac{(\xi - \eta)^2}{t-s} \right\} V(\eta) U(\eta, s) d\eta \right\} ds$$

for  $(\xi, t) \in R^1 \times (0, \infty)$  and the boundary condition

$$(1.2) \quad \int_a^b U(\xi, t) d\xi = E^w \left[ \exp \left\{ -\int_0^t V[x(s)] ds \right\}; a < x(t) < b \right]$$

for any  $(a, b) \subset R^1$ . The boundedness condition on  $V$  was then removed by the method of truncation. The integral equation (1.1) implies that  $U$  satisfies the differential equation

$$(1.3) \quad \frac{\partial U}{\partial t} = \frac{1}{2} \frac{\partial^2 U}{\partial \xi^2} - V(\xi) U \quad \text{for } (\xi, t) \in R^1 \times (0, \infty)$$

and the boundary condition (1.2) implies that  $U$  satisfies the initial condition

$$(1.4) \quad \lim_{t \rightarrow 0} \int_{-\varepsilon}^{\varepsilon} U(\xi, t) d\xi = 1 \quad \text{for every } \varepsilon > 0.$$

This result is summarized by the Kac-Feynman formula

$$(1.5) \quad U(\xi, t) = E^w \left[ \exp \left\{ -\int_0^t V[x(s)] ds \right\} \delta(x(t) - \xi) \right]$$

for the solution  $U$  of the differential system (1.3) and (1.4). Now let  $X_t$  and  $Y_t$  be two real valued functionals on  $C[0, t]$  defined by

$$X_t[x] = x(t) \quad \text{and} \quad Y_t[x] = \exp \left\{ -\int_0^t V[x(s)] ds \right\} \quad \text{for } x \in C[0, t].$$

The fact that the function  $U$  satisfies (1.2) indicates that it is related to the conditional Wiener integral  $E^w(Y_t|X_t)$ . Let us define a real valued function  $U$  on  $R^1 \times (0, \infty)$  by

$$(1.6) \quad U(\xi, t) = E^w(Y_t|X_t)(\xi) \frac{1}{\sqrt{2\pi t}} \exp \left\{ -\frac{\xi^2}{2t} \right\} \\ \text{for } (\xi, t) \in R^1 \times (0, \infty).$$

In §5 we show that by applying Theorem 1 to an appropriately chosen version of  $E^w(Y_t|X_t)$  we obtain the integral equation (1.1) for the function  $U$  defined by (1.6). The fact our function  $U$  satisfies (1.2) is obvious. This gives an alternate way of deriving the Kac-Feynman formula.<sup>1</sup>

2. **Conditional Wiener integral.** For a fixed  $t \in (0, \infty)$  consider the Wiener measure space  $(C[0, t], \mathfrak{B}^*, m_w)$  where  $\mathfrak{B}$  is the algebra of subsets  $W$  of  $C[0, t]$  of the type

$$(2.1) \quad W = \{x \in C[0, t]; [x(s_1), \dots, x(s_n)] \in B\}$$

where  $n$  is an arbitrary positive integer,  $0 = s_0 < s_1 < \dots < s_n \leq t$ , and  $B$  is an arbitrary member of the  $\sigma$ -algebra  $\mathfrak{B}^n$  of the Borel sets in the  $n$ -dimensional Euclidean space  $R^n$ ;  $m_w$  is a probability measure on the algebra  $\mathfrak{B}$  defined for  $W$  as in (2.1) by

$$(2.2) \quad m_w(W) = \left\{ (2\pi)^n \prod_{j=1}^n (s_j - s_{j-1}) \right\}^{-1/2} \\ \int_B \exp \left\{ -\frac{1}{2} \sum_{j=1}^n \frac{(\xi_j - \xi_{j-1})^2}{s_j - s_{j-1}} \right\} m_L(d\xi)$$

where  $\xi = (\xi_1, \dots, \xi_n) \in R^n$ ,  $\xi_0 = 0$  and  $m_L$  is the Lebesgue measure on  $(R^n, \mathfrak{B}^n)$ ;  $\mathfrak{B}^*$  is the  $\sigma$ -algebra of Carathéodory measurable subsets of  $C[0, t]$  with respect to the outer measure derived from the probability measure  $m_w$  on the algebra  $\mathfrak{B}$  which in particular contains the  $\sigma$ -algebra  $\sigma(\mathfrak{B})$  generated by  $\mathfrak{B}$ .

A real valued functional  $F$  on  $C[0, t]$  is said to be Wiener measurable if it is  $\mathfrak{B}^*$ -measurable, i.e. if it is a measurable transformation of  $(C[0, t], \mathfrak{B}^*)$  into  $(R^1, \mathfrak{B}^1)$ . For a Wiener measurable functional  $F$  we write

$$E^w[F] \quad \text{for} \quad \int_{C[0, t]} F[x] m_w(dx)$$

---

<sup>1</sup> The idea of deriving the Kac-Feynman formula by inversion goes back to M. D. Donsker. I wish to thank Professor Donsker for the conversations we had on this approach.

whenever the integral, i.e. the Wiener integral, exists. We say that  $F$  is Wiener integrable or  $m_w$ -integrable when the Wiener integral of  $F$  exists and is finite. The Wiener measurability and Wiener integrability of a complex valued functional on  $C[0, t]$  are defined in terms of its real and imaginary parts. An immediate consequence of the definition of the Wiener measure is that if  $F$  is a real or complex valued functional on  $C[0, t]$  of the type

$$(2.3) \quad F[x] = f[x(s_1), \dots, x(s_n)] \quad \text{for } x \in C[0, t]$$

where  $f$  is a real or complex valued Baire function on  $R^n$  and  $0 < s_1 < \dots < s_n \leq t$  then  $F$  is Wiener measurable and

$$(2.4) \quad E^w[F] = \left\{ (2\pi)^n \prod_{j=1}^n (s_j - s_{j-1}) \right\}^{-1/2} \int_{R^n} f(\xi) \exp \left\{ -\frac{1}{2} \sum_{j=1}^n \frac{(\xi_j - \xi_{j-1})^2}{s_j - s_{j-1}} \right\} m_L(d\xi)$$

in the sense that the existence of one side implies that of the other as well as the equality of the two.

In connection with the Wiener measure space let us remark that a real valued function  $X$  on  $[0, t] \times C[0, t]$  defined by

$$X(s, x) = x(s) \quad \text{for } (s, x) \in [0, t] \times C[0, t]$$

is a Brownian motion process on the probability space  $(C[0, t], \mathfrak{B}^*, m_w)$  and the domain  $[0, t]$  in which the space of sample functions  $X(\cdot, x)$ ,  $x \in C[0, t]$ , coincides with the sample space  $C[0, t]$  and thus every sample function is continuous. This last property implies in particular that the process is a measurable process. We shall refer to this realization of the Brownian motion process as the Wiener process on the domain  $[0, t]$ .

DEFINITION. Let  $X$  and  $Z$  be real valued Wiener measurable functionals on the Wiener measure space  $(C[0, t], \mathfrak{B}^*, m_w)$  and let  $Z$  be Wiener integrable on  $C[0, t]$ . Let  $P_X$  be the probability distribution of  $X$ , i.e. the probability measure on  $(R^1, \mathfrak{B}^1)$  determined by  $X$  by the definition

$$P_X(B) = m_w(X^{-1}(B)) \quad \text{for } B \in \mathfrak{B}^1.$$

The conditional Wiener integral of  $Z$  given  $X$ , written  $E^w(Z|X)$ , is by definition the equivalence class of  $\mathfrak{B}^1$ -measurable and  $P_X$ -integrable functions  $\psi$  on  $R^1$  modulo null functions on  $(R^1, \mathfrak{B}^1, P_X)$  such that

$$\int_{X^{-1}(B)} Z[x]m_w(dx) = \int_B \psi(\xi)P_X(d\xi) \quad \text{for } B \in \mathfrak{B}^1.$$

By the Radon-Nikodym theorem such a function  $\psi$  exists and is determined up to a null function on  $(R^1, \mathfrak{B}^1, P_X)$ . We shall also use  $E^w(Z|X)$  to mean a particular version, i.e. a particular representative of the equivalence class. Thus

$$(2.5) \quad \int_{X^{-1}(B)} Z[x]m_w(dx) = \int_B E^w(Z|X)(\xi)P_X(d\xi) \quad \text{for } B \in \mathfrak{B}^1.$$

### 3. Inversion formulae of conditional Wiener integrals.

LEMMA 1. *Let  $X$  and  $Z$  be measurable transformations of  $(C[0, t], \mathfrak{B}^*)$  into  $(R^1, \mathfrak{B}^1)$  with  $E^w[|Z|] < \infty$ . Let  $g$  be a measurable transformation of  $(R^1, \mathfrak{B}^1)$  into itself. Then*

$$E^w[(g \circ X)Z] = \int_{R^1} g(\xi)E^w(Z|X)(\xi)P_X(d\xi)$$

*in the sense that the existence of one side implies that of the other as well as the equality of the two.*

*Proof.* This lemma is a particular case of Proposition 3 in [4].

COROLLARY. *Let  $X$  and  $Z$  be as in Lemma 1. Assume that  $P_X \ll m_L$  on  $(R^1, \mathfrak{B}^1)$ . For  $\xi \in R^1$  and  $a > 0$  let  $\chi_{\xi, a}$  be a function on  $R^1$  defined by*

$$\chi_{\xi, a}(\eta) = \begin{cases} \frac{1}{2a} & \text{for } \eta \in [\xi - a, \xi + a] \\ 0 & \text{for } \eta \in [\xi - a, \xi + a]^c. \end{cases}$$

*Then there exists a version of  $E^w(Z|X)(dP_X/dm_L)$  such that*

$$E^w(Z|X)(\xi) \frac{dP_X}{dm_L}(\xi) = \lim_{a \rightarrow 0} E^w[(\chi_{\xi, a} \circ X)Z] \quad \text{for } \xi \in R^1.$$

*Proof.* Using  $\chi_{\xi, a}$  in the place of  $g$  in Lemma 1, we have

$$\begin{aligned} & \lim_{a \rightarrow 0} E^w[(\chi_{\xi, a} \circ X)Z] \\ &= \lim_{a \rightarrow 0} \int_{R^1} \chi_{\xi, a}(\eta) E^w(Z|X)(\eta) \frac{dP_X}{dm_L}(\eta) m_L(d\eta). \end{aligned}$$

It is well known that if  $f \in L^1_{loc}(R^1, \mathfrak{B}^1, m_L)$  then

$$\lim_{a \rightarrow 0} \int_{R^1} \chi_{\xi, a}(\eta) f(\eta) m_L(d\eta) = f(\xi) \quad \text{for } m_L\text{-a.e. } \xi \in R^1.$$

From this follows our corollary.

LEMMA 2. *Let  $X$  and  $Y$  be measurable transformations of  $(C[0, t], \mathfrak{B}^*)$  into  $(R^1, \mathfrak{B}^1)$  with  $E^w[|Y|] < \infty$ . Then*

$$E^w[e^{iuX} Y] = \int_{R^1} e^{iu\xi} E^w(Y|X)(\xi) P_X(d\xi) \text{ for } u \in R^1.$$

*Proof.* This lemma is the equality (3.19) in [4] adapted to our probability space  $(C[0, t], \mathfrak{B}^* m_w)$ .

THEOREM 1. *Let  $X$  and  $Y$  be measurable transformations of  $(C[0, t], \mathfrak{B}^*)$  into  $(R^1, \mathfrak{B}^1)$  with  $E^w[|Y|] < \infty$ . Assume that  $P_X \ll m_L$  on  $(R^1, \mathfrak{B}^1)$ . Then there exists a version of  $E^w(Y|X)(dP_X/dm_L)$  such that*

$$\begin{aligned} & E^w(Y|X)(\xi) \frac{dP_X}{dm_L}(\xi) \\ &= \lim_{a \rightarrow \infty} \frac{1}{2\pi} \int_{(-a, a)} \left(1 - \frac{|u|}{a}\right) e^{-iu\xi} E^w[e^{iuX} Y] m_L(du) \text{ for } \xi \in R^1. \end{aligned}$$

*Proof.* By Lemma 2

$$(3.1) \quad E^w[e^{iuX} Y] = \int_{R^1} e^{iu\xi} E^w(Y|X)(\xi) \frac{dP_X}{dm_L}(\xi) m_L(d\xi) \text{ for } u \in R^1.$$

Thus our  $E^w[e^{iuX} Y]$ ,  $u \in R^1$ , is the Fourier transform of the  $m_L$ -integrable function  $E^w(Y|X)(\xi) (dP_X/dm_L)(\xi)$ ,  $\xi \in R^1$ . According to a well known (C, 1) summability type inversion formula for Fourier transforms, if  $f$  is a  $m_L$ -integrable function on  $R^1$  and  $\hat{f}$  is its Fourier transform then

$$f(\xi) = \lim_{a \rightarrow \infty} \frac{1}{2\pi} \int_{(-a, a)} \left(1 - \frac{|u|}{a}\right) e^{-iu\xi} \hat{f}(u) m_L(du) \text{ for } m_L\text{-a.e. } \xi \in R^1$$

(see for instance [1]). From this follows our theorem.

THEOREM 2. *Let  $X$  and  $Y$  be as in Theorem 1. For  $a, b \in R^1$  such that  $a < b$  let  $\tilde{\chi}_{a,b}$  be a function on  $R^1$  defined by*

$$\tilde{\chi}_{a,b}(\eta) = \begin{cases} 1 & \text{for } \eta \in (a, b) \\ 0 & \text{for } \eta \in [a, b]^c \\ \frac{1}{2} & \text{for } \eta = a \text{ and for } \eta = b. \end{cases}$$

Then

$$\int_{R^1} \tilde{\chi}_{a,b}(\xi) E^w(Y|X)(\xi) \frac{dP_X(\xi)}{dm_L} m_L(d\xi) = \lim_{h \rightarrow \infty} \frac{1}{2\pi} \int_{(-h,h)} \frac{e^{-ib u} - e^{-ia u}}{-iu} E^w[e^{i u X} Y] m_L(du) .$$

*Proof.* As an immediate consequence of the Lévy inversion formula, if  $\Phi$  is a finite signed measure on  $(R^1, \mathfrak{B}^1)$  and if  $\varphi$  is the characteristic function  $\Phi$ , i.e.

$$\varphi(u) = \int_{R^1} e^{i u \xi} \Phi(d\xi) \quad \text{for } u \in R_1,$$

then we have

$$\int_{R^1} \tilde{\chi}_{a,b}(\xi) \Phi(d\xi) = \lim_{h \rightarrow \infty} \frac{1}{2\pi} \int_{(-h,h)} \frac{e^{-ib u} - e^{-ia u}}{-iu} \varphi(u) m_L(du) .$$

From this and from (3.1) follows our theorem.

**THEOREM 3.** *Let  $X$  and  $Y$  be as in Theorem 1. Assume further that  $E^w[e^{i u X} Y]$  is a  $m_L$ -integrable function of  $u$  on  $R^1$ . Then a version of  $E^w(Y|X)(dP_X/dm_L)$  is given by*

$$E^w(Y|X)(\xi) \frac{dP_X(\xi)}{dm_L} = \frac{1}{2\pi} \int_{R^1} e^{-i u \xi} E^w[e^{i u X} Y] m_L(du) \quad \text{for } \xi \in R^1 .$$

*Proof.* This theorem is a particular case of Theorem 2 in [4].

#### 4. Examples of evaluation of conditional Wiener integrals.

**EXAMPLE 1.<sup>2</sup>** For each  $x \in C[0, t]$  consider the average value of  $x$  over the time interval  $[0, t]$ , i.e.  $1/t \int_0^t x(s) ds$ . Let us find the conditional expectation of this average value given the condition that the final value of  $x, x(t)$ , is equal to  $\xi \in R^1$ . Thus we are to find  $E^w(Z|X)(\xi), \xi \in R^1$ , for

$$Z[x] = \frac{1}{t} \int_0^t x(s) ds \quad \text{and} \quad X[x] = x(t) \quad \text{for } x \in C[0, t] .$$

According to Corollary to Lemma 1, a version of  $E^w(Z|X)(dP_X/dm_L)$  is given by

$$(4.1) \quad E^w(Z|X)(\xi) \frac{dP_X(\xi)}{dm_L} = \lim_{a \rightarrow 0} E^w[(\chi_{\xi, \cdot, a} X) Z] \quad \text{for } \xi \in R^1 .$$

---

<sup>2</sup> According to Professor E. O. Thorp, this problem arises in the probabilistic study of the fluctuation of the price of a commodity as a Brownian motion process. The problem was to find the expectation of the time average of the price when the price at some fixed time point in the future is known for some reason.



With our  $Z$  and  $X$  we have

$$(4.2) \quad E^w[(\chi_{\xi, a} \circ X)Z] = E^w\left[\chi_{\xi, a}[x(t)]\frac{1}{t} \int_{[0, t]} x(s)m_L(ds)\right].$$

An interchange of the order of the two integrations on the right side of (4.2), one with respect to the Lebesgue measure and the other with respect to the Wiener measure, can be justified as follows. Recall first that the Wiener process is a measurable process, i.e.  $x(s)$  for  $(s, x) \in [0, t] \times C[0, t]$  is Lebesgue  $\times$  Wiener measurable. To apply the Fubini theorem observe that

$$|\chi_{\xi, a}[x(t)]x(s)| \leq \frac{1}{2a}|x(s)| \quad \text{for } (s, x) \in [0, t] \times C[0, t]$$

and by (2.4)

$$\begin{aligned} \int_{[0, t]} E^w\left[\frac{1}{2a}|x(s)|\right]m_L(ds) &= \frac{1}{2a} \int_{[0, t]} \left\{\frac{1}{\sqrt{2\pi s}} \int_{R^1} |\eta| e^{-\eta^2/2s} m_L(d\eta)\right\} m_L(ds) \\ &= \frac{1}{2a} \sqrt{\frac{2}{\pi}} \int_{[0, t]} \sqrt{s} m_L(ds) = \frac{1}{3a} \sqrt{\frac{2}{\pi}} t^{3/2} < \infty. \end{aligned}$$

Thus an interchange of the two integrals on the right side of (4.2) is justified and consequently

$$\begin{aligned} E^w[(\chi_{\xi, a} \circ X)Z] &= \frac{1}{t} \int_{[0, t]} E^w[\chi_{\xi, a}[x(t)]x(s)]m_L(ds) \\ (4.3) \quad &= \frac{1}{2at} \int_{[0, t]} \{(2\pi)^2 s(t-s)\}^{-1/2} \\ &\quad \times \left[ \int_{R^1 \times [\xi-a, \xi+a]} \eta \exp\left\{-\frac{1}{2} \frac{\eta^2}{s} - \frac{1}{2} \frac{(\zeta-\eta)^2}{t-s}\right\} m_L(d(\eta, \zeta)) \right] m_L(ds) \end{aligned}$$

by (2.4). Since

$$(4.4) \quad \frac{\eta^2}{s} + \frac{(\zeta-\eta)^2}{t-s} = \frac{t}{s(t-s)} \left(\eta - \frac{s}{t}\zeta\right)^2 + \frac{\zeta^2}{t}$$

and thus

$$\begin{aligned} &\eta \exp\left\{-\frac{1}{2} \frac{\eta^2}{s} - \frac{1}{2} \frac{(\zeta-\eta)^2}{t-s}\right\} \\ &= \left\{\left(\eta - \frac{s}{t}\zeta\right) + \frac{s}{t}\zeta\right\} \exp\left\{-\frac{1}{2} \frac{t}{s(t-s)} \left(\eta - \frac{s}{t}\zeta\right)^2 - \frac{1}{2} \frac{\zeta^2}{t}\right\} \end{aligned}$$

the integral with respect to  $m_L(d(\eta, \zeta))$  over  $R^1 \times [\xi - a, \xi + a]$  in (4.3) can be reduced by integrating with respect to  $m_L(d\eta)$  over  $R^1$  and using

$$(4.5) \quad \frac{1}{\sqrt{2\pi v}} \int_{R^1} y^p e^{-y^2/2v} m_L(dy) = 1, 0, v \text{ for } p = 0, 1, 2$$

respectively, with  $v > 0$

to

$$\sqrt{\frac{2s^3(t-s)\pi}{t^3}} \int_{[\xi-a, \xi+a]} \zeta \exp \left\{ -\frac{1}{2} \frac{\zeta^2}{t} \right\} m_L(d\zeta) .$$

Using this in (4.3) we have

$$(4.6) \quad \begin{aligned} & E^w[\chi_{\xi, a} \circ X | Z] \\ &= \frac{1}{\sqrt{2\pi t} 2at^2} \left\{ \int_{[0, t]} sm_L(ds) \right\} \left\{ \int_{[\xi-a, \xi+a]} \zeta \exp \left\{ -\frac{1}{2} \frac{\zeta^2}{t} \right\} m_L(d\zeta) \right\} \\ &= \frac{1}{2} \left[ \frac{1}{2a} \int_{[\xi-a, \xi+a]} \zeta \frac{1}{\sqrt{2\pi t}} \exp \left\{ -\frac{1}{2} \frac{\zeta^2}{t} \right\} m_L(d\zeta) \right] . \end{aligned}$$

From (4.1) and (4.6) we have

$$E^w(Z | X)(\xi) \frac{dP_X(\xi)}{dm_L(\xi)} = \frac{1}{2} \xi \frac{1}{\sqrt{2\pi t}} \exp \left\{ -\frac{1}{2} \frac{\xi^2}{t} \right\}$$

and thus

$$E^w(Z | X)(\xi) = \frac{1}{2} \xi .$$

EXAMPLE 2. Let us find  $E^w(Z | X)(\xi)$ ,  $\xi \in R^1$ , when

$$Z[x] = \int_0^t [x(s)]^2 ds \quad \text{and} \quad X[x] = x(t) \quad \text{for } x \in C[0, t] .$$

To apply Corollary of Lemma 1 we proceed as in Example 1 to obtain

$$(4.7) \quad \begin{aligned} E^w[\chi_{\xi, a} \circ X | Z] &= E^w \left[ \chi_{\xi, a}[x(t)] \int_0^t [x(s)]^2 ds \right] \\ &= \int_{[0, t]} E^w[\chi_{\xi, a}[x(t)] \{x(s)\}^2] m_L(ds) \\ &= \frac{1}{2a} \int_{[0, t]} \left\{ (2\pi)^2 s(t-s) \right\}^{-1/2} \\ &\quad \times \left[ \int_{R^1 \times [\xi-a, \xi+a]} \eta^2 \exp \left\{ -\frac{1}{2} \frac{\eta^2}{s} - \frac{1}{2} \frac{(\zeta - \eta)^2}{t-s} \right\} m_L(d(\eta, \zeta)) \right] m_L(ds) . \end{aligned}$$

By means of (4.4) we have

$$\begin{aligned} & \eta^2 \exp \left\{ -\frac{1}{2} \frac{\eta^2}{s} - \frac{1}{2} \frac{(\zeta - \eta)^2}{t-s} \right\} \\ &= \left\{ \left( \eta - \frac{s}{t} \zeta \right)^2 + 2 \frac{s}{t} \zeta \left( \eta - \frac{s}{t} \zeta \right) + \left( \frac{s}{t} \zeta \right)^2 \right\} \end{aligned}$$

$$\times \exp \left\{ -\frac{1}{2} \frac{t}{s(t-s)} \left( \eta - \frac{s}{t} \zeta \right)^2 - \frac{1}{2} \frac{\zeta^2}{t} \right\}.$$

Reducing the integral with respect to  $m_L(d(\eta, \zeta))$  over  $R^1 \times [\xi - a, \xi + a]$  by integrating with respect to  $m_L(d\eta)$  over  $R^1$  using (4.5) we obtain

$$\int_{[\xi-a, \xi+a]} \sqrt{\frac{2\pi s(t-s)}{t}} \left\{ \frac{s(t-s)}{t} + \left( \frac{s}{t} \zeta \right)^2 \right\} e^{-\zeta^2/2t} m_L(d\zeta).$$

Using this in (4.7) we have

$$\begin{aligned} E^w[(\chi_{\xi, a} \circ X)Z] &= \frac{1}{\sqrt{2\pi t} 2a} \int_{[\xi-a, \xi+a]} \left[ \int_{[0, t]} \left( s - \frac{s^2}{t} + \frac{s^2}{t^2} \zeta^2 \right) m_L(ds) \right] e^{-\zeta^2/2t} m_L(d\zeta) \\ &= \frac{1}{2a} \int_{[\xi-a, \xi+a]} \left( \frac{t^2}{6} + \frac{t}{3} \zeta^2 \right) \frac{1}{\sqrt{2\pi t}} e^{-\zeta^2/2t} m_L(d\zeta). \end{aligned}$$

Thus

$$\begin{aligned} (4.8) \quad E^w(Z|X)(\xi) \frac{dP_X}{dm_L}(\xi) &= \lim_{a \rightarrow 0} E^w[(\chi_{\xi, a} \circ X)Z] \\ &= \left( \frac{t^2}{6} + \frac{t}{3} \xi^2 \right) \frac{1}{\sqrt{2\pi t}} e^{-\xi^2/2t} \end{aligned}$$

and

$$E^w(Z|X)(\xi) = \frac{t^2}{6} + \frac{t}{3} \xi^2.$$

It is of interest to note that from (2.5) and (4.8)

$$\begin{aligned} E^w[Z] &= \int_{X^{-1}(R^1)} Z[x] m_w(dx) = \int_{R^1} E^w(Z|X)(\xi) P_X(d\xi) \\ &= \frac{1}{\sqrt{2\pi t}} \int_{R^1} \left( \frac{t^2}{6} + \frac{t}{3} \xi^2 \right) e^{-\xi^2/2t} m_L(d\xi) = \frac{t^2}{2} \end{aligned}$$

as can be obtained by a direct computation of  $E^w[Z]$ .

As an example of application of an inversion formula in evaluating a condition Wiener integral, consider the Wiener measure space  $(C[0, t], \mathfrak{B}^*, m_w)$  with fixed  $t \in (0, \infty)$ . Let  $\{\varphi_1, \dots, \varphi_n\}$  be an orthogonal system in the real Hilbert space  $L^2([0, t])$  with

$$\|\varphi_j\| = \sqrt{\int_{[0, t]} \{\varphi_j(s)\}^2 m_L(ds)} > 0 \quad \text{for } j = 1, 2, \dots, n$$

such that  $\varphi_j$  has a representative function which is of bounded variation

on  $[0, t]$  for  $j = 1, 2, \dots, n$ . Let  $F$  be a functional on  $C[0, t]$  defined by

$$(4.9) \quad F[x]f\left[\int_0^t \varphi_1(s)dx(s), \dots, \int_0^t \varphi_n(s)dx(s)\right] \quad \text{for } x \in C[0, t]$$

where  $f$  is a complex valued function on  $R^n$  whose real and imaginary parts are Baire functions. By a slight extension of a well known theorem by R. E. A. C. Paley, N. Wiener, and A. Zygmund [3]  $F$  is Wiener measurable and

$$(4.10) \quad E^w[F] = \left\{ (2\pi)^n \prod_{j=1}^n \|\varphi_j\|^2 \right\}^{-1/2} \int_{R^n} f(\eta) \exp \left\{ -\sum_{j=1}^n \frac{1}{2} \frac{\eta_j^2}{\|\varphi_j\|^2} \right\} m_L(d\eta)$$

with  $\eta = (\eta_1, \dots, \eta_n) \in R^n$ , in the sense that the existence of one side implies that of the other and the equality of the two.

**THEOREM 4.** *Let  $X$  and  $Y$  be two functionals on  $C[0, t]$  for fixed  $t \in (0, \infty)$  defined by*

$$X[x] = x(t)$$

and

$$(4.11) \quad Y[x] = g\left[\int_0^t \varphi_1(s)dx(s), \dots, \int_0^t \varphi_n(s)dx(s)\right] \quad \text{for } x \in C[0, t]$$

where  $\{\varphi_1, \dots, \varphi_n, 1\}$  is an orthogonal system in the real Hilbert space  $L^2([0, t])$ ,  $\|\varphi_j\| > 0$ ,  $\varphi_j$  has a representative function which is of bounded variation on  $[0, t]$  for  $j = 1, 2, \dots, n$  and  $g$  is a Baire function on  $R^n$  such that

$$(4.12) \quad A = \left\{ (2\pi)^n \prod_{j=1}^n \|\varphi_j\|^2 \right\}^{-1/2} \int_{R^n} g(\eta) \exp \left\{ -\sum_{j=1}^n \frac{1}{2} \frac{\eta_j^2}{\|\varphi_j\|^2} \right\} m_L(d\eta) < \infty$$

with  $\eta = (\eta_1, \dots, \eta_n) \in R^n$ . Then  $Y$  is Wiener integrable on  $C[0, t]$  and

$$(4.13) \quad E^w[Y] = A.$$

Also there exists a version of  $E^w(Y|X)$  such that

$$(4.14) \quad E^w(Y|X)(\xi) = A \quad \text{for } \xi \in R^1.$$

*Proof.* By applying (4.10) to (4.11) we have (4.13). To prove (4.14) note that

$$E^w[e^{iuX} Y] = E^w\left[\exp\left\{iu \int_0^t dx(s)\right\} g\left[\int_0^t \varphi_1(s)dx(s), \dots, \int_0^t \varphi_n(s)dx(s)\right]\right].$$

Applying (4.10) to the function

$$f(\eta_1, \dots, \eta_n, \xi) = e^{i u \xi} g(\eta_1, \dots, \eta_n) \quad \text{for } (\eta_1, \dots, \eta_n, \xi) \in \mathbb{R}^{n+1}$$

we have

$$E^w[e^{i u X} Y] = A \frac{1}{\sqrt{2\pi t}} \int_{\mathbb{R}^1} e^{i u \xi} e^{-\xi^2/2t} m_L(dt) = A e^{-u^2 t/2}$$

since

$$(4.15) \quad \int_{\mathbb{R}^1} e^{-(a\xi^2 + b\xi)} m_L(d\xi) = \sqrt{\frac{\pi}{a}} e^{b^2/4a} \quad \text{for } a > 0 \text{ and real or imaginary } b.$$

We have just shown that  $E^w[e^{i u X} Y]$  is a  $m_L$ -integrable function of  $u$  on  $\mathbb{R}^1$ . Thus by Theorem 3 there exists a version of  $E^w(Y|X)dP_X/dm_L$  such that

$$E^w(Y|X)(\xi) \frac{dP_X(\xi)}{dm_L} = \frac{A}{2\pi} \int_{\mathbb{R}^1} e^{-i u \xi} e^{-u^2 t/2} m_L(du) = \frac{A}{2\pi t} e^{-\xi^2/2t}$$

and

$$E^w(Y|X)(\xi) = A \quad \text{for } \xi \in \mathbb{R}^1.$$

5. The conditional Wiener integral of  $\exp\left\{-\int_0^t V[x(s)]ds\right\}$  given  $x(t)$ .

THEOREM 5. For  $t \in (0, \infty)$  let

$$(5.1) \quad X_t[x] = x(t) \text{ and } Y_t[x] = \exp\left\{-\int_0^t V[x(s)]ds\right\} \text{ for } x \in C[0, t]$$

where  $V$  is a nonnegative continuous function on  $\mathbb{R}^1$  satisfying the condition

$$(5.2) \quad \int_{\mathbb{R}^1} V(\xi) e^{-\xi^2/2t} m_L(d\xi) < \infty \quad \text{for every } t \in (0, \infty).$$

There exists a version of  $E^w(Y_t|X_t)$  such that the function  $U$  on  $\mathbb{R}^1 \times (0, \infty)$  defined by

$$(5.3) \quad U(\xi, t) = E^w(Y_t|X_t)(\xi) \frac{1}{\sqrt{2\pi t}} e^{-\xi^2/2t}$$

satisfies the integral equation

$$(5.4) \quad U(\xi, t) = \frac{1}{\sqrt{2\pi t}} e^{-\xi^2/2t} - \int_{[0, t]} \left\{ \int_{\mathbb{R}^1} V(\eta) \frac{1}{\sqrt{2\pi(t-s)}} e^{-1/2((\xi-\eta)^2/t-s)} U(\eta, s) m_L(d\eta) \right\} m_L(ds)$$

for  $(\xi, t) \in R^1 \times (0, \infty)$ .

REMARK. From the continuity of  $V$  on  $R^1$  and the continuity of  $x$  on  $[0, t]$ ,  $\int_0^t V[x(s)]ds$  can be given as the limit of a sequence of Riemann sums corresponding to a sequence of partitions on  $[0, t]$  which does not depend on  $x$ . From this follows the Wiener measurability of  $Y_t$  on  $C[0, t]$ . Its Wiener integrability is obvious since it is bounded by 1.

The condition (5.2) on  $V$  is satisfied if for instance  $V$  satisfies the order of growth condition

$$V(\xi) = O(e^{\xi^2 - \delta}) \text{ as } \xi \longrightarrow \pm \infty$$

for some  $\delta \in (0, 2)$ . Note also that under (5.2), if we define

$$(5.5) \quad \varphi(t) = \frac{1}{\sqrt{2\pi t}} \int_{R^1} V(\xi) e^{-\xi^2/2t} m_L(d\xi) \quad \text{for } t \in (0, \infty)$$

then  $\varphi$  is a nonnegative continuous function on  $(0, \infty)$  and furthermore  $\lim_{t \rightarrow 0} \varphi(t) = V(0)$ . Let us define

$$\varphi(0) = \lim_{t \rightarrow 0} \varphi(t)$$

so that  $\varphi$  is continuous on  $[0, \infty)$ .

*Proof of Theorem 5.* According to Theorem 1, there exists a version of  $E^w(Y_t | X_t)$  such that our  $U$  defined by (5.3) can be given as

$$(5.6) \quad U(\xi, t) = \lim_{a \rightarrow \infty} \frac{1}{2\pi} \int_{(-a, a)} \left(1 - \frac{|u|}{a}\right) e^{-iu\xi} E^w[e^{-iuX_t} Y_t] m_L(du) \\ \text{for } (\xi, t) \in R^1 \times (0, \infty).$$

Since

$$\frac{d}{ds} \exp \left\{ - \int_0^s V[x(s)] dr \right\} = - \exp \left\{ - \int_0^s V[x(r)] dr \right\} V[x(s)]$$

we have by integrating with respect to  $s$  on  $[0, t]$

$$(5.7) \quad \exp \left\{ - \int_0^t V[x(s)] ds \right\} - 1 = - \int_0^t V[x(s)] \exp \left\{ - \int_0^s V[x(r)] dr \right\} ds.$$

Applying (5.7) to  $Y_t[x]$  as defined by (5.1) we have

$$(5.8) \quad Y_t[x] = 1 - \int_0^t V[x(s)] \exp \left\{ - \int_0^s V[x(r)] dr \right\} ds.$$

Substituting (5.8) in (5.6) we have

$$(5.9) \quad U(\xi, t) = I_1(\xi, t) - I_2(\xi, t)$$

where

$$(5.10) \quad I_1(\xi, t) = \lim_{a \rightarrow \infty} \frac{1}{2\pi} \int_{(-a, a)} \left(1 - \frac{|u|}{a}\right) e^{-iu\xi} E^w[e^{iux(t)}] m_L(du)$$

and

$$(5.11) \quad I_2(\xi, t) = \lim_{a \rightarrow \infty} \frac{1}{2\pi} \int_{(-a, a)} \left(1 - \frac{|u|}{a}\right) e^{-iu\xi} J(t) m_L(du)$$

with

$$(5.12) \quad J(t) = E^w \left[ e^{iux(t)} \int_{[0, t]} V[x(s)] \exp \left\{ - \int_0^s V[x(r)] dr \right\} m_L(ds) \right].$$

To evaluate  $I_1$ , note that by (2.4) and (4.15) we have

$$E^w[e^{iux(t)}] = \frac{1}{\sqrt{2\pi t}} \int_{R^1} e^{i u \eta} e^{-\eta^2/2t} m_L(d\eta) = e^{-u^2 t/2}$$

and thus

$$(5.13) \quad \begin{aligned} I_1(\xi, t) &= \lim_{a \rightarrow \infty} \left\{ \frac{1}{2\pi} \int_{(-a, a)} e^{-iu\xi} e^{-u^2 t/2} m_L(du) - \frac{1}{2\pi} \right. \\ &\quad \left. \int_{(-a, a)} \frac{|u|}{a} e^{-iu\xi} e^{-u^2 t/2} m_L(du) \right\} \\ &= \frac{1}{2\pi} \int_{R^1} e^{-iu\xi} e^{-u^2 t/2} m_L(du) - \frac{1}{2\pi} \left( \lim_{u \rightarrow \infty} + \lim_{u \rightarrow -\infty} \right) \left( \frac{|u|}{a} e^{-iu\xi} e^{-u^2 t/2} \right) \\ &= \frac{1}{\sqrt{2\pi t}} e^{-\xi^2/2t}. \end{aligned}$$

To interchange the order of the Wiener integral and the integral with respect to  $m_L(ds)$  on  $[0, t]$  in (5.12), note that

$$\left| e^{iux(t)} V[x(s)] \exp \left\{ - \int_0^s V[x(r)] dr \right\} \right| \leq V[x(s)]$$

for  $(s, x) \in [0, t] \times C[0, t]$

and that by (2.4), (4.15), (5.5) and the continuity of  $\varphi$  on  $[0, \infty)$

$$\begin{aligned} \int_{[0, t]} E^w[V[x(s)]] m_L(ds) &= \int_{[0, t]} \frac{1}{\sqrt{2\pi s}} \left\{ \int_{R^1} V(\zeta) e^{-\zeta^2/2s} m_L(d\zeta) \right\} m_L(ds) \\ &= \int_{[0, t]} \varphi(s) m_L(ds) < \infty \end{aligned}$$

so that the Fubini theorem is applicable and thus

$$(5.14) \quad J(t) = \int_{[0,t]} E^w \left[ e^{iux(t)} V[x(s)] \exp \left\{ - \int_0^s V[x(r)] dr \right\} \right] m_L(ds) .$$

Let us write

$$e^{iux(t)} = e^{iu\{x(t)-x(s)\}} e^{iux(s)} .$$

By the fact that  $\{x(t) - x(s), x(r)\}$  is an independent system of random variables on  $(C[0, t], \mathfrak{B}^*, m_w)$  for every  $r \in (0, s]$  and by

$$E^w [e^{iux\{x(t)-x(s)\}}] = e^{-(u^2/2)(t-s)}$$

which follows from (2.4) and (4.15), we have from (5.14)

$$(5.15) \quad J(t) = \int_{[0,t]} e^{-(u^2/2)(t-s)} E^w \left[ e^{iux(s)} V[x(s)] \exp \left\{ - \int_0^s V[x(r)] dr \right\} \right] m_L(ds) .$$

Applying Lemma 1 to the Wiener integral in (5.15) and recalling (5.1) and (5.3) we have

$$(5.16) \quad \begin{aligned} J(t) &= \int_{[0,t]} e^{-(u^2/2)(t-s)} \left[ \int_{R^1} e^{i u \eta} V(\eta) E(Y_s | X_s)(\eta) P_{X_s}(d\eta) \right] m_L(ds) \\ &= \int_{[0,t]} e^{-(u^2/2)(t-s)} \left[ \int_{R^1} e^{i u \eta} V(\eta) U(\eta, s) m_L(d\eta) \right] m_L(ds) . \end{aligned}$$

Let us use (5.16) in (5.11). To interchange the order of the three integrals with respect to  $m_L(d\eta)$  over  $R^1$ ,  $m_L(ds)$  over  $[0, t]$  and  $m_L(du)$  over  $(-a, a)$  in the resulting expression for  $I_2$ , observe that

$$\left| \left( 1 - \frac{|u|}{a} \right) e^{-i u \xi} e^{i u \eta} V(\eta) U(\eta, s) e^{-(u^2/2)(t-s)} \right| \leq V(\eta) U(\eta, s) e^{-(u^2/2)(t-s)}$$

$$\text{for } (\eta, s, u) \in R^1 \times [0, t] \times (-a, a) .$$

Recalling (5.3) and Lemma 1 we have by (2.4) and (5.5)

$$\begin{aligned} \int_{R^1} V(\eta) U(\eta, s) m_L(d\eta) &= E^w \left[ V[x(s)] \exp \left\{ - \int_0^s V[x(r)] dr \right\} \right] \\ &\leq E^w [V[x(s)]] = \frac{1}{\sqrt{2\pi s}} \int_{R^1} V(\zeta) e^{-\zeta^2/(2s)} m_L(d\zeta) = \varphi(s) . \end{aligned}$$

Now

$$\int_{[0,t]} \varphi(s) e^{-(u^2/2)(t-s)} m_L(ds) = A e^{-u^2 t/2}$$

where

$$A = \int_{[0,t]} \varphi(s) e^{u^2 s/2} m_L(ds) < \infty$$

by the continuity of  $\varphi$  on  $[0, \infty)$  and thus



$$\int_{R^1} A e^{-u^2 t/2} m_L(du) < \infty .$$

This verifies the applicability of the Fubini theorem in interchanging the order of the three integrals. Thus we have

$$(5.17) \quad I_2(\xi, t) = \lim_{a \rightarrow \infty} \frac{1}{2\pi} \int_{[0, t]} \left\{ \int_{R^1} V(\eta) U(\eta, s) \left[ \int_{R^1} \chi_{(-a, a)}(u) \left( 1 - \frac{|u|}{a} \right) \right. \right. \\ \left. \left. \times e^{-i u (\xi - \eta)} e^{-(u^2/2)(t-s)} m_L(du) \right] m_L(d\eta) \right\} m_L(ds) .$$

For all  $a > 0$  the integrand in (5.17) is bounded by

$$V(\eta) U(\eta, s) e^{-(u^2/2)(t-s)} \quad \text{for } (\eta, s, u) \in R^1 \times [0, t] \times R^1$$

which is integrable with respect to  $(m_L \times m_L \times m_L)(d(\eta, s, u))$  on  $R^1 \times [0, t] \times R^1$  as we saw above. Thus by the Dominated Convergence Theorem

$$(5.18) \quad I_2(\xi, t) = \frac{1}{2\pi} \int_{[0, t]} \left\{ \int_{R^1} V(\eta) U(\eta, s) \right. \\ \left. \times \left[ \int_{R^1} e^{-i u (\xi - \eta)} e^{-(u^2/2)(t-s)} m_L(du) \right] m_L(d\eta) \right\} m_L(ds) \\ = \int_{[0, t]} \left\{ \int_{R^1} V(\eta) U(\eta, s) \frac{1}{\sqrt{2\pi(t-s)}} e^{-(\xi - \eta)^2 / (2(t-s))} m_L(d\eta) \right\} m_L(ds) .$$

Using (5.13) and (5.18) in (5.9) we have (5.4).

#### REFERENCES

1. R. R. Goldberg, *Fourier Transforms*, Cambridge University Press, 1961.
2. M. Kac, *Probability and Related Topics in Physical Sciences*, Interscience Publishers, New York, 1959.
3. R. E. A. C. Paley, N. Wiener and A. Zygmund, *Notes on random functions*, *Mathematische Zeitschrift*, **37** (1933), 647-668.
4. J. Yeh, *Inversion of conditional expectations*, *Pacific J. Math.*, **52** (1974), 631-640.

Received May 12, 1975.

UNIVERSITY OF CALIFORNIA, IRVINE

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Aharon Atzmon, <i>A moment problem for positive measures on the unit disc</i> .....	317
Peter W. Bates and Grant Bernard Gustafson, <i>Green's function inequalities for two-point boundary value problems</i> .....	327
Howard Edwin Bell, <i>Infinite subrings of infinite rings and near-rings</i> .....	345
Grahame Bennett, Victor Wayne Goodman and Charles Michael Newman, <i>Norms of random matrices</i> .....	359
Beverly L. Brechner, <i>Almost periodic homeomorphisms of <math>E^2</math> are periodic</i> .....	367
Beverly L. Brechner and R. Daniel Mauldin, <i>Homeomorphisms of the plane</i> .....	375
Jia-Arng Chao, <i>Lusin area functions on local fields</i> .....	383
Frank Rimi DeMeyer, <i>The Brauer group of polynomial rings</i> .....	391
M. V. Deshpande, <i>Collectively compact sets and the ergodic theory of semi-groups</i> .....	399
Raymond Frank Dickman and Jack Ray Porter, <i><math>\theta</math>-closed subsets of Hausdorff spaces</i> .....	407
Charles P. Downey, <i>Classification of singular integrals over a local field</i> .....	417
Daniel Reuven Farkas, <i>Miscellany on Bieberbach group algebras</i> .....	427
Peter A. Fowler, <i>Infimum and domination principles in vector lattices</i> .....	437
Barry J. Gardner, <i>Some aspects of T-nilpotence. II: Lifting properties over T-nilpotent ideals</i> .....	445
Gary Fred Gruenhagen and Phillip Lee Zenor, <i>Metritzation of spaces with countable large basis dimension</i> .....	455
J. L. Hickman, <i>Reducing series of ordinals</i> .....	461
Hugh M. Hilden, <i>Generators for two groups related to the braid group</i> .....	475
Tom (Roy Thomas Jr.) Jacob, <i>Some matrix transformations on analytic sequence spaces</i> .....	487
Elyahu Katz, <i>Free products in the category of <math>k_w</math>-groups</i> .....	493
Tsang Hai Kuo, <i>On conjugate Banach spaces with the Radon-Nikodým property</i> .....	497
Norman Eugene Liden, <i>K-spaces, their antispace and related mappings</i> .....	505
Clinton M. Petty, <i>Radon partitions in real linear spaces</i> .....	515
Alan Saleski, <i>A conditional entropy for the space of pseudo-Menger maps</i> .....	525
Michael Singer, <i>Elementary solutions of differential equations</i> .....	535
Eugene Spiegel and Allan Trojan, <i>On semi-simple group algebras. I</i> .....	549
Charles Madison Stanton, <i>Bounded analytic functions on a class of open Riemann surfaces</i> .....	557
Sherman K. Stein, <i>Transversals of Latin squares and their generalizations</i> .....	567
Ivan Ernest Stux, <i>Distribution of squarefree integers in non-linear sequences</i> .....	577
Lowell G. Sweet, <i>On homogeneous algebras</i> .....	585
Lowell G. Sweet, <i>On doubly homogeneous algebras</i> .....	595
Florian Vasilescu, <i>The closed range modulus of operators</i> .....	599
Arthur Anthony Yanushka, <i>A characterization of the symplectic groups <math>P\text{Sp}(2m, q)</math> as rank 3 permutation groups</i> .....	611
James Juei-Chin Yeh, <i>Inversion of conditional Wiener integrals</i> .....	623