

# Pacific Journal of Mathematics

**COMPACTNESS-LIKE PROPERTIES FOR GENERALIZED  
WEAK TOPOLOGICAL SUMS**

W. WISTAR (WILLIAM) COMFORT

# COMPACTNESS-LIKE PROPERTIES FOR GENERALIZED WEAK TOPOLOGICAL SUMS

W. W. COMFORT

It is shown that under suitable conditions (involving cardinal numbers) on a family of spaces  $\{X_i : i \in I\}$  with  $p_i \in X_i$  for  $i \in I$ , their  $\gamma$ -weak sum  $\{x \in \prod_{i \in I} X_i : |\{i \in I : x_i \neq p_i\}| < \gamma\}$  is  $\alpha$ -compact in the  $\kappa$ -box topology. For example, there is Corollary 2.5: If  $\alpha$  is regular and uncountable and  $|X_i| < \alpha$  for all  $i \in I$ , then the  $\omega$ -weak sum (= direct sum) is  $\alpha$ -compact in the  $\alpha$ -box topology; in particular, the direct sum of any set of finite spaces is  $\alpha$ -compact in the  $\alpha$ -complete topology for regular  $\alpha > \omega$ .

**1. Notation and definitions.** We denote the smallest infinite cardinal by  $\omega$ , and for a cardinal  $\alpha$  we denote by  $\alpha^+$  the smallest cardinal  $\beta$  such that  $\beta > \alpha$ . For  $\alpha \geq \omega$  we denote by  $\text{cf}(\alpha)$  the smallest cardinal  $\beta$  for which there is a family  $\{\alpha_\xi : \xi < \beta\}$  of cardinals such that

$$\alpha_\xi < \alpha \quad \text{for all } \xi < \beta \quad \text{and}$$

$$\sum_{\xi < \beta} \alpha_\xi = \alpha.$$

The (infinite) cardinal  $\alpha$  is *regular* if  $\alpha = \text{cf}(\alpha)$ , *singular* otherwise. For cardinals  $\alpha$  and  $\gamma$  we set

$$\alpha^\gamma = \sum \{\alpha^\beta : \beta \text{ is a cardinal and } \beta < \gamma\}.$$

and if  $I$  is a set we define

$$\mathcal{P}(I) = \{J : J \subset I\} \quad \text{and}$$

$$\mathcal{P}_\gamma(I) = \{J \in \mathcal{P}(I) : |J| < \gamma\}.$$

It is well-known and easy to prove (see for example §1 of [5]) that

$$\gamma \leq \alpha^\gamma \quad \text{if } 2 \leq \alpha, \quad \text{and}$$

$$|\mathcal{P}_\gamma(\alpha)| = \alpha^\gamma \quad \text{if } \omega + \gamma \leq \alpha^+.$$

Throughout this paper we denote by  $\{X_i : i \in I\}$  or by  $\{X_\xi : \xi < \alpha\}$  a set of nonempty topological spaces (not assumed to satisfy any special

separation axioms), we fix  $p_i \in X_i$ , and for every infinite cardinal  $\gamma$  we set

$$\gamma \prod_{i \in I} X_i = \left\{ x \in \prod_{i \in I} X_i : |\{i \in I : x_i \neq p_i\}| < \gamma \right\}.$$

When  $\gamma = \omega$ , the space  $\gamma \prod_{i \in I} X_i$  is called the *weak topological sum* of  $\{X_i : i \in I\}$ .

If  $U_i \subset X_i$  and  $U = \prod_{i \in I} U_i$ , we set

$$R(U) = \{i \in I : U_i \neq X_i\};$$

for a family  $\mathcal{U}$  of such sets  $U$  we set  $R(\mathcal{U}) = \cup \{R(U) : U \in \mathcal{U}\}$ . For an infinite cardinal  $\kappa$ , the  $\kappa$ -*box topology* on  $\prod_{i \in I} X_i$  is the topology generated by sets of the form  $U = \prod_{i \in I} U_i$  for which

$U_i$  is open in  $X_i$  for all  $i$ , and

$$|R(U)| < \kappa.$$

The space  $\prod_{i \in I} X_i$  with the  $\kappa$ -box topology is denoted  $(\prod_{i \in I} X_i)_\kappa$ , and for  $Y \subset \prod_{i \in I} X_i$  we denote by  $Y_\kappa$  the set  $Y$  with the topology inherited from  $(\prod_{i \in I} X_i)_\kappa$ . We note that  $(\prod_{i \in I} X_i)_\omega$  is  $\prod_{i \in I} X_i$  with the usual product topology, and that if  $|I| < \kappa$  then  $(\prod_{i \in I} X_i)_\kappa$  is  $\prod_{i \in I} X_i$  with the usual box topology.

The sets  $U = \prod_{i \in I} U_i$  as above are elements of the *canonical basis* for  $(\prod_{i \in I} X_i)_\kappa$  and if  $Y \subset \prod_{i \in I} X_i$  then the *canonical basis* for  $Y_\kappa$  consists of all sets of the form  $U \cap Y$  (with  $U$  as above).

Let  $\alpha \leq \beta$ . A space  $X$  is said to be  $[\alpha, \beta]$ -compact if for every open cover  $\mathcal{U}$  of  $X$  such that  $|\mathcal{U}| < \beta$  there is  $\mathcal{V} \in \mathcal{P}_\alpha(\mathcal{U})$  such that  $X = \cup \mathcal{V}$ ; and  $X$  is  $\alpha$ -compact if  $X$  is  $[\alpha, |\mathcal{P}(X)|^+]$ -compact, i.e., if for every open cover  $\mathcal{U}$  of  $X$  there is  $\mathcal{V} \in \mathcal{P}_\alpha(\mathcal{U})$  such that  $X = \cup \mathcal{V}$ . If  $\mathcal{B}$  is a basis for  $X$  then  $X$  is  $[\alpha, \beta]$ -compact with respect to  $\mathcal{B}$  if for every cover  $\mathcal{U} \in \mathcal{P}_\beta(\mathcal{B})$  there is  $\mathcal{V} \in \mathcal{P}_\alpha(\mathcal{U})$  such that  $X = \cup \mathcal{V}$ .

We note that in this terminology a space is compact if and only if it is  $\omega$ -compact, and Lindelöf if and only if it is  $\omega^+$ -compact.

For a survey of properties related to  $[\alpha, \beta]$ -compactness see Vaughan [14], [15]; for connections with topological groups, see Comfort and Saks [8]; and for product-space theorems relating to  $[\alpha, \beta]$ -compact spaces, see Vaughan [13] and Greene [12]. In these, notation and terminology differ slightly from that above.

There is at least one assertion concerning  $\alpha$ -compactness of the space  $X = \omega \prod_{i \in I} X_i$  already available in the literature: It is pointed out by Corson [9] (Proposition 4) that if each of the spaces  $X_i$  is  $\sigma$ -compact, then  $X$  is  $\sigma$ -compact (and hence is a Lindelöf space).

It is not pleasing to be forced to introduce into the picture the bizarre concept “[ $\alpha, \beta$ ]-compact with respect to  $\mathcal{B}$ ”. As partial justification, and as an explanation of its appearance in Theorem 2.3 below, we note that for every  $\beta \cong \omega$  there are a completely regular Hausdorff space  $X$  and a basis  $\mathcal{B}$  for  $X$  such that  $X$  is [ $\alpha, \beta$ ]-compact with respect to  $\mathcal{B}$  for all cardinals  $\alpha$  such that  $2 \cong \alpha \cong \beta$ , and  $X$  is not [ $\alpha, \beta$ ]-compact for any cardinal  $\alpha < \beta$ . Indeed, we may take for  $X$  the discrete space  $\beta$  and for  $\mathcal{B}$  the basis

$$\mathcal{B} = \{X\} \cup \{\{\xi\}: \xi < \beta\}.$$

## 2. Compactness for weak topological sums.

LEMMA 2.1. *Let  $\omega \cong \kappa$ , and let  $I$  be a set and  $\alpha, \beta$  and  $\gamma$  cardinals such that*

$$\omega \cong \gamma \cong |I|^z < \text{cf}(\alpha) \cong \alpha \cong \beta.$$

*If  $(\prod_{i \in I} X_i)_\kappa$  is [ $\alpha, \beta$ ]-compact for all  $J \in \mathcal{P}_\gamma(I)$ , then  $(\gamma \prod_{i \in I} X_i)_\kappa$  is [ $\alpha, \beta$ ]-compact.*

*Proof.* If  $|I| < \gamma$  then with  $J = I$  we have

$$\left(\gamma \prod_{i \in I} X_i\right)_\kappa = \left(\prod_{i \in I} X_i\right)_\kappa = \left(\prod_{i \in J} X_i\right)_\kappa$$

and the statement is obvious. We assume therefore that  $|I| \cong \gamma$ .

For  $J \in \mathcal{P}_\gamma(I)$  we set

$$Y(J) = \left\{x \in \gamma \prod_{i \in I} X_i: x_i = p_i \text{ for all } i \in I \setminus J\right\}.$$

Clearly  $\gamma \prod_{i \in I} X_i = \cup \{Y(J): J \in \mathcal{P}_\gamma(I)\}$ , and for  $J \in \mathcal{P}_\gamma(I)$  the space  $(Y(J))_\kappa$  is homeomorphic to  $(\prod_{i \in J} X_i)_\kappa$ . Now let  $\mathcal{U}$  be an open cover of  $(\gamma \prod_{i \in I} X_i)_\kappa$  such that  $|\mathcal{U}| < \beta$ . For every  $J \in \mathcal{P}_\gamma(I)$  there is  $\mathcal{V}(J) \in \mathcal{P}_\alpha(\mathcal{U})$  such that  $Y(J) \subset \cup \mathcal{V}(J)$ . Then  $\cup \{\mathcal{V}(J): J \in \mathcal{P}_\gamma(I)\}$  is a cover of  $\gamma \prod_{i \in I} X_i$  by elements of  $\mathcal{U}$ , and since  $|\mathcal{P}_\gamma(I)| = |I|^z < \text{cf}(\alpha)$  and  $|\mathcal{V}(J)| < \alpha$  for all  $J \in \mathcal{P}_\gamma(I)$  we have

$$|\cup \{\mathcal{V}(J): J \in \mathcal{P}_\gamma(I)\}| \cong \sum \{|\mathcal{V}(J)|: J \in \mathcal{P}_\gamma(I)\} < \alpha.$$

The proof is complete.

We note that if in Lemma 2.1 it is assumed simply that  $(\prod_{i \in J} X_i)_\kappa$  is  $[\alpha, \beta]$ -compact with respect to its canonical basis for all  $J \in \mathcal{P}_\gamma(I)$ , then (as the proof shows) the space  $(\gamma \prod_{i \in I} X_i)_\kappa$  is  $[\alpha, \beta]$ -compact with respect to its canonical basis.

**REMARK 2.2.** We note that if for every  $i \in I$  the space  $X_i$  is a  $T_1$ -space such that  $|X_i| \geq 2$ , and if further  $\gamma \leq \kappa$ ,  $\gamma \leq |I|$  and  $\beta \geq (2^\gamma)^+$ , then we have  $2^\lambda < \alpha$  for all  $\lambda < \gamma$ . For under these (additional) assumptions the space  $(2^J)_\kappa$  is a closed, discrete subspace of  $(\prod_{i \in J} X_i)_\kappa$  (and is therefore  $\alpha$ -compact) for all  $J \in \mathcal{P}_\gamma(I)$ , and hence  $2^{|\mu|} < \alpha$  for all  $J \in \mathcal{P}_\gamma(I)$ .

**DEFINITION.** Let  $\alpha$  and  $\gamma$  be cardinals such that  $\gamma \leq \alpha$ . Then  $\alpha$  is *strongly  $\gamma$ -inaccessible* if  $\beta^\lambda < \alpha$  whenever  $\beta < \alpha$  and  $\lambda < \gamma$ .

We note that if  $\omega \leq \gamma < \alpha$  and  $\alpha$  is regular then the following three conditions are equivalent:

- (i)  $\alpha$  is strongly  $\gamma$ -inaccessible;
- (ii)  $\prod_{\xi < \lambda} \alpha_\xi < \alpha$  whenever  $\lambda < \gamma$  and  $\alpha_\xi < \alpha$  for all  $\xi < \lambda$ ;
- (iii)  $\beta^\lambda < \alpha$  for all  $\beta < \alpha$ .

It has been shown by Erdős and Rado [10] that for  $\alpha$  a regular infinite cardinal and  $\gamma < \alpha$ , the following combinatorial condition is equivalent to (i)–(iii).

(iv) if  $\{S_\xi : \xi < \alpha\}$  is a family of sets such that  $|S_\xi| < \gamma$  for all  $\xi < \alpha$ , then there is  $A \subset \alpha$  and a set  $B$  such that  $|A| = \alpha$  and  $S_\xi \cap S_\eta = B$  whenever  $\xi, \eta \in A$  and  $\xi \neq \eta$ .

For another proof of the equivalence (i)  $\Leftrightarrow$  (iv) and several topological equivalences, see [3]; and for applications to generalized  $\Sigma$ -spaces see [4] and [6]. The notion of pairs  $\gamma, \alpha$  such that  $\omega \leq \gamma < \alpha$  and  $\alpha$  is strongly  $\gamma$ -inaccessible has been introduced independently and used by Fuhrken [11] in connection with a generalization of Loś' ultraproduct theorem.

**THEOREM 2.3.** Let  $\omega \leq \kappa \leq \alpha$  and  $\omega \leq \gamma \leq \alpha \leq \beta$  with  $\alpha$  regular and strongly  $\gamma$ -inaccessible, let  $\{X_i : i \in I\}$  be a family of spaces such that  $(\prod_{i \in J} X_i)_\kappa$  is  $[\alpha, \beta]$ -compact for all  $J \in \mathcal{P}_\gamma(\alpha)$ , and define  $X = (\gamma \prod_{i \in I} X_i)_\kappa$ . Then  $X$  is  $[\alpha, \beta]$ -compact with respect to the canonical basis.

*Proof.* Let  $\mathcal{U}$  be a cover of  $X$  such that  $|\mathcal{U}| < \beta$  and the elements of  $\mathcal{U}$  are basic open sets, i.e., they are sets of the form  $U \cap X$  where  $U = \prod_{i \in I} U_i$  with  $U_i$  open in  $X_i$  for all  $i \in I$ , and  $|R(U)| < \kappa$  for all  $i \in I$ . We show there is  $\mathcal{V} \in \mathcal{P}_\alpha(\mathcal{U})$  such that  $\cup \mathcal{V} = X$ .

For  $J \subset I$  we set

$$Y(J) = \{x \in X : x_i = p_i \text{ for all } i \in I \setminus J\}$$

and we note that if  $J \in \mathcal{P}_\alpha(I)$  then since  $|J|^\gamma < \alpha = \text{cf}(\alpha)$  and  $(Y(J))_\kappa$  is homeomorphic to  $(\gamma \prod_{i \in J} X_i)_\kappa$  the space  $(Y(J))_\kappa$  is  $[\alpha, \beta]$ -compact by Lemma 2.1.

We set

$$\begin{aligned}\bar{\gamma} &= \gamma \text{ if } \gamma \text{ is regular} \\ &= \gamma^+ \text{ if } \gamma \text{ is singular}\end{aligned}$$

and we note that  $\bar{\gamma} < \alpha$  (since if  $\gamma$  is singular then  $\bar{\gamma} = \gamma^+ \leq \gamma^{\text{cf}(\gamma)} < \alpha$ ).

We observe now that there are families  $\{J_\xi: \xi < \bar{\gamma}\} \subset \mathcal{P}(I)$  and  $\{\mathcal{V}_\xi: \xi < \bar{\gamma}\} \subset \mathcal{P}(\mathcal{U})$  such that

- (i)  $J_0 = \emptyset$ ;
- (ii)  $\mathcal{V}_0 = \emptyset$ ;
- (iii)  $J_{\xi+1} = J_\xi \cup \cup \{R(V): V \in \mathcal{V}_\xi\}$  for  $\xi < \bar{\gamma}$ ;
- (iv)  $\mathcal{V}_{\xi+1}$  is a cover of  $Y(J_{\xi+1})$  and  $|\mathcal{V}_{\xi+1}| < \alpha$  for  $\xi < \bar{\gamma}$ ; and
- (v)  $J_\xi = \bigcup_{\eta < \xi} J_\eta$  and  $\mathcal{V}_\xi = \bigcup_{\eta < \xi} \mathcal{V}_\eta$  for nonzero limit ordinals

$\xi < \bar{\gamma}$ .

Indeed, these families may be defined recursively, it being noted inductively that  $|J_{\xi+1}| < \alpha$  for  $\xi < \bar{\gamma}$  (so that  $(Y(J_{\xi+1}))_\kappa$  is  $[\alpha, \beta]$ -compact and  $\mathcal{V}_{\xi+1}$  may be defined as required in (iv)).

The definition of the families  $\{J_\xi: \xi < \bar{\gamma}\}$  and  $\{\mathcal{V}_\xi: \xi < \bar{\gamma}\}$  being complete, we set

$$J = \bigcup_{\xi < \bar{\gamma}} J_\xi \quad \text{and} \quad \mathcal{V} = \bigcup_{\xi < \bar{\gamma}} \mathcal{V}_\xi.$$

Since  $|J_\xi| < \alpha$  and  $|\mathcal{V}_\xi| < \alpha$  for all  $\xi < \bar{\gamma}$  (by (iii), (iv) and (v) above) we have  $|J| < \alpha$  and  $|\mathcal{V}| < \alpha$ .

We note that if  $y \in Y(J)$  and  $y_i \neq p_i$  then  $i \in J$  and hence there is  $\xi(i) < \bar{\gamma}$  such that  $i \in J_{\xi(i)}$ . Since  $|\{i \in I: y_i \neq p_i\}| < \gamma \leq \bar{\gamma}$  and  $\bar{\gamma}$  is regular there is  $\bar{\xi} < \bar{\gamma}$  such that  $\xi(i) < \bar{\xi}$  for all  $i$  with  $y_i \neq p_i$ ; it follows that

$$y \in Y(J_{\bar{\xi}}) \subset \cup \mathcal{V}_{\bar{\xi}} \subset \cup \mathcal{V}.$$

We conclude that  $Y(J) \subset \cup \mathcal{V}$ .

We claim that in fact  $X \subset \cup \mathcal{V}$ . Indeed let  $x \in X$  and define

$$\begin{aligned}y_i &= x_i \quad \text{if } i \in J \\ &= p_i \quad \text{if } i \in I \setminus J.\end{aligned}$$

It is clear that  $(y \in X \text{ and}) y \in Y(J)$ . From the preceding paragraph there is  $V = \prod_{i \in I} V_i \in \mathcal{V}$  such that  $y \in V$ , and since  $R(V) \subset J$  we have  $V_i = X_i$  for all  $i \in I \setminus J$  and hence  $x \in V$ . The proof is complete.

**COROLLARY 2.4.** *Let  $\omega \leq \kappa \leq \alpha$  and  $\omega \leq \gamma < \alpha$  with  $\alpha$  regular and strongly  $\gamma$ -inaccessible, let  $\{X_i: i \in I\}$  be a family of spaces such that  $|X_i| < \alpha$  for all  $i \in I$ , and set  $X = (\gamma \prod_{i \in I} X_i)_\kappa$ . Then  $X$  is  $\alpha$ -compact.*

*Proof.* From the equivalence (i)  $\Leftrightarrow$  (ii) remarked above we have  $|\prod_{i \in J} X_i| < \alpha$  whenever  $J \in \mathcal{P}_\gamma(I)$ , so that  $(\prod_{i \in J} X_i)_\kappa$  is  $[\alpha, \beta]$ -compact for all  $J \in \mathcal{P}_\gamma(I)$  and  $\beta \geq \alpha$ . Thus the result follows from Theorem 2.3.

We remark that in Corollary 2.4 (and earlier) the hypothesis that  $\alpha$  is strongly  $\gamma$ -inaccessible cannot be omitted (or even substantially modified). Indeed suppose that  $\omega \leq \gamma < \alpha$  and that there are cardinals  $\beta < \alpha$  and  $\lambda < \gamma$  such that  $\beta^\lambda \geq \alpha$ ; then the power space  $(\beta^\lambda)_\alpha$ , which is  $(\gamma\beta^\lambda)_\alpha$ , is a discrete space of cardinality at least  $\alpha$  and is not  $\alpha$ -compact.

In the positive direction, every infinite cardinal is strongly  $\omega$ -inaccessible. Thus from Corollary 2.4 we have the following further specialization of Theorem 2.3.

**COROLLARY 2.5.** *Let  $\alpha$  be an uncountable, regular cardinal, let  $\{X_i: i \in I\}$  be a family of spaces such that  $|X_i| < \alpha$  for all  $i \in I$  and let  $X$  be the "weak sum in the  $\alpha$ -box topology", i.e.,  $X = (\omega \prod_{i \in I} X_i)_\alpha$ . Then  $X$  is  $\alpha$ -compact.*

**3. An application.** For a nondiscrete space  $X$  we denote by  $pX$  the smallest possible number of open subsets of  $X$  with nonopen intersection. Now the space  $2$  is an (additive) topological group with identity element  $0$ , and it is clear that for regular  $\alpha > \omega$  and with the choice  $p_\xi = 0_\xi = 0$  for all  $\xi < \beta$ , the space  $G(\alpha, \beta) = (\omega \prod_{\xi < \beta} 2_\xi)_\alpha$  is a topological group such that  $|G| = \beta$  and  $pG = \alpha$ . It is shown in [1] that the groups  $G(\alpha, \alpha)$  are *fine*, i.e., every continuous real-valued function on  $G(\alpha, \alpha)$  is uniformly continuous, and the question is raised there whether for regular  $\alpha > \omega$  there are fine groups  $G$  of arbitrarily large cardinality such that  $pG = \alpha$ . It follows from Theorem 2.3. (iii) of [1] and the fact that  $G(\alpha, \beta)$  is  $\alpha$ -compact for all  $\beta \geq \alpha$  (a special case of Corollary 2.5 above) that the answer to this question is "Yes". The present paper sheds no light on the several other questions posed in [1].

**4. Concluding remarks.** My understanding of the full generality under which the conclusions of Theorem 2.3 and Corollary 2.4 can be established has been achieved slowly, over an extend period of time, and with the substantial help of other mathematicians. It is a pleasure to cite in particular my indebtedness to:

(a) K. A. Ross. The case of Corollary 2.4 in which  $\gamma = \omega < \omega^+ = \alpha = \kappa = I$  and each  $X_i = 2$  is Example 3.2 of [7];

(b) A. W. Hager. Portions of the argument required for 2.4 above appear in 5.4 and 5.7 of [1], where we considered however only the case in which  $\gamma = \omega$  and there are no limit cardinals between  $\alpha$  and  $|I|$ .

(c) A. Hajnal and I. Juhász. Conversations following my presentation of a colloquium with this title at the Bolyai János Mathematical Society resulted in a proof [2] with combinatorial emphasis of the case  $X_i = 2$  of Corollary 2.4.

(d) K. Kunen. As referee of the present paper, he has strengthened the statement of Theorem 2.3 and shortened its proof.

### REFERENCES

1. W. W. Comfort and Anthony W. Hager, *Uniform continuity in topological groups*, in Proc. January, 1974 Rome symposium on topological groups and Lie groups, Rome, 1975, to appear.
2. W. W. Comfort, A. Hajnal, and I. Juhász, *Compactness-like properties of generalized weak products*, to appear.
3. W. W. Comfort and S. Negrepointis, *On families of large oscillation*, *Fundamenta Math.*, **75** (1972), 275–290.
4. W. W. Comfort and S. Negrepointis, *Continuous functions on products with strong topologies*, in *General topology and its relations to modern analysis and algebra III*, Proc. third (1971) Prague topological symposium, pp. 89–92, 1972.
5. W. W. Comfort and S. Negrepointis, *The theory of ultrafilters*, *Grundlehren der math. Wissenschaften Band 211*, Springer-Verlag, Heidelberg, 1974.
6. W. W. Comfort and S. Negrepointis, manuscript in preparation.
7. W. W. Comfort and Kenneth A. Ross, *Pseudocompactness and uniform continuity in topological groups*, *Pacific J. Math.*, **16** (1966), 483–486.
8. W. W. Comfort and Victor Saks, *Countably compact groups and finest totally bounded topologies*, *Pacific J. Math.*, **49** (1973), 33–44.
9. H. H. Corson, *Normality in subsets of product spaces*, *American J. Math.*, **81** (1959), 784–796.
10. P. Erdős and R. Rado, *Intersection theorems for systems of sets II*, *J. London Math. Soc.*, **44** (1969), 467–479.
11. G. Fuhrken, *Languages with added quantifier “There exist at least  $\aleph_\alpha$ ”*, in *The theory of models*, Proc. 1963 International Berkeley Symposium, Amsterdam 1965, pp. 121–131.
12. Elwood W. Greene, *A survey of  $[\alpha, \beta]$ -compactness and related properties*, Master’s thesis, Wesleyan University, 1975.
13. J. E. Vaughan, *Product spaces with compactness-like properties*, *Duke Math. J.*, **39** (1972), 611–617.
14. ———, *Some properties related to  $[\alpha, \beta]$ -compactness*, to appear.
15. ———, *Convergence, closed projections, and compactness*, to appear.

Received June 3, 1974. The author gratefully acknowledges partial support received from the National Science Foundation under grant NSF-GF-39263 A 1.





D. E. Bennett, <i>Strongly unicoherent continua</i> .....	1
Walter R. Bloom, <i>Sets of <math>p</math>-spectral synthesis</i> .....	7
R. T. Bumby and D. E. Dobbs, <i>Amitsur cohomology of quadratic extensions: Formulas and number-theoretic examples</i> .....	21
W. W. Comfort, <i>Compactness-like properties for generalized weak topological sums</i> .....	31
D. R. Dunninger and J. Locker, <i>Monotone operators and nonlinear biharmonic boundary value problems</i> .....	39
T. S. Erickson, W. S. Martindale, 3rd and J. M. Osborn, <i>Prime nonassociative algebras</i> .....	49
P. Fischer, <i>On the inequality <math>\sum_{i=1}^n p_i \frac{f(p_i)}{f(q_i)} \geq 1</math></i> .....	65
G. Fox and P. Morales, <i>Compact subsets of a Tychonoff set</i> .....	75
R. Gilmer and J. F. Hoffmann, <i>A characterization of Prüfer domains in terms of polynomials</i> .....	81
L. C. Glaser, <i>On tame Cantor sets in spheres having the same projection in each direction</i> .....	87
Z. Goseki, <i>On semigroups in which <math>X = XYX = XZX</math> if and only if <math>X = XYZX</math></i> .....	103
E. Grosswald, <i>Rational valued series of exponentials and divisor functions</i> .....	111
D. Handelman, <i>Strongly semiprime rings</i> .....	115
J. N. Henry and D. C. Taylor, <i>The <math>\bar{\beta}</math> topology for <math>w^*</math>-algebras</i> .....	123
M. J. Hodel, <i>Enumeration of weighted <math>p</math>-line arrays</i> .....	141
S. K. Jain and S. Singh, <i>Rings with quasiprojective left ideals</i> .....	169
S. Jeyaratnam, <i>The diophantine equation <math>Y(Y + m)(Y + 2m) \times (Y + 3m) = 2X(X + m)(X + 2m)(X + 3m)</math></i> .....	183
R. Kane, <i>On loop spaces without <math>p</math> torsion</i> .....	189
Alvin J. Kay, <i>Nonlinear integral equations and product integrals</i> .....	203
A. S. Kechris, <i>Countable ordinals and the analytic hierarchy, I</i> .....	223
Ka-Sing Lau, <i>A representation theorem for isometries of <math>C(X, E)</math></i> .....	229
I. Madsen, <i>On the action of the Dyer-Lashof algebra in <math>H_*(G)</math></i> .....	235
R. C. Metzler, <i>Positive linear functions, integration, and Choquet's theorem</i> .....	277
A. Nobile, <i>Some properties of the Nash blowing-up</i> .....	297
G. E. Petersen and G. V. Welland, <i>Plessner's theorem for Riesz conjugates</i> .....	307

Donald Earl Bennett, <i>Strongly uncoherent continua</i> .....	1
Walter Russell Bloom, <i>Sets of <math>p</math>-spectral synthesis</i> .....	7
Richard Thomas Bumby and David Earl Dobbs, <i>Amitsur cohomology of quadratic extensions: formulas and number-theoretic examples</i> .....	21
W. Wistar (William) Comfort, <i>Compactness-like properties for generalized weak topological sums</i> .....	31
Dennis Robert Dunninger and John Stewart Locker, <i>Monotone operators and nonlinear biharmonic boundary value problems</i> .....	39
Theodore Erickson, Wallace Smith Martindale, III and J. Marshall Osborn, <i>Prime nonassociative algebras</i> .....	49
Pál Fischer, <i>On the inequality <math>\sum_{i=0}^n [f(p_i)/f(q_i)]p_i \geq i</math></i> .....	65
Geoffrey Fox and Pedro Morales, <i>Compact subsets of a Tychonoff set</i> .....	75
Robert William Gilmer, Jr. and Joseph F. Hoffmann, <i>A characterization of Prüfer domains in terms of polynomials</i> .....	81
Leslie C. Glaser, <i>On tame Cantor sets in spheres having the same projection in each direction</i> .....	87
Zensiro Goseki, <i>On semigroups in which <math>x = xyx = xzx</math> if and only if <math>x = xyzx</math></i> .....	103
Emil Grosswald, <i>Rational valued series of exponentials and divisor functions</i> .....	111
David E. Handelman, <i>Strongly semiprime rings</i> .....	115
Jackson Neal Henry and Donald Curtis Taylor, <i>The <math>\tilde{\beta}</math> topology for <math>W^*</math>-algebras</i> .....	123
Margaret Jones Hodel, <i>Enumeration of weighted <math>p</math>-line arrays</i> .....	141
Surender Kumar Jain and Surjeet Singh, <i>Rings with quasi-projective left ideals</i> .....	169
S. Jeyaratnam, <i>The Diophantine equation <math>Y(Y + m)(Y + 2m)(Y + 3m) = 2X(X + m)(X + 2m)(X + 3m)</math></i> .....	183
Richard Michael Kane, <i>On loop spaces without <math>p</math> torsion</i> .....	189
Alvin John Kay, <i>Nonlinear integral equations and product integrals</i> .....	203
Alexander S. Kechris, <i>Countable ordinals and the analytical hierarchy. I</i> .....	223
Ka-Sing Lau, <i>A representation theorem for isometries of <math>C(X, E)</math></i> .....	229
Ib Henning Madsen, <i>On the action of the Dyer-Lashof algebra in <math>H_*(G)</math></i> ...	235
Richard C. Metzler, <i>Positive linear functions, integration, and Choquet's theorem</i> .....	277
Augusto Nobile, <i>Some properties of the Nash blowing-up</i> .....	297
Gerald E. Peterson and Grant Welland, <i>Plessner's theorem for Riesz conjugates</i> .....	307