

Pacific Journal of Mathematics

NORMAL HYPERSURFACES

JOSEPH BECKER

NORMAL HYPERSURFACES

JOSEPH BECKER

The purpose of this note is to give a simple analytic proof of a theorem of Oka: If V is a complex analytic hypersurface whose singular locus has codimension at least two, then V is normal. In other words, every weakly holomorphic function is holomorphic.

This result has since been generalized by Abhankar and Thimm to the case when V is an algebraic complete intersection (which is to say that the ideal of functions holomorphic in the ambient space vanishing on V is generated by k functions, where k is the codimension of V in the ambient space).

Actually we prove a slightly stronger result than Oka's.

THEOREM. *Let V be a complex analytic hypersurface, A a complex analytic subset of V with codimension at least 2. Then there is a bounded linear operator $\phi: \mathcal{O}(V - A) \rightarrow \mathcal{O}(V)$ such that $\phi(f) | V - A = f$.*

Proof. Suppose $V \subset \mathbb{C}^n$ and the projection $\pi: \mathbb{C}^n \rightarrow \mathbb{C}^{n-1}$ to the first $n - 1$ co-ordinates gives an r -sheeted branched cover of V in some neighborhood of the origin with branch set B , $B' = \pi(B)$, $A' = \pi(A)$ and $z' = \pi(z)$. Now π induces a homomorphism ${}_{n-1}\mathcal{O} \rightarrow {}_n\mathcal{O} / I(V) = \mathcal{O}(V)$ making $\mathcal{O}(V)$ into a finitely generated ${}_{n-1}\mathcal{O}$ module with generators $1, z_n, \dots, z_n^{r-1}$. Let $P(z', z_n)$ be the minimal degree polynomial for z_n over ${}_{n-1}\mathcal{O}$; for any $f \in \mathcal{O}(V)$ by the Weierstrass division theorem we have $f = QP + R$ where $R \in {}_{n-1}\mathcal{O}[z_n]$ is a holomorphic polynomial of 'degree $\leq r - 1$. Hence f can be written as $\sum_{i=0}^{r-1} b_i(z') z_n^{r-i-1} \bmod I(V)$. However the $b_i(z')$'s are unique.

For every $z' \notin B'$, let $\alpha_1(z'), \dots, \alpha_r(z')$ be the values of z_n on the fiber $\pi^{-1}(z')$ and $f_j = f(z', \alpha_j(z'))$ for $j = 1, \dots, r$. Then

$$f_j = \sum_{i=0}^{r-1} b_i(z') \alpha_j(z')^{r-i-1}.$$

These equations can be viewed as a system of r linear equations in the r unknowns $b_i(z')$ and solved by Cramer's rule:

$$b_i(z') = \frac{\det [1, \alpha_j, \alpha_j^2, \dots, \alpha_j^{r-i-2}, f_j, \alpha_j^{r-i}, \dots, \alpha_j^{r-1}]}{\det [1, \alpha_j, \dots, \alpha_j^{r-1}]}$$

where in both determinants the entries in the j 'th row are indicated. The denominator is the Vandermonde determinant $\Delta(\alpha_1, \dots, \alpha_r)$ and

equals $\prod_{i \neq k} (\alpha_i(z') - \alpha_k(z'))$. It is an easy exercise in linear algebra to verify that the numerator equals

$$\sum_{j=1}^r \sigma_i(\alpha_1, \dots, \hat{\alpha}_j, \dots, \alpha_r) \Delta(\alpha_1, \dots, \hat{\alpha}_j, \dots, \alpha_r) f_j$$

where σ_i is the elementary symmetric polynomial of degree i , and $\hat{\alpha}_j$ means that α_j is to be deleted. Therefore

$$b_i(z') = \sum_{j=1}^r \frac{\sigma_i(\alpha_1(z'), \dots, \hat{\alpha}_j(z'), \dots, \alpha_r(z'))}{\prod_{k \neq j} (\alpha_j(z') - \alpha_k(z'))} f(z', \alpha_j(z')) .$$

Since f is holomorphic so is

$$b_i(z') = \frac{1}{(r-i-1)!} \frac{\partial^{r-i-1} f}{\partial z_n} (z', \alpha_j(z'))$$

and hence the $b_i(z')$ extend holomorphically across B' . (Since f and $f - QP = R$ have the same values on V , it makes no difference which is used in these formulas.)

Now suppose $h \in \mathcal{O}(V - A)$, and for $z' \notin B'$ define

$$\phi(h) = \sum_{j=1}^r \left(\prod_{k \neq j} \frac{z_n - \alpha_k(z')}{\alpha_j(z') - \alpha_k(z')} \right) h(z', \alpha_j(z')) = \sum_{i=0}^{r-1} b_i(z') z_n^{r-i-1}$$

where b_i is defined above. Then the functions are holomorphic in a neighborhood of z' provided h is holomorphic near each point in $\pi^{-1}(z')$. Thus $b_i(z')$ extend to holomorphic functions near any point of $B' - A'$ and then by Hartogs' theorem, b_i extends across A' , because $\text{codim } A \geq 2$.

The operator ϕ whose existence has been demonstrated is obviously linear and clearly bounded with norm one because both $\mathcal{O}(V - A)$ and $\mathcal{O}(V)$ are Fréchet spaces [3] with seminorms being sup on compact subsets; $V - A$ is dense in V so $\sup_{x \in V} |\phi(f)| = \sup_{V \in x-A} |f(x)|$.

Incidentally notice that we did not use the result [4] that a function holomorphic in the complement of an analytic set of codim two is bounded in a neighborhood of that set, e.g. $\mathcal{O}(V - A) = \mathcal{O}(V)$.

REMARK. The same proof works when $\mathcal{O}(V)$ is a Cohen Macaulay ring, that is the direct image sheaf of $\mathcal{O}(V)$ by a local parametrization is a free sheaf.

REFERENCES

1. S. Abhyankar, *Concepts of order and rank on a complex space, and a condition for normality*, Math. Ann., **141**, (1960), 171-192.

2. R. C. Gunning, *Lectures on Complex Analytic Varieties*, Mathematical Notes, Princeton, N.J., Princeton University Press 1970.
3. R. C. Gunning and H. Rossi, *Analytic Functions of Several Complex Variables*, Englewood Cliffs, N.J., Prentice-Hall 1965.
4. R. Narasimhan, *Introduction to the Theory of Analytic Spaces*, Lecture Notes in Mathematics 25, Berlin-Heidelberg-New York, Springer 1966.
5. K. Oka, *Sur les fonctions analytique de plusieurs variables VIII*, J. Math. Soc. Japan, **3** (1951).
6. W. Thimm, *Untersuchungen über das Spurproblem von holomorphen Funktionen auf analytischen Mengen*, **139** (1959), 95-114.

Received February 5, 1974.

PURDUE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics

Vol. 61, No. 1

November, 1975

Jiří Adámek, V. Koubek and Věra Trnková, <i>Sums of Boolean spaces represent every group</i>	1
Richard Neal Ball, <i>Full convex l-subgroups and the existence of a^*-closures of lattice ordered groups</i>	7
Joseph Becker, <i>Normal hypersurfaces</i>	17
Gerald A. Beer, <i>Starshaped sets and the Hausdorff metric</i>	21
Dennis Dale Berkey and Alan Cecil Lazer, <i>Linear differential systems with measurable coefficients</i>	29
Harald Boehme, <i>Glättungen von Abbildungen 3-dimensionaler Mannigfaltigkeiten</i>	45
Stephen LaVern Campbell, <i>Linear operators for which T^*T and $T + T^*$ commute</i>	53
H. P. Dikshit and Arun Kumar, <i>Absolute summability of Fourier series with factors</i>	59
Andrew George Earnest and John Sollion Hsia, <i>Spinor norms of local integral rotations. II</i>	71
Erik Maurice Ellentuck, <i>Semigroups, Horn sentences and isolic structures</i>	87
Ingrid Fotino, <i>Generalized convolution ring of arithmetic functions</i>	103
Michael Randy Gabel, <i>Lower bounds on the stable range of polynomial rings</i>	117
Fergus John Gaines, <i>Kato-Taussky-Wielandt commutator relations and characteristic curves</i>	121
Theodore William Gamelin, <i>The polynomial hulls of certain subsets of C^2</i>	129
R. J. Gazik and Darrell Conley Kent, <i>Coarse uniform convergence spaces</i>	143
Paul R. Goodey, <i>A note on starshaped sets</i>	151
Eloise A. Hamann, <i>On power-invariance</i>	153
M. Jayachandran and M. Rajagopalan, <i>Scattered compactification for $N \cup \{P\}$</i>	161
V. Karunakaran, <i>Certain classes of regular univalent functions</i>	173
John Cronan Kieffer, <i>A ratio limit theorem for a strongly subadditive set function in a locally compact amenable group</i>	183
Siu Kwong Lo and Harald G. Niederreiter, <i>Banach-Buck measure, density, and uniform distribution in rings of algebraic integers</i>	191
Harold W. Martin, <i>Contractibility of topological spaces onto metric spaces</i>	209
Harold W. Martin, <i>Local connectedness in developable spaces</i>	219
A. Meir and John W. Moon, <i>Relations between packing and covering numbers of a tree</i>	225
Hiroshi Mori, <i>Notes on stable currents</i>	235
Donald J. Newman and I. J. Schoenberg, <i>Splines and the logarithmic function</i>	241
M. Ann Piech, <i>Locality of the number of particles operator</i>	259
Fred Richman, <i>The constructive theory of KT-modules</i>	263
Gerard Sierksma, <i>Carathéodory and Helly-numbers of convex-product-structures</i>	275
Raymond Earl Smithson, <i>Subcontinuity for multifunctions</i>	283
Gary Roy Spoor, <i>Differentiability conditions and bounds on singular points</i>	289
Rosario Strano, <i>Azumaya algebras over Hensel rings</i>	295