

# Pacific Journal of Mathematics

**THE WORD PROBLEM AND POWER PROBLEM IN  
1-RELATOR GROUPS ARE PRIMITIVE RECURSIVE**

FRANK BENJAMIN CANNONITO AND RONALD WALLACE GATTERDAM

## THE WORD PROBLEM AND POWER PROBLEM IN 1-RELATOR GROUPS ARE PRIMITIVE RECURSIVE

F. B. CANNONITO AND R. W. GATTERDAM

**The purpose of this paper is to show the solution to the word problem in a 1-relator group can be computed with respect to an effective indexing of the group by an algorithm at level at most  $2+\sigma(R)$  of the Grzegorzcyk hierarchy, where  $\sigma(R)$  is the length of the relator, and by a primitive recursive function, always. As a consequence, it is shown that the power problem in a 1-relator group can be solved similarly. An example is given in which the Magnus algorithm for the extended word problem is at level 4 but not 3 of the Grzegorzcyk hierarchy even though the word problem is solvable at level 3.**

Our theorem provides a negative solution to problem 20 of [2], p. 643, which reads "is there a one-relator group whose word problem is not solvable by a primitive recursive function?" These results do not depend on a particular presentation of the group, rather they are algebraic invariants in the sense of [3]. (The algebraic invariance of the order and power problem can be shown in a manner similar to the proof of the invariance of the word problem in [3].) Since each finitely generated subgroup of a group with primitive recursive word problem has a primitive recursive word problem ([4] Corollary 3.7, p. 380) and there are finitely generated and finitely presented groups with recursive but not primitive recursive word problem ([5], [6] and [1] we see that a finitely generated group with solvable word problem cannot necessarily be embedded in 1-relator group. (This result can be obtained in the stronger form: a finitely presented group with primitive recursive word problem is not necessarily embeddable in a 1-relator group, from B. B. Newman's work [11] which shows a free abelian group of rank 3 cannot be a subgroup of a 1-relator group and the easily shown primitive recursive solution to the word problem in such groups.)

The proof of the main theorem follows by analyzing the proof of Magnus' theorem of the solvability of the word problem in 1-relator groups as given in [9], with respect to the Grzegorzcyk hierarchy of primitive recursive functions [8]. (See also [12].) We assume a knowledge of [4], [5] and [9]. We prove that the Magnus algorithm for solving the "extended word problem" and hence the word problem in a one relator group is  $\mathcal{E}^{2+\sigma(R)}$  computable in the Grzegorzcyk hierarchy where  $\sigma(R)$  is the length of the relator. In

section 4 we verify that the order and power problem in one-relator groups are also  $\mathcal{E}^{2+\sigma(R)}$  computable.

Finally we note that the Magnus algorithm is not always the "easiest" algorithm in the sense of the Grzegorzcyk hierarchy, for solving the word problem in a one relator group. In §5 we give an example of a one relator group in which the Magnus algorithm is  $\mathcal{E}^4$  but not  $\mathcal{E}^3$  computable and yet for which the word problem is  $\mathcal{E}^3$  computable.

2. Notation and definitions. We abbreviate finitely generated, finitely presented, word problem, primitive recursive by "f.g.", "f.p.", "w.p." and "p.r." respectively. The levels of the Grzegorzcyk hierarchy are denoted by  $\mathcal{E}^\alpha$ ,  $\alpha \geq 0$ . If  $R$  is a word on  $a, b, c, \dots$  then  $\sigma(R)$  denotes the length of  $R$  and  $\sigma_a(R)$  denotes the exponent sum of  $a$  in  $R$ .

The following definition is basic to this paper. It is discussed in detail in [4] and [5].

A countable group  $G$  is  $\mathcal{E}^\alpha$  computable, or briefly an  $\mathcal{E}^\alpha$  group if the following three conditions are met:

(1) There is an injection  $i: G \rightarrow \omega$  such that  $i(G)$  is an  $\mathcal{E}^\alpha$ -decidable subset of  $\omega$ . (If  $r \in i(G)$ ,  $i^{-1}(r)$  will be denoted by  $g_r$ .)

(2) The function  $m: i(G) \times i(G) \rightarrow i(G)$  defined by  $m(r, s) = i(g_r \cdot g_s)$ ,  $r, s \in i(G)$  is an  $\mathcal{E}^\alpha$  function. (More precisely the restriction of an  $\mathcal{E}^\alpha$  function to  $i(G) \times i(G)$ .)

(3) The function  $\text{In}: i(G) \rightarrow i(G)$  defined by

$$\text{In}(r) = i(g_r^{-1}), r \in i(G)$$

is an  $\mathcal{E}^\alpha$  function. (This condition is redundant for  $\mathcal{E}^\alpha =$  the recursive functions.)

The function  $i$  of (1) is called an  $\mathcal{E}^\alpha$  index of  $G$ . Of particular importance is the notion of a *standard index* which is obtained as follows. Let  $G$  have a presentation  $G = gp(a_1, a_2, a_3, \dots; R, S, T, \dots)$  where  $R, S, T, \dots$  are words on  $a_1, a_2, a_3, \dots$ . We fix a Gödel numbering of the free group  $F$  on  $a_1, a_2, a_3, \dots$  and assign to each  $g \in G$  the index  $i(g)$  equal to the smallest index in  $\rho^{-1}(g)$  where  $\rho$  is the canonical epimorphism  $F \rightarrow G$ . If  $G$  is  $\mathcal{E}^\alpha$  computable with respect to a f.g. standard index we say  $G$  is  $\mathcal{E}^\alpha$  standard. It is known, in this case, that  $G$  is then  $\mathcal{E}^\alpha$  standard with respect to all f.g. standard indices for this is another way of expressing the algebraic invariance of the w.p. for f.g. groups with respect to the Grzegorzcyk hierarchy. See [3] and [6] for the details on the equivalence of  $\mathcal{E}^\alpha$  w.p. and having an  $\mathcal{E}^\alpha$  standard index for f.g. groups.

3. The main result. We now state and prove the

**THEOREM.** *The word problem in a one relator group*

$$G = gp(a, b, \dots; R(a, b, \dots))$$

*is at most  $\mathcal{E}^{2+\sigma(R)}$  solvable. In particular it is p.r.*

*Proof.* We prove the “extended word problem” (henceforth “e.w.p.”) for  $G$  is  $\mathcal{E}^{2+\sigma(R)}$  where we mean the problem of determining if a freely reduced word on the generators of  $G$  represents an element of the subgroup  $H < G$  generated by a given proper subset of the generators and, if so, to rewrite it in terms of the generators of  $H$ . The proof will involve an induction on  $\sigma(R)$  which raises the degree of computational complexity by at most 1 at each stage.

It should be observed that we may assume  $R$  involves all of the generators of  $G$ , and in particular that  $G$  is f.g., for otherwise we can write  $G$  as the free product of a free group with a 1-relator group in which the relator involves all of the generators. Such a free product has an  $\mathcal{E}^n$  e.w.p. if the 1-relator factor does, by virtue of the normal form theorem for free products and the obvious  $\mathcal{E}^3$  e.w.p. for free groups. Also, observe that by the Freiheitssatz the subgroup of  $G$  generated by deleting one relator is free. Thus, if  $H < G$  is a subgroup generated by a proper subset of the generators, say  $a \notin H$ , and  $W$  is an arbitrary word, the e.w.p. for  $W$  reduces to the question of  $W$  representing a word in  $K = gp(b, c, \dots) < G$  and of rewriting  $W$  in terms of  $b, c, \dots$ , because the e.w.p. in  $K$  (with respect to generators  $b, c, \dots$ ) is  $\mathcal{E}^3$ . This last remark holds in particular if  $H = \{1\}$ , so an  $\mathcal{E}^n$  solution to the e.w.p. implies an  $\mathcal{E}^n$  solution to the w.p.

Now if  $G$  has one generator,  $G$  is either trivial or finite cyclic and the e.w.p. is identical to the w.p. and is  $\mathcal{E}^3$ . Thus, we assume  $G$  has at least two generators, say  $a$  and  $b$ , and hence that  $R$  has length at least two. The proof proceeds by recursion on the length of the relator.

Given a subgroup  $H < G$  generated by all but one of the generators of  $G$  and a word  $W$  representing an element of  $G$  the recursion reduces the question of  $W$  representing an element of  $H$  in  $G$  (we write  $W \in H < G$ ) to the question  $W' \in H' < G'$ , where  $G'$  is a 1-relator group with relator of length less than that of  $R$  and  $H'$  is generated by a proper subset of the generators of  $G'$ . We show at each stage of this recursion the encoding of  $W'$  is  $\mathcal{E}^3$  computable from the result of the previous stage and that at

each stage the level of computability is raised by at most 1, involving an (inner) recursion. The result then follows since the number of steps in the recursion is bounded by  $\sigma(R)$  independent of  $\sigma(W)$ .

First, assume  $\sigma_a(R) = 0$ . Also assume  $H < G$ , is generated by all generators of  $G$  excluding  $a$ . Let  $N < G$  be the normalizer of the set of generators of  $G$  excluding  $a$ . Then since  $\sigma_a(R) = 0$ , the map  $G \rightarrow gp(a;)$  by  $a \mapsto a$  and all other generators  $\mapsto 1$  extends to an epimorphism  $G \rightarrow gp(a;)$  given by  $W(a, b, \dots) \mapsto a^{\sigma_a(W(a,b,\dots))}$ . Thus,  $W \in N$  iff  $\sigma_a(W) = 0$  (an  $\mathcal{E}^3$  decision) and since  $H < N$ ,  $W \notin H$  in  $G$  otherwise. By a Reidemeister-Schreier rewriting process (using coset representatives  $a^k$ )  $N$  has a presentation

$$N = gp(\dots, b_{-1}, b_0, b_1, \dots, c_{-1}, c_0, c_1, \dots; \dots, P_{-1}, P_0, P_1, \dots)$$
 where

$b_k = a^k b a^{-k}$ ,  $c_k = a^k c a^{-k}$ , etc. and  $P_k$  is  $a^k R a^{-k} \in N$  rewritten in terms of the  $b_k$ ,  $c_k$ , etc. For  $W \in N$ ,  $W$  is rewritten by replacing  $b^j$  (or  $c^j$  etc.) by  $b_k^j$  for  $k$  equal to the exponent sum on  $a$  of the symbols preceding  $b_k^j$ . (For example  $a^2 b^4 a c^{-1} a^{-1}$  is rewritten  $b_2^4 c_3^{-1}$ .) In particular the rewriting lowers the length of  $W$ . Notice also that  $P_k$  may be obtained from  $P_0$  by raising the subscripts by  $k$ .

Now for  $W \in N$ ,  $W \in H$  iff its rewrite  $W' \in H'$  for  $H' = gp(b_0, c_0, \dots) < N$ . Note  $\sigma(W') \leq \sigma(W)$ . For simplicity of notation, assume  $R$  begins with the symbol  $b$  (conjugate  $R$  if necessary) and let  $n$  be the minimum subscript of  $b$  in  $P_0$  ( $n \leq 0$  by the simplification) and  $m$  the maximum subscript of  $b$  in  $P_0$  ( $m \geq 0$ ). Set

$$N_0 = gp(b_n, \dots, b_m, \dots, c_{-1}, c_0, c_1, \dots; P_0).$$

Note  $H' < N_0$  since  $n \leq 0 \leq m$ . Recursively form

$$N_k = gp(b_{n+k}, \dots, b_{m+k}, \dots, c_{-1}, c_0, c_1, \dots; P_k).$$

It should be observed that the  $N_k$  are all isomorphic to  $N_0$  the isomorphism given by lowering the subscripts of the  $b$  generators by  $k$ . Also note the group  $N_0$  (and subsequent groups in the recursion playing the role of  $N_0$ ) are determined by  $R$  independent of  $W$ . Thus (formal) questions and rewritings of words in  $N_k$  may be treated as questions and rewritings in  $N_0$  by an  $\mathcal{E}^3$  modification of subscripts. The significance of the above is that in the following the groups  $N_{-\sigma(W)}$ ,  $N_{-\sigma(W)+1}$ ,  $\dots$ ,  $N_0$ ,  $\dots$ ,  $N_{\sigma(W)}$  must be considered, the number dependent on  $W$ . Nevertheless, the algorithms involved depend only on  $R$ .

Form  $N_{0,1} = N_0 *_{\phi} N_1$  where  $\phi$  amalgamates the free subgroup generated by  $b_{n+1}, \dots, b_m, \dots, c_{-1}, c_0, c_1, \dots$  (all but  $b_n$ ) in  $N_0$  with the free subgroup having the same generators in  $N_1$ . Next, form  $N_{-1,1} = N_0 *_{\phi} N_{-1}$  for  $\phi$  amalgamating the free subgroups generated

by  $b_n, \dots, b_{m-1}, \dots, c_{-1}, c_0, c_1, \dots$  and proceed to form  $N_0 < N_{0,1} < N_{-1,1} < N_{-1,2} < N_{-2,2} < \dots$ . Now  $N$  is the union of this chain so if  $W \in N$ ,  $W$  is eventually in one of these groups (at least by stage  $N_{-\sigma(W), \sigma(W)}$ ). Note  $H' < N_0$ . Also observe that up to isomorphism by change of subscripts of the  $b$  symbols, there are two types of amalgams namely one generated by  $b_{n+k}, \dots, b_{m+k-1}, \dots, c_{-1}, c_0, c_1, \dots$ , and one generated by  $b_{n+k+1}, \dots, b_{m+k}, \dots, c_{-1}, c_0, c_1, \dots$ . Thus questions and rewrites in the amalgams do not depend on  $k$  and hence not on  $W$  up to an  $\mathcal{E}^3$  process on the subscripts.

We must show both the question  $W' \in H'$  and, if true, the process of rewriting  $W'$  in terms of  $b_0, c_0, \dots$  are  $\mathcal{E}^{2+\sigma(R)}$ . Then the rewriting in terms of  $b, c, \dots$  (i.e. renumbering to delete the subscripts) is  $\mathcal{E}^3$  and so the e.w.p. is  $\mathcal{E}^{2+\sigma(R)}$  in this case.

By an  $\mathcal{E}^3$  process,  $W' \equiv W_1 \dots W_p$  in syllables, each  $W_i$  in a different  $N_k$ . We describe a recursion involving the e.w.p. in  $N_k$  (or in  $N_0$ ). Inductively (recursively) the e.w.p. in  $N_k$  is  $\mathcal{E}^{2+\sigma(P_0)}$  and  $\sigma(P_0) < \sigma(R)$  so the computability of the (inner) recursion is at most  $\mathcal{E}^{2+\sigma(P_0)+1} \subseteq \mathcal{E}^{2+\sigma(R)}$ . We consider each pair of syllables  $W_i W_{i+1}$ . Let  $W_i \in N_k$  and  $W_{i+1} \in N_l$  and assume  $k < l$  (for simplicity; the case  $k > l$  is similar). Then we ask: can  $W_i$  be rewritten without the  $b_{n+k}$  symbol? and if so rewrite it (an  $\mathcal{E}^{2+\sigma(P_0)}$  process). Next, we ask: can  $W_{i+1}$  be rewritten without the  $b_{m+k-1}$  symbol? and if so rewrite it. The rewrites are in  $N_{k+1}$  and  $N_{l-1}$  if possible. We continue until neither type of rewrite is possible or  $W_i$  has been rewritten  $W'_i \in N_j$  and  $W_{i+1}$  rewritten  $W'_{i+1} \in N_j$  for  $k \leq j \leq l$ . In the former event,  $W_i W_{i+1}$  is left as in the original; in the latter  $W_i W_{i+1}$  is replaced by the single syllable  $(W'_i W'_{i+1})$  reducing the syllable length. The process continues until  $W'$  has been rewritten  $W''$  with minimal syllable length. Now the possibilities for  $W''$  are:

(i)  $W'' = \Lambda$  (the empty word) whence  $W \in H$  with rewrite  $\Lambda$ ,  
(ii) the syllable length of  $W''$  is greater than 1 whence  $W \notin H$ ,  
(iii) the syllable length of  $W''$  is 1 with  $W'' \in N_k$  in which case we ask if  $W''$  can be rewritten  $W'''$  involving  $b_0$  as the only  $b$  symbol if  $n+k \leq 0 \leq m+k$  or no  $b$  symbol otherwise (an  $\mathcal{E}^{2+\sigma(P_0)}$  decision and process) so either

(iii) (a)  $W'''$  can not be obtained hence  $W \notin H$

(iii) (b)  $W''' \in N_0$  can be obtained but  $W''' \notin H'$  (an  $\mathcal{E}^{2+\sigma(P_0)}$  decision) so  $W \notin H$  or

(iii) (c)  $W''' \in N_0$  can be obtained and  $W''' \in H'$  so  $W \in H$  and  $W$  is rewritten in terms of  $b, c, \dots$  by rewriting  $W'''$  in terms of  $b_0, c_0, \dots$  and deleting the subscripts.

Recall that we had assumed above that  $\sigma_a(R) = 0$  and  $a \notin H$ . Now assume  $\sigma_a(R) = 0$  but  $b \notin H$  (so  $a \in H$ ). In this case  $W \in H$

does not imply  $W \in N$  but for  $\alpha = \sigma_a(W)$ ,  $W \in H$  iff  $Wa^{-\alpha} \in H$  and  $Wa^{-\alpha} \in N$ . Thus the above argument is carried out replacing  $W$  by  $Wa^{-\alpha}$  and using  $H'$ , the subgroup generated by all generators of  $N$  excluding the  $b_k$  for all  $k$ . It should be observed that  $H' < N_0$  and also that the rewrite  $W'$  of  $Wa^{-\alpha}$  satisfies  $\sigma(W') \leq \sigma(W)$ . Step (iii) c) must be replaced by  $W''' \in H$  generated by  $\dots c_{-1}, c_0, c_1, \dots, d_{-1}, d_0, \dots$  and  $W$  is rewritten in terms of  $a, c, \dots$  by rewriting  $W'''$  replacing  $c_i$  (respectively  $d_i$  etc.) by  $a^i c a^{-i}$  ( $d^i c d^{-i}$  respectively), concatenating  $a^\alpha$  and freely reducing (all  $\mathcal{E}^3$  processes).

Next, assume none of the exponent sums of  $R$  is zero and the problem is to determine if  $W(a, b, c, \dots)$  is equal in  $G$  to a word not involving  $b$  and if so rewrite  $W$  without  $b$ . Let  $\sigma_a(R) = \alpha \neq 0$  and  $\sigma_b(R) = \beta \neq 0$ . Consider  $E = gp(\eta, b, c, \dots; R(\eta^\beta, b, \dots)) = G *_{\phi} gp(\eta;)$  where  $\phi$  amalgamates the infinite cyclic groups generated by  $a$  and  $\eta^\beta$ . By an  $\mathcal{E}^3$  process  $W(a, b, \dots) \in G$  can be rewritten  $W(\eta^\beta, b, \dots) \in E$ . By a Tietze transformation

$$\begin{aligned} E &= gp(\eta, \zeta, b, c, \dots; R(\eta^\beta, b, \dots), b = \zeta\eta^{-\alpha}) \\ &= gp(\eta, \zeta, c, \dots; R(\eta^\beta, \zeta\eta^{-\alpha}, \dots)) \end{aligned}$$

where  $\sigma_\eta(R(\eta^\beta, \zeta\eta^{-\alpha}, \dots)) = 0$ . Now  $W(a, b, c, \dots)$  is equal in  $G$  to a word involving only  $a, c, \dots$  iff  $W(\eta^\beta, \zeta\eta^{-\alpha}, \dots)$  is equal in  $E$  to a word involving only  $\eta^\beta, c, \dots$ . Note that expressing  $W(\eta^\beta, \zeta\eta^{-\alpha}, \dots)$  in terms of  $W(a, b, c, \dots)$  is an  $\mathcal{E}^3$  process. We proceed with the previous analysis letting  $N$  be the normalizer of  $\{\eta, c, \dots\}$  (excluding  $\zeta$ ) in  $E$ . Thus, by the recursion we can determine if  $W(\eta^\beta, \zeta\eta^{-\alpha}, c, \dots)$  is equal in  $E$  to a word say  $V(\eta, c, \dots)$  (excluding  $\zeta$  symbols) and if so compute  $V$  by an  $\mathcal{E}^{2+\sigma(R)}$  process. Now since the subgroup of  $E$  generated by  $\eta, c, \dots$  (excluding  $\zeta$ ) is free, it is an  $\mathcal{E}^3$  process to determine if all powers of  $\eta$  in  $V$  are multiples of  $\beta$ . If so, replacing  $\eta^\beta$  by  $a$  in  $V$  is also an  $\mathcal{E}^3$  process. Thus, this situation has been reduced to the previous one. This completes the proof.

4. The power problem. We now turn to the power problem in 1-relator groups. The reader will recall that the generalized word problem (g.w.p.) for a subgroup  $H < G = gp(S; D)$  is the algorithmic problem of deciding whether or not an arbitrary word  $W \in G$  defines an element of  $H$ . If the g.w.p. is solvable for every cyclic subgroup of  $G$  then  $G$  is said to have solvable power problem. The object of this section is to prove that the power problem is  $\mathcal{E}^{2+\sigma(R)}$ -decidable in 1-relator groups. We followed the discussion in McCool [10], indicating where the argument McCool must be modified to obtain the information needed to locate the decidability level with respect to the hierarchy  $\{\mathcal{E}^\alpha\}$ .

The first step after noting that McCool's lemma [10, p. 428] is identical to parts of our theorem above, is to show the order problem is  $\mathcal{E}^{2+\sigma(R)}$ -decidable. That is, the algorithmic problem of deciding the order of the element defined by a word  $W$  in a 1-relator group is  $\mathcal{E}^{2+\sigma(R)}$ . We follow the discussion in [9] Theorem 4.13, page 269. Now a 1-relator group has an element of finite order if and only if the relator  $R$  is a  $k^{\text{th}}$  power,  $k > 1$ , of some non-empty word  $V$  in the free group on the generators of the 1-relator group). Thus we assume  $G = gp(a_1, \dots, a_n; V^k(a_1, \dots, a_n))$ ,  $k > 1$ . Furthermore, we may assume  $V$  is cyclically reduced since we assume  $V^k$  is. (The reduction of a word  $W$  to a cyclically reduced word can be accomplished by an  $\mathcal{E}^3$  process because a word is cyclically reduced if and only if each of its cyclic permutations is freely reduced. Clearly all cyclic permutations and their subsequent free reductions are obtainable from the index of  $W$  by an  $\mathcal{E}^3$  process.) Thus given  $V^k$ , since  $k \leq \sigma(V^k)$  and  $\sigma(V) \leq \sigma(V^k)$  we can use bounded minimalization, which preserves level  $\mathcal{E}^3$ , to obtain  $V$  and  $k$  by an  $\mathcal{E}^3$  process. Then  $W \in G$  has finite order ( $\leq k$ ) if and only if  $W$  is conjugate to a power of  $V$  ([9] Theorem 4.13, p. 269). Therefore  $W$  has finite order if and only if  $W^k = 1$ , and  $\mathcal{E}^{2+\sigma(R)}$  decision, since the word problem in  $G$  is  $\mathcal{E}^{2+\sigma(R)}$ . If the order of  $W$  is finite, it is the minimal  $m \leq k$  such that  $W^m = 1$  and so is  $\mathcal{E}^{2+\sigma(R)}$  computable by a bounded minimalization on the decision procedure for the word problem. The remainder of Case 1 of McCool [10], page 428 can be decided at level  $\mathcal{E}^{2+\sigma(R)}$ , since the induction used is identical to the recursion given above for the  $\mathcal{E}^{2+\sigma(R)}$ -decidability of the e.w.p. As above, Case 2 reduces to Case 1. Thus we have the

**THEOREM.** *The order and power problems in a 1-relator group are  $\mathcal{E}^{2+\sigma(R)}$ -decidable. In particular, they are p.r.*

5. Remarks, example and questions. We remark again that the groups considered in the Magnus process are determined, up to isomorphism by relabelling subscripts, only by the relator and not by the word under consideration. The induction on the length of the relator introduces at each stage two new groups, the amalgams. Thus the number of groups which must be considered is  $\mathcal{E}^{\alpha(R)+1} - 1$  where  $\alpha(R)$  is the number of stages in the induction and depends only on the relator  $R$ . It is also clear that a better bound on the computability level is  $\mathcal{E}^{3+\alpha(R)}$  where  $\alpha(R) \leq \sigma(R)-1$ .

In fact each stage of the recursion need not raise the computability level since in many cases the length of the word currently under consideration is reduced permitting a bound on the recursion. However every application of the case type  $\sigma_{\text{generator}}(\text{Relator}) \neq 0$  in



the course of the recursion can increase the length of the rewriting. This being the only instance in which the length of the rewrite of  $W$  can increase, a better bound than  $\mathcal{E}^{3+\alpha(R)}$  can be obtained for particular groups.

In view of the above, the authors had long erroneously conjectured that the computability bound  $\mathcal{E}^3$  could be obtained for the Magnus process on any one relator group. The following example which shows the conjecture false has been brought to their attention. Let  $G = gp(a, b; a^{-1}b^{-1}aba^{-1}bab^{-2})$ ,  $X_0 = b$ ,  $X_{k+1} = a^{-1}X_k^{-1}aba^{-1}X_k a$ . Then  $\sigma(X_k) = 2^k + 5(2^k - 1)$ , an  $\mathcal{E}^3$  function of  $k$  and  $X_{\frac{k}{K}} \stackrel{\text{def}}{=} b^{f_4(2^k, k)}$  for  $f_4$  as defined in [4], an  $\mathcal{E}^4 - \mathcal{E}^3$  function of  $k$ . This shows the Magnus process for  $G$  has a lower computability bound of  $\mathcal{E}^4$ . In fact a careful consideration of the recursion in the Magnus process for  $G$  shows one occurrence of the  $\sigma_{\text{generator}}(\text{Relator}) \neq 0$  case and exactly the  $\mathcal{E}^4 - \mathcal{E}^3$  computability of the Magnus algorithm.

It is tempting to conjecture that the group  $G$  in the above example has  $\mathcal{E}^4 - \mathcal{E}^3$  computable w.p. To see however that  $G$  has an  $\mathcal{E}^3$  computable w.p. first observe  $G \cong gp(a, b, c; c^{-1}bc = b^2, a^{-1}ba = c)$  by a Tietze transformation. Let  $F = gp(b;)$  and  $\phi_1: F \rightarrow F$  by  $b \mapsto b^2$  (an  $\mathcal{E}^3$  isomorphism in  $F$  as defined in [7]). Then by Tietze transformations the strong Britton extension  $F_{\phi_1} = gp(b, c; c^{-1}bc = b^2) < gp(b, c;)$   $*_{\psi_1} gp(\hat{b}, s;)$   $= M$  where  $\psi_1$  amalgamates  $H$ , the subgroup of  $gp(\hat{b}, c;)$  generated (freely) by  $\hat{b}$  and  $c^{-1}bc$  with the subgroup  $\hat{H}$  of  $gp(\hat{b}, s;)$  generated (freely) by  $\hat{b}$  and  $s^{-1}\hat{b}^2s$  (see [7] proof of Theorem 3.1). We show  $M$  has  $\mathcal{E}^3$  computable w.p. and hence so does  $F_{\phi_1}$ . To see the w.p. in  $M$  is  $\mathcal{E}^3$  let  $W$  be a word on  $b, c, \hat{b}, s$ . Then following the proof of Theorem 4.10 of [4],  $W = W_1 W_2 \dots W_p$  in syllables and the w.p. for  $M$  is solved by rewriting the syllables to obtain a word of shorter syllable length. Since the rewriting are  $\hat{b} \leftrightarrow b$  and  $c^{-1}bc \leftrightarrow s^{-1}\hat{b}^2s$ , the length of a rewrite is at most twice the length of the original and since each rewrite lowers the syllable length, the length of any rewrite is bounded by  $2^p \sigma(W) \leq 2^{\sigma(W)} \sigma(W)$ , an  $\mathcal{E}^3$  function of  $W$ . Thus the recursion has an  $\mathcal{E}^3$  bound and the w.p. in  $M$  is  $\mathcal{E}^3$ . Similarly,  $\phi_2$  given by  $\phi_2: b \mapsto c$  is an  $\mathcal{E}^3$  isomorphism in  $F_{\phi_1}$  and hence in  $M$  so  $F_{\phi_1, \phi_2} < M_{\phi_2}$  a subgroup of a free product with amalgamation where in this case the rewrites do not increase the length of the word under consideration. Thus  $M_{\phi_2}$  hence  $F_{\phi_1, \phi_2} = gp(a, b, c; c^{-1}bc = b^2, a^{-1}ba = c) \cong G$  has  $\mathcal{E}^3$  w.p.

Thus the question of whether or not the w.p. for one relator groups is  $\mathcal{E}^3$  remains open. One might also ask if the example given can be generalized to obtain one relator groups with strictly  $\mathcal{E}^n$  Magnus process and  $\mathcal{E}^3$  w.p.

The authors would like to thank Charles F. Miller III for his constructive advice.

## REFERENCES

1. W. W. Boone, *Word problems and recursively enumerable degrees of unsolvability. A sequel on finitely presented groups*, Ann. of Math., **84** (1966), 49-84.
2. W. W. Boone, F. B. Cannonito and R. C. Lyndon, editors, *Word Problems*, North-Holland Pub. Co., Amsterdam, 1973, 646 pp.
3. F. B. Cannonito, *The algebraic invariance of the word problem in groups*, in Word Problems, op. cit., 349-364.
4. F. B. Cannonito and R. W. Gatterdam, *The computability of group construction*, Part I, in Word Problems, op. cit., 365-400.
5. F. B. Cannonito, *Hierarchies of computable groups and the word problem*, J. Symbolic Logic, **31**, No. **3** (1966), 376-392.
6. R. W. Gatterdam, *The computability of group constructions, II*, Bull. Australian Math. Soc., **8**, No. **1** (1973), 27-60.
7. ———, *Embeddings of primitive recursive computable groups*, doctoral dissertation, University of California, Irvine 1970.
8. A. Grzegorzcyk, *Some classes of recursive functions*, Rozprawy Mat', Warsaw, **4** (1953), 1-45.
9. W. Magnus, A. Karass and D. Solitar, *Combinatorial Group Theory*, Interscience Pub., New York, 1966.
10. J. McCool, *The power problem for groups with one defining relator*, Proc. Amer. Math. Soc., **28**, No. **2** (1971), 427-430.
11. B. B. Newman, *Some results on one-relator groups*, Bull. Amer. Math. Soc., **74**, No. **3** (1968), 568-571.
12. A. Yasuhara, *Recursive Function Theory and Logic*, Academic Press, New York, 1971.

Received July 5, 1974 and in revised form August 5, 1975. Air Force Office of Scientific Research Grant 1321-67. Wisconsin Alumni Research Fund.

UNIVERSITY OF CALIFORNIA, IRVINE  
AND  
UNIVERSITY OF WISCONSIN, PARKSIDE



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

Graham Donald Allen, Francis Joseph Narcowich and James Patrick Williams, <i>An operator version of a theorem of Kolmogorov</i> .....	305
Joel Hilary Anderson and Ciprian Foias, <i>Properties which normal operators share with normal derivations and related operators</i> .....	313
Constantin Gelu Apostol and Norberto Salinas, <i>Nilpotent approximations and quasinilpotent operators</i> .....	327
James M. Briggs, Jr., <i>Finitely generated ideals in regular <math>F</math>-algebras</i> .....	339
Frank Benjamin Cannonito and Ronald Wallace Gatterdam, <i>The word problem and power problem in 1-relator groups are primitive recursive</i> .....	351
Clifton Earle Corzatt, <i>Permutation polynomials over the rational numbers</i> .....	361
L. S. Dube, <i>An inversion of the <math>S_2</math> transform for generalized functions</i> .....	383
William Richard Emerson, <i>Averaging strongly subadditive set functions in unimodular amenable groups. I</i> .....	391
Barry J. Gardner, <i>Semi-simple radical classes of algebras and attainability of identities</i> .....	401
Irving Leonard Glicksberg, <i>Removable discontinuities of <math>A</math>-holomorphic functions</i> ...	417
Fred Halpern, <i>Transfer theorems for topological structures</i> .....	427
H. B. Hamilton, T. E. Nordahl and Takayuki Tamura, <i>Commutative cancellative semigroups without idempotents</i> .....	441
Melvin Hochster, <i>An obstruction to lifting cyclic modules</i> .....	457
Alistair H. Lachlan, <i>Theories with a finite number of models in an uncountable power are categorical</i> .....	465
Kjeld Laursen, <i>Continuity of linear maps from <math>C^*</math>-algebras</i> .....	483
Tsai Sheng Liu, <i>Oscillation of even order differential equations with deviating arguments</i> .....	493
Jorge Martinez, <i>Doubling chains, singular elements and hyper-<math>Z</math> <math>l</math>-groups</i> .....	503
Mehdi Radjabalipour and Heydar Radjavi, <i>On the geometry of numerical ranges</i> .....	507
Thomas I. Seidman, <i>The solution of singular equations, I. Linear equations in Hilbert space</i> .....	513
R. James Tomkins, <i>Properties of martingale-like sequences</i> .....	521
Alfons Van Daele, <i>A Radon Nikodým theorem for weights on von Neumann algebras</i> .....	527
Kenneth S. Williams, <i>On Euler's criterion for quintic nonresidues</i> .....	543
Manfred Wischnewsky, <i>On linear representations of affine groups. I</i> .....	551
Scott Andrew Wolpert, <i>Noncompleteness of the Weil-Petersson metric for Teichmüller space</i> .....	573
Volker Wrobel, <i>Some generalizations of Schauder's theorem in locally convex spaces</i> .....	579
Birge Huisgen-Zimmermann, <i>Endomorphism rings of self-generators</i> .....	587
Kelly Denis McKennon, <i>Corrections to: "Multipliers of type <math>(p, p)</math>"; "Multipliers of type <math>(p, p)</math> and multipliers of the group <math>L_p</math>-algebras"; "Multipliers and the group <math>L_p</math>-algebras"</i> .....	603
Andrew M. W. Glass, W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>Correction to: "<math>a^*</math>-closures to completely distributive lattice-ordered groups"</i> .....	606
Zvi Arad and George Isaac Glauberman, <i>Correction to: "A characteristic subgroup of a group of odd order"</i> .....	607
Roger W. Barnard and John Lawson Lewis, <i>Correction to: "Subordination theorems for some classes of starlike functions"</i> .....	607
David Westreich, <i>Corrections to: "Bifurcation of operator equations with unbounded linearized part"</i> .....	608