

# Pacific Journal of Mathematics

**CONTINUITY OF LINEAR MAPS FROM  $C^*$ -ALGEBRAS**

KJELD LAURSEN

## CONTINUITY OF LINEAR MAPS FROM $C^*$ -ALGEBRAS

K. B. LAURSEN

The celebrated problem of automatic continuity in Banach algebras—whether or not an arbitrary homomorphism from the algebra  $C(X)$  of all complex continuous functions on a compact Hausdorff space  $X$  is continuous—remains unsolved.

The lack of success on this point has generated quite a bit of effort to determine 'the extent' to which a homomorphism is continuous. In the basic work of W. G. Bade and P. C. Curtis around 1960 it was shown that a homomorphism is necessarily continuous on some dense subalgebra of the algebra  $C(X)$ .

Many of these results have later been shown to carry over to a much larger class of mappings, namely the separable maps (cf. Definition 1.1 below).

Recently, A. M. Sinclair has taken a new look at the homomorphism problems and succeeded in extending much of Bade's and Curtis's work to general  $C^*$ -algebras. In this paper we employ some of Sinclair's methods and obtain extensions of his main results, notably we prove (Theorem 3.7) that a separable linear map defined on a  $C^*$ -algebra  $A$  is necessarily continuous on a dense subalgebra of  $A$ .

The class of mappings considered here was introduced in [4]; we employ the generalized notion from [6].

**DEFINITION 1.1.** Let  $A, B, X, Y$  be normed linear spaces,  $q: A \times B \rightarrow X$  a continuous bilinear form. Let  $T: X \rightarrow Y$  be linear;  $T$  is said to be *separable* (with respect to  $q$ ) if there are functions  $f: A \rightarrow \mathbb{R}^+$  and  $g: B \rightarrow \mathbb{R}^+$  (where  $\mathbb{R}^+$  is the positive reals) such that  $\|T(q(a, b))\| \leq f(a)g(b)$  for all  $a \in A, b \in B$ .

As mentioned in [4] this class of mappings contains all algebra-homomorphisms, derivations, multipliers (centralizers); in fact, as an easy norm-estimate shows, if  $T$  is a continuous linear mapping, then  $T$  is separable with respect to any continuous bilinear  $q: A \times B \rightarrow X$ .

Since discontinuous derivations are known (cf. e.g. [2]), not every separable linear map is continuous. Nevertheless, as the following will show, certain results concerning the degree to which separable linear maps are continuous can be obtained.

The techniques employed here owe much to the work of Sinclair [5]; it is only appropriate to add to the list of examples of separable

maps the module homomorphisms considered in [5]: with the notation of Definition 1.1, let  $A$  be a Banach algebra,  $B = X$  a Banach left  $A$ -module,  $Y$  a Banach space with continuous module operations (i.e.  $Y$  is an  $A$ -module and for each  $a \in A$ ,  $y \rightarrow ay$  is a continuous operator) and  $T: X \rightarrow Y$  a module homomorphism. If  $q(a, x) = ax$  is the module multiplication on  $X$  we have  $\|T(q(a, x))\| = \|T(ax)\| = \|aT(x)\| \leq C_a \|T(x)\|$  where  $C_a$  is the norm-bound of the operator  $y \rightarrow ay: Y \rightarrow Y$ .

We list now the basic technical facts

**LEMMA 1.2** ([6 Lemma 1.1]). *If  $A, B$  are Banach spaces,  $X, Y$  normed spaces,  $q: A \times B \rightarrow X$  is a continuous bilinear form and  $T: X \rightarrow Y$  is a linear map separable with respect to  $q$ , then, with  $\{a_n\} \subset A$ ,  $\{b_k\} \subset B$  such that  $q(a_n, b_k) = 0$  whenever  $n \neq k$ , we have that*

$$\sup_n \|T(q(a_n, b_n))\| / \|a_n\| \|b_n\| < \infty .$$

*Proof.* Similar to the proof of [4, Lemma 1.1]

**COROLLARY 1.3** (cf. [5, Lemma 2.1 (b)]). *Suppose  $A$  is a Banach algebra,  $X$  a Banach left  $A$ -module and  $Y$  a normed linear space. If  $T: X \rightarrow Y$  is a linear map separable with respect to module multiplication, if  $\{a_n\}, \{b_k\} \subset A$  are sequences for which  $a_n b_k = 0$  whenever  $n \neq k$  then  $T \circ a_n b_n: X \rightarrow Y$  is continuous for all but finitely many  $n$  and the set  $\{\|T \circ a_n b_n\| / \|a_n\| \|b_n\| \mid T \circ a_n b_n \text{ is continuous}\}$  is bounded.*

*Proof.* If the conclusion fails then we may assume all the maps  $T \circ a_n b_n$  to be discontinuous and hence we can find  $\{x_n\} \subset X$  such that  $\|x_n\| \leq (\|a_n\| \|b_n\|)^{-1}$  but  $\|T(a_n b_n x_n)\| \geq n$  for  $n = 1, 2, \dots$ . Letting  $b_n x_n$  play the rôle of  $b_n$  in Lemma 1.2 we have a contradiction. Then rest of the corollary is then immediate.

**2. Commutative regular algebras.** Throughout this section  $A$  is assumed to be a regular semi-simple commutative unital Banach algebra,  $X$  is a Banach left  $A$ -module,  $Y$  a normed linear space and  $T: X \rightarrow Y$  a linear map, separable with respect to the module multiplication.

We now introduce the first version of the continuity ideal of  $T$  and study its size in  $A$ . The following results are analogues and generalizations of those of [6, §3]; the methods of proof that Stein employs are due to Bade and Curtis [1] and could be used here, too. Instead we have chosen the not dissimilar approach of [5, §2].

DEFINITION 2.1. With the assumptions of the first paragraph of this section let the continuity ideal of  $T$ ,  $I_T$ , be

$$I_T = \{a \in A \mid x \longrightarrow T(ax): X \longrightarrow Y \text{ continuous}\}.$$

It is clear that  $I_T$  is indeed an ideal.

As usual, if  $I$  is an ideal in a commutative Banach algebra  $B$  with maximal ideal space  $M_B$ , the hull of  $I$  is the set  $\{\varphi \in M_B \mid \varphi(I) = \{0\}\}$ . If  $F$  is a closed subset of  $M_B$  we let  $J(F)$  denote the set

$$J(F) = \{x \in B \mid \varphi(x) = 0 \text{ for every } \varphi \text{ in a neighborhood of } F\}.$$

We assume known the fact that if  $B$  is regular and semisimple then  $J(F)$  is minimal among all ideals in  $B$  with hull  $F$ .

The following generalizes [5, Theorem 2.2]

THEOREM 2.2. *Let  $A$  be a regular semisimple commutative unital Banach algebra, let  $X$  be a left Banach  $A$ -module and  $Y$  be a normed linear space. Suppose  $T: X \rightarrow Y$  is separable with respect to module multiplication. Then  $I_T = \{a \in A \mid x \rightarrow T(ax) \text{ is continuous}\}$  is an ideal in  $A$  with finite hull  $F$ ; moreover  $\exists$  constant  $C$  such that*

$$\|T \circ ab\| \leq C \|a\| \|b\|$$

for all  $a, b \in J(F)$ .

*Proof.* Let  $F$  be the hull of  $I_T$  and suppose  $F$  is infinite; then we can extract from  $F$  a sequence  $\{\varphi_n\}$  which does not converge to any of the points  $\varphi_n$ . We can therefore choose open sets  $U_1, U_2, \dots$  in  $M_A$  such that  $\varphi_n \in U_n$ ,  $n = 1, 2, \dots$  and such that  $U_j \cap U_n = \emptyset$  for all  $j < n$  and  $n = 1, 2, \dots$ . Now employ the regularity of  $A$  to find functions  $\{a_n\}$  such that  $\hat{a}_n$  vanishes off  $U_n$  and  $\hat{a}_n(\varphi_n) = 1$ . By choice of  $a_n$  we have  $a_n^2 \notin I_T$ ,  $n = 1, 2, \dots$ ; also  $a_n a_m = 0$  if  $n \neq m$ . Since  $a_n^2 \notin I_T$ ,  $T \circ a_n^2$  is discontinuous for each  $n$ . This contradicts Corollary 1.3.

To prove the boundedness of  $f(a, b) = \|T \circ ab\| / \|a\| \|b\|$  on  $J(F) \times J(F)$  we proceed as in [5, proof of Theorem 2.2] and suppose  $f$  unbounded on  $J(F) \times J(F)$ .

First we show that if  $U \subset M_A$  is an open neighborhood of  $F$  then

$$K_U = \{f(a, b) \mid a, b \in J(F) \cap J(M_A \setminus U)\}$$

is unbounded. Suppose  $K_U$  is bounded by  $C$ . Since  $A$  is normal on  $M_A$  we can find  $h \in A$  for which  $h$  is zero in a neighborhood of  $F$  and 1 in a neighborhood of  $M_A \setminus U$ . Since  $h$  is in  $J(F)$  and thus in  $I_T$ ,  $T \circ h$  is continuous. Let  $a, b \in J(F)$ ; then

$$\begin{aligned} \| T \circ ab \| &= \| T \circ (a - ha)(b - hb) + T \circ h(2 - h)ab \| \\ &\leq \| T \circ (a - ha)(b - hb) \| + \| T \circ h(2 - h)ab \| ; \end{aligned}$$

since  $a - hb, b - ha \in J(F) \cap J(M_A \setminus U)$  we have  $\| T \circ (a - ha)(b - hb) \| \leq C \| a - ha \| \| b - hb \|$ ; also

$$\| T \circ h(2 - h)ab \| \leq \| ab \| \| T \circ h(2 - h) \| ,$$

since  $h(2 - h) \in I_T$ . Collecting these estimates we get

$$\| T \circ ab \| \leq C \| 1 - h \|^2 \| a \| \| b \| + \| T \circ h(2 - h) \| \| a \| \| b \|$$

which contradicts the assumption that  $f$  be unbounded.

Thus  $K_U$  is an unbounded set for any open  $U \supseteq F$ . We use this to construct sequences that will contradict Corollary 1.3.

First choose  $a_1, b_1$  in  $J(F)$  such that  $f(a_1, b_1) \geq 1$  and let  $U_1$  be an open proper neighborhood of  $\{\varphi \in M_A \mid \varphi(a_1) = \varphi(b_1) = 0\}$ . Assume next that  $a_1, \dots, a_n, b_1, \dots, b_n$  have been found in  $J(F)$  such that  $f(a_j, b_j) \geq j$  and  $a_j, b_j \in J(M_A \setminus U_{j-1})$  where  $U_{j-1}$  is a proper open neighborhood of

$$\{\varphi \in M_A \mid \varphi(a_k) = \varphi(b_k) = 0, k = 1, \dots, j - 1\}$$

for  $j = 2, \dots, n$ . Since  $K_{U_n}$  is unbounded we can find  $a_{n+1}, b_{n+1} \in J(F) \cap J(M_A \setminus U_n)$  for which  $f(a_{n+1}, b_{n+1}) \geq n + 1$ . Since  $a_n$  vanishes where  $b_m$  is supported (if  $n \neq m$ ) the semi-simplicity of  $A$  implies that  $a_n b_m = 0$  whenever  $n \neq m$ . We have obtained a contradiction of Corollary 1.3.

The following is a bilinear variant of Theorem 2.2 which will be the basis of the work in the next section.

**THEOREM 2.3.** *Let  $A$  be a commutative regular semi-simple unital Banach algebra,  $X$  a Banach algebra and  $\Pi: A \rightarrow X$  a homomorphism. Let  $Y$  be a normed linear space,  $T: X \rightarrow Y$  a linear map, separable with respect to the algebra multiplication in  $X$ . Let*

$$I_T = \{a \in A \mid (b, c) \longrightarrow T(b\Pi(a)c): X \times X \longrightarrow Y \text{ is continuous}\}$$

*be the continuity ideal. Then  $I_T$  has finite hull  $F$  and  $\exists$  constant  $C$  such that the bilinear map*

$$\theta_{a,b}: X \times X \longrightarrow Y: (x_1, x_2) \longrightarrow T(x_1\Pi(ab)x_2)$$

*satisfies  $\| \theta_{a,b} \| \leq C \| \Pi(a) \| \| \Pi(b) \|$  for all  $a, b \in J(F)$ .*

*Proof.* Except for the changes necessitated by the linearity

being replaced by bilinearity the proof proceeds exactly as that of the preceding Theorem, so we omit it.

**THEOREM 2.4.** *In addition to the assumptions of Theorem 2.2 suppose  $J(F)$  has a bounded approximate identity. Then  $T$  is continuous on the subspace  $J(F)X$  of  $X$ .*

*Proof.* Since  $J(F)X = \text{span} \{ax \mid a \in J(F), x \in X\}$  we see that if  $z \in J(F)X$  then there is a  $b \in J(F)$  such that  $z = bz$ . Suppose  $\{e_\alpha\}$  is an approximate identity for  $J(F)$  with bound  $M$  and let  $C$  be the bound of Theorem 2.2. Since  $\{e_\alpha^2\}$  is also an approximate identity we may choose  $e \in \{e_\alpha\}$  such that  $\|b - e^2b\| \leq 1/\|b\|$ . Then

$$\begin{aligned} \|Tz\| &= \|Tb^2z\| \leq \|T(b - e^2b)bz\| + \|Te^2z\| \\ &\leq C\|b - e^2b\|\|b\|\|z\| + C\|e\|^2\|z\| \leq C(1 + M^2)\|z\|. \end{aligned}$$

**REMARK.** Recall that a commutative unital Banach algebra  $A$  is said to satisfy Ditkin's condition if  $J(\{\varphi\})$  contains a bounded approximate identity for each  $\varphi \in M_A$ . Then any ideal  $J(F)$  for which  $F$  is finite will contain a bounded approximate identity. Thus, the above result is a generalization of [6, Proposition 3.2] and is also an analogue of part of [5, Theorem 2.3].

The technique of the proof of Theorem 2.4 yields an analogous bilinear result, based on Theorem 2.3.

**THEOREM 2.5.** *With the assumptions of Theorem 2.3 and the additional assumptions that  $\Pi: A \rightarrow X$  be continuous and  $J(F) \subset A$  have a bounded approximate identity the bilinear map*

$$X \times J(F)X \longrightarrow Y: (x, t) \longrightarrow T(xt)$$

*is continuous.*

*Proof.* As in the proof of Theorem 2.4, if  $t \in J(F)X$ ,  $t = \sum \Pi(a_i)x_i$ , then  $\exists b \in J(F)$  such that  $t = \Pi(b)t$ ; also we can find  $e \in J(F)$  with  $\|\Pi(e)\| \leq \|\Pi\|M$  such that  $\|\Pi(b - e^2b)\| \leq 1/\|\Pi(b)\|$ . If  $x \in X$  then  $\|T(xt)\| \leq \|T(x\Pi(b - e^2b)\Pi(b)t)\| + \|T(x\Pi(e)^2t)\| \leq C\|\Pi(b - e^2b)\| \times \|\Pi(b)\|\|x\|\|t\| + C\|\Pi\|^2M^2\|x\|\|t\| \leq C(1 + \|\Pi\|^2M^2)\|x\|\|t\|$ .

**3.  $C^*$ -algebras.** In this section we restrict attention to  $C^*$ -algebras; we let  $A$  be a unital  $C^*$ -algebra,  $B$  a normed linear space and  $T: A \rightarrow B$  a linear map, separable with respect to the algebra multiplication on  $A$ . The main result is Theorem 3.7 which states that  $T$  is continuous on a dense subalgebra of  $A$  thus establishing the general version of [4, Theorem 5.6]. As in the previous sections

the techniques are mainly those of [5].

**PROPOSITION 3.1.** *Let  $C$  be a unital commutative  $C^*$ -subalgebra of  $A$ . Then there is a closed cofinite ideal  $\ker F$  in  $C$  such that*

$$\theta_1: A \ker F \times I_T \longrightarrow B$$

and

$$\theta_2: I_T \times \ker FA \longrightarrow B$$

defined by  $\theta_1(a, b) = T(ab)$ ,  $\theta_2(b, a) = T(ba)$ , are continuous bilinear maps. Here

$$I_T = \{b \in A \mid (a, c) \longrightarrow T(abc): A \times A \longrightarrow B \text{ is continuous}\}.$$

*Proof.* Let  $F_1$  be the singularity set of  $T$  as in Theorem 2.3, let  $a \in A \ker F_1$  and let  $b \in I_T$ . Let  $e$  denote the unit of  $A$  and choose  $\varepsilon > 0$ . By the Cohen factorization theorem applied to  $A \ker F_1$  there is  $c_1 \in A \ker F_1$  and  $c_2 \in \ker F_1$  such that  $a = c_1 c_2$  with  $\|c_1\| \leq 2\|a\|$ ,  $\|c_2\| = 1$ . Choose  $c_3, c_4$  and  $c_5$  in  $J(F)$  such that  $c_3 = c_4 c_5$ , such that  $\|c_2 - c_3\| < \varepsilon_1$  ( $\varepsilon_1$  to be specified) and such that  $\|c_4\| \|c_5\| \leq 2\|c_3\|$ . This is yet another application of Cohen's factorization theorem. By the analogous version of Theorem 2.5 the bilinear map  $(x, z) \rightarrow T(xz): AJ(F_1) \times A \rightarrow B$  is continuous and has a norm bound which does not exceed  $2C_1$ , where  $C_1$  is the constant of Theorem 2.3. Let  $C_b$  denote the norm bound of the continuous bilinear operator  $(x_1, x_2) \rightarrow T(x_1 b x_2)$ ; and note that  $\|c_2 - c_3\| < \varepsilon_1$ , so  $\|c_3\| < 1 + \varepsilon_1$ . We then have

$$\begin{aligned} \|T(ab)\| &= \|T(c_1 c_2 b)\| \\ &\leq \|T(c_1(c_2 - c_3)be)\| + \|T(c_1 c_3 b)\| \\ &\leq \|c_1\| \|c_2 - c_3\| C_b \|e\| + 2C_1 \|c_1\| \|c_3\| \|b\| \\ &\leq \varepsilon_1 \|c_1\| C_b + 2C_1 \cdot 2 \|a\| (1 + \varepsilon_1) \|b\| \\ &= \varepsilon_1(2C_b + 4C_1 \|b\|) \|a\| + 4C_1 \|a\| \|b\|. \end{aligned}$$

If

$$\varepsilon_1 < \frac{\varepsilon}{(2C_b + 4C_1 \|b\|) \|a\|}$$

then

$$\|T(ab)\| \leq 4C_1 \|a\| \|b\| + \varepsilon$$

which proves that  $\theta_1$  is continuous on  $A \ker F_1 \times I_T$ . Similarly we get a finite set  $F_2$  such that  $\theta_2$  is continuous on  $I_T \times \ker F_2 A$ . If

we take  $F = F_1 \cup F_2$  the proposition follows.

**COROLLARY 3.2.** *The trilinear maps*

$$\Psi_1: A \times \ker F \times I_T \longrightarrow B$$

$$\Psi_2: I_T \times \ker F \times A \longrightarrow B$$

defined by  $\Psi_1(a, b, c) = T(abc)$ ,  $\Psi_2(c, b, a) = T(cba)$  are continuous.

On the basis of the above corollary we make the following definition.

**DEFINITION 3.3.**

$$K_T = \{a \in A \mid (c, d) \longrightarrow T(cad): A \times I_T \longrightarrow B \text{ and} \\ (d, c) \longrightarrow T(dac): I_T \times A \longrightarrow B \text{ are continuous}\}.$$

About this set we have the following

**PROPOSITION 3.4.**  *$K_T$  is a closed cofinite two-sided ideal that contains  $I_T$ .*

*Proof.* It is immediate that  $K_T$  is a two-sided ideal that contains  $I_T$ . By [5, Lemma 3.2], to show that  $K_T$  is closed and cofinite, it is sufficient to show that  $K_T \cap C$  is closed and cofinite for any commutative unital  $C^*$ -subalgebra of  $C$ . Corollary 3.2 shows that the hull of  $K_T \cap C$  is finite. If  $\ker F \cong K_T \cap C$  let  $c_1, c_2, \dots, c_n$  be chosen in  $K_T \cap C$  such that  $c_i c_j = 0$  if  $i \neq j$  and such that  $c_i = 1$  in a neighborhood of  $\varphi_i$  where

$$\{\varphi_1, \dots, \varphi_n\} = F \setminus \text{hull}(K_T \cap C)$$

and such that  $c_i(\varphi_j) = 0$  if  $j \neq i$ .

If  $a \in K_T \cap C$  then  $a - \sum a(\varphi_i) c_i \in \ker F$  and since  $\ker F$  is closed it follows from the definition of  $K_T$  that  $K_T \cap C$  is closed: suppose  $\{a_m\} \subset K_T \cap C$  converges to  $a$ ; then  $a_m - \sum a_m(\varphi_i) c_i \rightarrow a - \sum a(\varphi_i) c_i$ , so if  $c \in X$  and  $d \in I_T$  then

$$T(c(a - \sum a(\varphi_i) c_i) d) = T(cad) - \sum a(\varphi_i) T(cc_i d)$$

which shows that  $(c, d) \rightarrow T(cad)$  is continuous, i.e. that  $a \in K_T \cap C$ .

**COROLLARY 3.5.** *The bilinear maps*

$$(a, b) \longrightarrow T(ab): K_T \times I_T \longrightarrow B$$

and

$$(b, a) \longrightarrow T(ba): I_T \times K_T \longrightarrow B$$



are continuous.

The next result corresponds to [5, Theorem 3.8] and is a generalization of [4, Theorem 5.1].

**THEOREM 3.6.** *In addition to the above assumptions suppose  $B$  to be a Banach space. Then  $T$  is continuous on the two sided ideal  $K_T I_T$  which is dense in  $I_T$ .*

*Proof.* Let  $\Psi$  be the extension by continuity to  $K_T \times I_T^-$  of the mapping  $(a, b) \rightarrow T(ab)$  mentioned in Corollary 3.5. We may also consider  $\Psi$  a continuous linear operator from the projective tensor product  $K_T \otimes_r I_T^-$  into  $B$ . Let  $\theta: K_T \otimes_r I_T^- \rightarrow I_T^-$  be defined by  $\theta(\sum x_i \otimes y_i) = \sum x_i y_i$ . By Cohen's factorization theorem  $\theta$  is surjective so by the open mapping theorem  $\theta^{-1}: I_T^- \rightarrow K_T \otimes_r I_T^- / \ker(\theta)$  is continuous.

As we shall see below,  $\ker \theta \subseteq \ker \Psi$  so the composition  $\Phi = \Psi \circ \theta^{-1}: I_T^- \rightarrow B$  is well defined and continuous. On  $K_T I_T \subseteq I_T^-$  we have  $\Phi(ab) = \Psi(a, b) = T(ab)$  and consequently  $T$  is continuous on  $K_T I_T$ .

It remains to see that  $\ker \theta \subseteq \ker \Psi$ .

To that end we can refer to Sinclair's proof in [5, Theorem 3.8] in which a similar claim is established.

The density of  $K_T I_T$  in  $I_T$  is clear from the fact that  $K_T \supseteq I_T$ . That completes the proof.

We are now in a position to state and prove the main result of this section (cf. the introduction to this section).

**THEOREM 3.7.** *Let  $A$  be a  $C^*$ -algebra,  $B$  a Banach space,  $T$  a linear map of  $A$  into  $B$ , separable with respect to algebra multiplication. Then  $T$  is continuous on a dense subalgebra of  $A$ .*

*Proof.* By Theorem 3.6  $T$  is continuous on the two-sided ideal  $K_T I_T$  which has cofinite closure [4, p. 501]. By [3, Theorem 6] there is a finite dimensional subspace  $F$  of  $A$  such that  $K_T I_T + F$  is a dense subalgebra on which  $T$  is clearly continuous. That proves the theorem.

## REFERENCES

1. W. G. Bade and P. C. Curtis Jr., *Homomorphisms of commutative Banach algebras*, Amer. J. Math., **82** (1960) 589-608.
2. A. Browder, *Introduction to Function Algebras*, Benjamin, New York, 1969.
3. K. B. Laursen and A. M. Sinclair, *Lifting matrix units in  $C^*$ -algebras II*, Math. Scand., **37** (1975), 167-172.

4. K. B. Laursen and J. D. Stein Jr., *Automatic continuity in Banach spaces and algebras*, Amer. J. Math., **95** (1973), 495-505.
5. A. M. Sinclair, *Homomorphisms from  $C^*$ -algebras*, Proc. London Math. Soc., (3) **29** (1974), 435-452.
6. J. D. Stein Jr. *Some aspects of automatic continuity*, Pacific J. Math., **50** (1974), 187-204.

Received January 30, 1975 and in revised form November 4, 1975.

MATHEMATICAL INSTITUTE  
UNIVERSITETSPARKEN 5  
2100 COPENHAGEN  
DENMARK



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)

University of California  
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics  
Manufactured and first issued in Japan

# Pacific Journal of Mathematics

Vol. 61, No. 2

December, 1975

|   |     |
|---|-----|
| Graham Donald Allen, Francis Joseph Narcowich and James Patrick Williams, <i>An operator version of a theorem of Kolmogorov</i> .....   | 305 |
| Joel Hilary Anderson and Ciprian Foias, <i>Properties which normal operators share with normal derivations and related operators</i> .....  | 313 |
| Constantin Gelu Apostol and Norberto Salinas, <i>Nilpotent approximations and quasinilpotent operators</i> .....  | 327 |
| James M. Briggs, Jr., <i>Finitely generated ideals in regular <math>F</math>-algebras</i> .....   | 339 |
| Frank Benjamin Cannonito and Ronald Wallace Gatterdam, <i>The word problem and power problem in 1-relator groups are primitive recursive</i> .....  | 351 |
| Clifton Earle Corzatt, <i>Permutation polynomials over the rational numbers</i> .....   | 361 |
| L. S. Dube, <i>An inversion of the <math>S_2</math> transform for generalized functions</i> .....   | 383 |
| William Richard Emerson, <i>Averaging strongly subadditive set functions in unimodular amenable groups. I</i> .....   | 391 |
| Barry J. Gardner, <i>Semi-simple radical classes of algebras and attainability of identities</i> .....  | 401 |
| Irving Leonard Glicksberg, <i>Removable discontinuities of <math>A</math>-holomorphic functions</i> .....   | 417 |
| Fred Halpern, <i>Transfer theorems for topological structures</i> .....   | 427 |
| H. B. Hamilton, T. E. Nordahl and Takayuki Tamura, <i>Commutative cancellative semigroups without idempotents</i> .....   | 441 |
| Melvin Hochster, <i>An obstruction to lifting cyclic modules</i> .....  | 457 |
| Alistair H. Lachlan, <i>Theories with a finite number of models in an uncountable power are categorical</i> .....   | 465 |
| Kjeld Laursen, <i>Continuity of linear maps from <math>C^*</math>-algebras</i> .....  | 483 |
| Tsai Sheng Liu, <i>Oscillation of even order differential equations with deviating arguments</i> .....  | 493 |
| Jorge Martinez, <i>Doubling chains, singular elements and hyper-<math>Z</math> <math>l</math>-groups</i> .....  | 503 |
| Mehdi Radjabalipour and Heydar Radjavi, <i>On the geometry of numerical ranges</i> .....  | 507 |
| Thomas I. Seidman, <i>The solution of singular equations, I. Linear equations in Hilbert space</i> .....  | 513 |
| R. James Tomkins, <i>Properties of martingale-like sequences</i> .....  | 521 |
| Alfons Van Daele, <i>A Radon Nikodým theorem for weights on von Neumann algebras</i> .....  | 527 |
| Kenneth S. Williams, <i>On Euler's criterion for quintic nonresidues</i> .....  | 543 |
| Manfred Wischnewsky, <i>On linear representations of affine groups. I</i> .....   | 551 |
| Scott Andrew Wolpert, <i>Noncompleteness of the Weil-Petersson metric for Teichmüller space</i> .....   | 573 |
| Volker Wrobel, <i>Some generalizations of Schauder's theorem in locally convex spaces</i> .....   | 579 |
| Birge Huisgen-Zimmermann, <i>Endomorphism rings of self-generators</i> .....  | 587 |
| Kelly Denis McKennon, <i>Corrections to: "Multipliers of type <math>(p, p)</math>"; "Multipliers of type <math>(p, p)</math> and multipliers of the group <math>L_p</math>-algebras"; "Multipliers and the group <math>L_p</math>-algebras"</i> ..... | 603 |
| Andrew M. W. Glass, W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>Correction to: "<math>a^*</math>-closures to completely distributive lattice-ordered groups"</i> .....  | 606 |
| Zvi Arad and George Isaac Glauberman, <i>Correction to: "A characteristic subgroup of a group of odd order"</i> .....   | 607 |
| Roger W. Barnard and John Lawson Lewis, <i>Correction to: "Subordination theorems for some classes of starlike functions"</i> .....   | 607 |
| David Westreich, <i>Corrections to: "Bifurcation of operator equations with unbounded linearized part"</i> .....  | 608 |