

Pacific Journal of Mathematics

ON THE GEOMETRY OF NUMERICAL RANGES

MEHDI RADJABALIPOUR AND HEYDAR RADJAVI

ON THE GEOMETRY OF NUMERICAL RANGES

MEHDI RADJABALIPOUR AND HEYDAR RADJAVI

A bounded convex set G in the plane is the numerical range of an operator on a separable Hilbert space if $G \setminus G^\circ$ is a countable union of arcs of conic sections and singletons. This result answers, in particular, a question raised by Joel Anderson.

O. Introduction. Throughout this paper T denotes a (bounded linear) operator on a separable complex Hilbert space H . The numerical range of T , denoted by $W(T)$, is defined as the set of all complex numbers $(Tx | x)$ with $\|x\| = 1$. The problem of determining all (bounded and convex) subsets of the plane which are attainable as numerical ranges seems to be hard. In fact, the question of attainability is still unsettled for some very simple sets.

Joel Anderson has made several (still unpublished) contributions to the subject; they include the following elegant result in the case where H has finite dimension n : If $W(T)$ is contained in the closed unit disk and if it intersects the unit circle in more than n points, then it coincides with the unit disk. Anderson asked whether the assertion would be true if n were replaced by \aleph_0 . He asked, in particular, whether the closed half-disk is attainable as the numerical range of an operator T ; he proved that it is not if T is compact. See [4]. We shall show, as a corollary of a more general result, that the closed half-disk is attainable by a rank-one perturbation of a hermitian operator.

It should be noted, perhaps, that if the space has dimension $\geq 2^{\aleph_0}$ then every bounded convex subset of the plane is the numerical range of a normal operator [5]. The fact that this is not true in the separable case is well known and easily demonstrated by a cardinality argument. In fact, if G is any convex open set whose closure contains uncountably many extreme points, then there exist convex sets with interior G which are not attainable. No concrete examples of non-attainable bounded convex sets are known to us. It is proved in this paper that a convex set G is the numerical range of an operator T if $G \setminus G^\circ = E_0 \cup E_1$, where E_0 is countable and E_1 is a union of quadratic arcs.

We are grateful to J. Bastian and P. Fillmore for stimulating conversations.

1. Preliminaries. Let G be a convex set in the plane and let λ be in the boundary of G . A line L is called a *support* of G at λ

if $\lambda \in L$ and G lies in one of the closed half-planes determined by L . A point $\lambda \in G$ is called an *extreme point* of G if λ does not belong to any open line segment lying in $W(T)$. An extreme point λ of G is called a *sharp point* of G if G has more than one support at λ . (We use the notations G^- and G^0 for the closure and the interior of a set G , respectively.)

We will make use of the following Theorems *A – E*. Most of them are known; we have included the references wherever possible and have provided proofs otherwise.

THEOREM A. *The numerical range $W(T)$ of T is convex and $W(aT + b) = aW(T) + b$ for all complex numbers a and b . Moreover if P is an arbitrary nonzero projection, then $W(PT|PH) \subseteq W(T)$. (See [3, Problem 166] and the references cited there.)*

THEOREM B. *Let λ be a sharp point of $W(T)$. Then $T = S \oplus \lambda I$, where $\lambda \notin W(S)$. (See [1, Theorem 1].)*

THEOREM C. *Let T be the direct sum of any family $\{T_\alpha\}_{\alpha \in J}$ of operators. Then $W(T) = \text{Co}(\bigcup_{\alpha \in J} W(T_\alpha))$. (Here $\text{Co}(G)$ denotes the convex hull of a set G .)*

The proof follows from the fact that if G is a convex set and if $\{\lambda_i\}$ is a sequence in G , then $\sum \alpha_i \lambda_i \in G$ for all sequences $\{\alpha_i\}$ of nonnegative numbers satisfying $\sum \alpha_i = 1$.

The following theorem is a combination of the results of [2] and [6]. We leave the proof to the reader.

THEOREM D. *Let T be hyponormal and let L be a support of $W(T)$. Then the set*

$$M = \{x \in H: (Tx | x) = \lambda \|x\|^2 \text{ for some } \lambda \in L\}$$

is a reducing (trivial or nontrivial) invariant subspace of T and $W(T|M) \subseteq L$. In particular $W(T) \setminus W(T)^0 = E_0 \cup E_1$, where E_0 is a countable set and E_1 is a countable union of (not necessarily closed) line segments. Moreover if $W(T) \setminus W(T)^0 \neq \emptyset$, then T has a non-trivial normal part.

The following theorem can be regarded as a converse to Theorem D.

THEOREM E. *Let G be a convex set with countably many extreme points. Then there exists a normal operator T such that $W(T) = G$.*

Proof. It follows from the assumption on G that

$$G \setminus G^0 = \bigcup_{\alpha \in J} E_\alpha,$$

where J is a (possibly empty) countable index set and where E_α is either a singleton or a line segment ($\alpha \in J$). Let T_α be a normal operator with numerical range E_α and let M_z be the multiplication by z in $L^2(G, dx dy)$. In view of Theorems C and D, $W(M_z \oplus \sum_{\alpha \in J} \oplus T_\alpha) = G$. (Note that the numerical range of M_z is G^0 .)

2. **Main results.** In what follows a *quadratic arc* means a (not necessarily closed) smooth subarc of a conic section.

THEOREM 1. *Let G be any closed region whose boundary consists of a closed line segment L and a closed quadratic arc C . Then there exists an operator T with the following properties:*

- (i) T is irreducible,
- (ii) $T = A + B$ where $W(A) \subseteq L$ and B is a normal operator of rank 1,
- (iii) $W(T) = G \setminus (L \cap C)$.

Proof. In view of Theorem A we can assume, without loss of generality, that L is the closed real interval $[-2, 2]$ and C lies in the upper half plane. Let (α, β) be the point on C at which the tangent to C is parallel to the x -axis. It is easy to verify that the equation of an arbitrary conic section having the above properties is given by the formula

$$(1) \quad \beta x^2 + b\beta y^2 - 2\alpha xy + (4 + \alpha^2 - b\beta^2)y - 4\beta = 0,$$

where

$$b > (\alpha^2 - 4)/\beta^2.$$

(The latter inequality follows from the fact that β is the maximum ordinate on the arc C .)

Let $\gamma = (1/2)(b\beta^2 - \alpha^2 + 4)^{1/2}$. Let H be a Hilbert space with an orthonormal basis $\{e_n\}_{1 \leq n < \infty}$. Define an operator T as follows:

$$Te_1 = (\alpha + \beta i)e_1 + \gamma e_2, \quad Te_2 = \gamma e_1 + e_3,$$

and

$$Te_n = e_{n-1} + e_{n+1} \quad \text{for } n = 3, 4, \dots$$

We claim that T has the required properties (i)-(iii). The first two properties are easy; we prove only the last one. For $t \in [0, 1]$, let

M_t be the set of all unit vectors u in H such that $|(u | e_1)|^2 = t$. Let S be the weighted shift defined by $Se_1 = \gamma e_2$ and $Se_n = e_{n+1}$ for $n = 2, 3, \dots$. Let $\Delta_t = \{2\operatorname{Re}(Su | u) : u \in M_t\}$. Computation shows that the points of $W(T)^-$ are determined by the following relations.

$$(2) \quad \begin{cases} \alpha t + \inf \Delta_t \leq x \leq \alpha t + \sup \Delta_t, \\ y = \beta t, \quad 0 \leq t \leq 1. \end{cases}$$

Since $|(Su | u)|^2 \leq (1-t)(\gamma^2 t - t + 1)$, it follows that $W(T)^-$ is a subset of all points (x, y) such that

$$\begin{cases} \beta x^2 + b\beta y^2 - 2\alpha xy + (4 + \alpha^2 - b\beta^2)y - 4\beta \leq 0, \\ \text{and} \quad 0 \leq y \leq \beta. \end{cases}$$

For $t \neq 0$, let $\lambda = \pm[(1-t)/(1-t+\gamma^2 t)]^{1/2}$ and $u = t^{1/2}(e_1 + \lambda\gamma e_2 + \dots + \lambda^{n-1}\gamma e_n + \dots)$. It is easy to see that $\|u\| = 1$ and $(Su | u) = \pm[(1-t)(\gamma^2 t - t + 1)]^{1/2}$. Thus the arc

$$\begin{cases} \beta x^2 + b\beta y^2 - 2\alpha xy + (4 + \alpha^2 - b\beta^2)y - 4\beta = 0, \\ \text{and} \quad 0 < y \leq \beta \end{cases}$$

lies in $W(T)$.

Let P be the projection onto the closed span of $\{e_3, e_4, \dots\}$. Since $(1/2)PT|PH$ is the real part of a simple shift, $W(T) \supseteq W(PT|PH) = (-2, 2)$. Therefore $W(T) = G \setminus \{-2, 2\}$ because $W(T)$ cannot have any sharp point (see Theorem B).

COROLLARY 1. *Let G be a convex set such that $G \setminus G^0 = E_0 \cup E_1$ where E_0 is a countable set and E_1 is a union of quadratic arcs. Then G is the numerical range of an operator.*

The proof is an easy consequence of Theorems C and 1 and the proof of Theorem E.

3. Remarks on the topology of $W(T)$. (a) Although $W(T)$ may not be closed, there always exists an operator S similar to T with $W(S)$ closed. The proof follows from the existence [7, Theorem 3] of an operator F of rank 2 such that $1 + F$ is invertible and

$$W((1 + F)^{-1}T(1 + F)) \supseteq W_e(T),$$

where $W_e(T) = \bigcap \{W(T + K)^- : K \text{ compact}\}$. Now let $S = (1 + F)^{-1}T(1 + F)$. Since $W(S) \supseteq W_e(S)$, it follows from [4, Theorem 1] that $W(S)$ is closed.

(b) Let $U = \{x \in H : \|x\| = 1\}$ and let $f(x) = (Tx | x)$ for $x \in U$. Obviously f is continuous in norm and $W(T) \setminus W(T)^0$ is the image of the closed set $f^{-1}(\partial W(T))$ under f . This shows that $W(T) \setminus W(T)^0$ is

an analytic set. The following question seems to be open: is $W(T) \setminus W(T)^{\circ}$ a Borel set? In view of [4, Theorem 1] the answer is in the affirmative if $W_{\circ}(T)$ has a countable number of extreme points.

(c) Corollary 1 suggests another (probably open) question: is every compact convex set the numerical range of some operator on H ? The corresponding question on open convex sets is easily answered in Theorem E.

REFERENCES

1. W. F. Donoghue, *On the numerical range of a bounded operator*, Michigan Math. J., **4** (1957), 261-263.
2. M. R. Embry, *the numerical range of an operator*, Pacific J. Math., **32** (1970), 647-650.
3. P. R. Halmos, *A. Hilbert Space Problem Book*, Van Nostrand, Princeton (1967).
4. J. S. Lancaster, *The boundary of the numerical range*, Proc. Amer. Math. Soc., **49** (1975), 393-398.
5. F. M. Pollack, *Numerical range and convex sets*, Canad. Math. Bull., **17** (1974), 295-296.
6. J. G. Stampfli, *Extreme points of the numerical range of hyponormal operators*, Michigan Math. J., **13** (1966), 87-89.
7. J. P. Williams, *Similarity and the numerical range*, J. Math. Anal. Appl., **36** (1969), 307-314.

Received May 5, 1975. This research was supported by a grant from N. R. C.

DALHOUSIE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 61, No. 2

December, 1975

Graham Donald Allen, Francis Joseph Narcowich and James Patrick Williams, <i>An operator version of a theorem of Kolmogorov</i>	305
Joel Hilary Anderson and Ciprian Foias, <i>Properties which normal operators share with normal derivations and related operators</i>	313
Constantin Gelu Apostol and Norberto Salinas, <i>Nilpotent approximations and quasinilpotent operators</i>	327
James M. Briggs, Jr., <i>Finitely generated ideals in regular F-algebras</i>	339
Frank Benjamin Cannonito and Ronald Wallace Gatterdam, <i>The word problem and power problem in 1-relator groups are primitive recursive</i>	351
Clifton Earle Corzatt, <i>Permutation polynomials over the rational numbers</i>	361
L. S. Dube, <i>An inversion of the S_2 transform for generalized functions</i>	383
William Richard Emerson, <i>Averaging strongly subadditive set functions in unimodular amenable groups. I</i>	391
Barry J. Gardner, <i>Semi-simple radical classes of algebras and attainability of identities</i>	401
Irving Leonard Glicksberg, <i>Removable discontinuities of A-holomorphic functions</i>	417
Fred Halpern, <i>Transfer theorems for topological structures</i>	427
H. B. Hamilton, T. E. Nordahl and Takayuki Tamura, <i>Commutative cancellative semigroups without idempotents</i>	441
Melvin Hochster, <i>An obstruction to lifting cyclic modules</i>	457
Alistair H. Lachlan, <i>Theories with a finite number of models in an uncountable power are categorical</i>	465
Kjeld Laursen, <i>Continuity of linear maps from C^*-algebras</i>	483
Tsai Sheng Liu, <i>Oscillation of even order differential equations with deviating arguments</i>	493
Jorge Martinez, <i>Doubling chains, singular elements and hyper-Z l-groups</i>	503
Mehdi Radjabalipour and Heydar Radjavi, <i>On the geometry of numerical ranges</i>	507
Thomas I. Seidman, <i>The solution of singular equations, I. Linear equations in Hilbert space</i>	513
R. James Tomkins, <i>Properties of martingale-like sequences</i>	521
Alfons Van Daele, <i>A Radon Nikodým theorem for weights on von Neumann algebras</i>	527
Kenneth S. Williams, <i>On Euler's criterion for quintic nonresidues</i>	543
Manfred Wischnewsky, <i>On linear representations of affine groups. I</i>	551
Scott Andrew Wolpert, <i>Noncompleteness of the Weil-Petersson metric for Teichmüller space</i>	573
Volker Wrobel, <i>Some generalizations of Schauder's theorem in locally convex spaces</i>	579
Birge Huisgen-Zimmermann, <i>Endomorphism rings of self-generators</i>	587
Kelly Denis McKennon, <i>Corrections to: "Multipliers of type (p, p)"; "Multipliers of type (p, p) and multipliers of the group L_p-algebras"; "Multipliers and the group L_p-algebras"</i>	603
Andrew M. W. Glass, W. Charles (Wilbur) Holland Jr. and Stephen H. McCleary, <i>Correction to: "a^*-closures to completely distributive lattice-ordered groups"</i>	606
Zvi Arad and George Isaac Glauberman, <i>Correction to: "A characteristic subgroup of a group of odd order"</i>	607
Roger W. Barnard and John Lawson Lewis, <i>Correction to: "Subordination theorems for some classes of starlike functions"</i>	607
David Westreich, <i>Corrections to: "Bifurcation of operator equations with unbounded linearized part"</i>	608