

Pacific Journal of Mathematics

**ABELIAN AND NILPOTENT SUBGROUPS OF MAXIMAL
ORDER OF GROUPS OF ODD ORDER**

ZVI ARAD

ABELIAN AND NILPOTENT SUBGROUPS OF MAXIMAL ORDER OF GROUPS OF ODD ORDER

ZVI ARAD

Denote the *maximum* of the orders of all nilpotent subgroups A of class at most c , of a finite group G , by $d_c(G)$. Let $A_c(G)$ be the set of all nilpotent subgroups of class at most c and having order $d_c(G)$ in G . Let $A_\infty(G)$ denote the set of all nilpotent subgroups of maximal order of a group G .

The aim of this paper is to investigate the set $A_\infty(G)$ of groups G of odd order and the structure of the groups G with the property $A_2(G) \subseteq A_\infty(G)$. Theorem 1 gives an expression for the number of elements in $A_\infty(G)$. Theorem 2 gives criteria for the nilpotency of groups of odd order.

In this paper G is a finite group, and π is a set of primes. If G is of odd order, then G is solvable [6].

1. Introduction. Denote the *maximum* of the orders of all nilpotent subgroups A of class at most c , of a finite group G , by $d_c(G)$. Let $A_c(G)$ be the set of all nilpotent subgroups of class at most c and having order $d_c(G)$ in G . Then $J_c(G)$ is the subgroup of G generated by $A_c(G)$. In particular, $J_1(G) = J(G)$ is the Thompson subgroup of G . Moreover, $A_\infty(G)$ is the set of all nilpotent subgroups of maximal order of a group G . Here $J_\infty(G)$ is the subgroup of G generated by the elements of $A_\infty(G)$.

In this paper G is a finite group, and π is a set of primes. If G is of odd order, then G is solvable [6].

The aim of this paper is to investigate the set $A_\infty(G)$ for groups G of odd order and the structure of the groups G with the property $A_2(G) \subseteq A_\infty(G)$.

We shall give, in Theorem 1, an expression for the number of elements in $A_\infty(G)$. In Theorem 2 we shall state criteria for the nilpotency of groups of odd order.

For groups G with the property $A_2(G) \subseteq A_\infty(G)$, we have the following:

THEOREM 3. *Let G be a π -solvable group with an S_π -subgroup K of G . Assume that $O_{\pi'}(G) = 1$ and that $A \in A_2(K) \cap A_\infty(K) \neq \emptyset$, then*

- (i) If either $2, 3 \notin \pi$ or $O_2(A)$ is Abelian, then $O_p(A) = O_p(G) = O_p(K)$, for every $p \in \pi$.
- (ii) If $F(G)$ is odd and if G has an Abelian S_2 -subgroup, then $J_\infty(K) = J_\infty(G) = F(G) = F(K) = A$.

Three corollaries of Theorem 3 and further information can be found in Chapter 2.

Our notation is standard and is taken mainly from [8]. In particular, $\pi(G)$ will designate the set of primes dividing $|G|$ and G_π denotes an S_π -subgroup of G . For the definitions of *Sylow system*, *system normalizers* and *Carter subgroups* of a group G see [11], Definition 11.1 p. 726 and Definition 12.1 p. 736.

2. Statements and proofs of the main theorems. The next result is needed for the proofs of the main Theorems.

PROPOSITION 1. *Suppose G is a group. Assume that A normalizes a nilpotent subgroup B of G , and assume that at least one of the following conditions holds:*

- (i) $A \in A_1(G)$, and B is Abelian ([3], Proposition 1).
- (ii) $A \in A_1(G)$, $|A|$ is odd, and an S_2 -subgroup of B is Abelian ([3], Proposition 1).
- (iii) $A \in A_2(G)$ ([7]).
- (iv) $A \in A_c(G)$, $c \geq 2$, $|B|$ is odd, and an S_2 -subgroup of A is Abelian ([4]).
- (v) $A \in A_x(G)$, $|B|$ is odd and an S_2 -subgroup of A is Abelian ([4]).

Then AB is nilpotent.

Define $A_x^p(G)$ to be the set $\{A_p/A \in A_x(G)\}$ of distinct p -subgroups of a group G , where p is a prime.

THEOREM 1. *Let G be a group of odd order. Then*

- (i) $|A_x(G)| = [G : N_G(A)] = \prod_{p \in \pi(A)} |A_x^p(G)|$, where $A \in A_x(G)$.
- (ii) $|A_x^p(G)| \equiv 1 \pmod{p}$
- (iii) $|A_x^p(G)|/h_p$, where $h_p = [G : N_G(G_p)]$
- (iv) $G = \langle N_G(A)/A \in A_x(G) \rangle$.
- (v) If $A \in A_x(G)$ and $A_p \subseteq G_p$ then there exists $x \in G$ such that $G_p \cap G_p^x = A_p$.

For a discussion of the number h_p see [9], Theorem 9.3.1.

Proof. Proposition 1(v) implies that every element of $A_x(G)$ con-

tains $F(G)$. Therefore [13], Theorem 1 implies that $A_\infty(G)$ is a conjugate class. If $A, B \in A_\infty(G)$ then $[A_p, B_q] = 1$ for every two distinct primes p and q by [11], Theorem 7.18, p. 705, proving (i). Let $A_p \subseteq G_p$. We shall prove that $N_G(G_p) \subseteq N_G(A_p)$. If $x \in N_G(G_p)$ then $\langle A_p^x, A_p \rangle \subseteq G_p$ is a p -group. Hence, $[\langle A_p^x, A_p \rangle, A_p] = 1$. Therefore $\langle A_p^x, A_p \rangle A_p$ is nilpotent. Consequently, $A_p^x = A_p$, and $N_G(G_p) \subseteq N_G(A_p)$. Part (i) implies that $A_\infty^2(G)$ is a conjugate class. Hence, $|A_\infty^2(G)| = [G : N_G(A_p)]$. By [14], Theorem 6.2.3, $|A_\infty^2(G)| \equiv 1 \pmod{p}$. Thus (ii)–(iii) also hold. Let $\mathcal{F} = \{G_p/p \in \pi(G)\}$ be a Sylow system of G . Let us denote the intersection of the normalizers of the subgroups of the given Sylow system by $N(\mathcal{F})$. By definition $N(\mathcal{F})$ is a system normalizer of G . Since $A_\infty(G)$ is a conjugate class of G there exists $A(p) \in A_\infty(G)$ such that $[A(p)]_p \subseteq G_p$, for every $p \in \pi[A(p)]$.

As above $A = x[A(p)]_p$ is an element of $A_\infty(G)$, moreover $N_G(A) \supseteq N(\mathcal{F})$. Since G is generated by the set of system normalizers of G [11], we obtain that $G = \langle N_G(A) / A \in A_\infty(G) \rangle$, proving (iv). Proposition 1 implies that $A_p = O_p(N_G(A_p))$. As mentioned before $G_p \subseteq N_G(A_p)$. Therefore by Ito's theorem [12] there exists $x \in G$ such that $G_p \cap G_p^x = A_p$.

The author knows of no counterexample to the conjecture that if G is an arbitrary group then $A_\infty(G)$ is a conjugate class.

Let ϕ be the class of finite solvable groups in which the system normalizers are Carter subgroups. ϕ -groups are discussed in [5] and [11], pp. 743–751.

THEOREM 2. *Let G be a group of odd order. Then*

(i) *If $G = BC$, where $B \in A_b(G)$, $C \in A_c(G)$, $b \geq 1$, $c \geq 1$, then G is nilpotent.*

(ii) *G is nilpotent if and only if $G \in \phi$ and the Carter subgroups of G are elements of $A_c(G)$, for some integer c .*

Proof. (i) Proposition 1 implies that $BF(G)$ and $CF(G)$ are nilpotent. Therefore, Theorem 1(i) and [13], Theorem 1 imply that there exist $x \in G$ and $A \in A_\infty(G)$ such that $B \subseteq A$ and $C \subseteq A^x$. Hence $G = AA^x$. By [11], Theorem 7.18, p. 705 G is nilpotent.

(ii) Let C be a Carter subgroup of G . Assume that $C \in A_c(G)$, for some integer c . Since $CF(G)$ is nilpotent by Proposition 1, there exists $A \in A_\infty(G)$ such that $C \subseteq A$. Since $N_G(C) = C$, we obtain that $C = A$. By assumption $G \in \phi$. Therefore A is a system normalizer. By the definition of system normalizer, Proposition 1 implies that G is nilpotent.

REMARK. Theorem 2(i) is true without assuming [6].

If $G = BC$, where $B \in A_b(G)$, $C \in A_c(G)$, $b \geq 1$, $c \geq 1$, then G is solvable by [14], Theorem 13.2.9.

We shall say that G is a D_π -group if all the maximal π -subgroups of G are conjugate S_π -subgroups of G . If G and every normal subgroup of G is a D_π -group we will call G a D_π^N -group.

Let $[A, B, C]$ denotes the triple commutator $[[A, B], C]$ of three subgroups A, B, C of G . We say that G is a π -stable group if it satisfies the following condition:

Let K be an arbitrary π -subgroup of G . Let A be an arbitrary π -subgroup of $N_G(K)$. Then if $[K, A, A] = 1$, we have $AC_G(K)/C_G(K) \subseteq O_\pi(N_G(K)/C_G(K))$.

The next result is needed for the proof of Theorem 3.

PROPOSITION 2. Let G be a π -stable D_π^N -group. Let K be an S_π -subgroup of G . Assume that $A_2(K) \subseteq A_\infty(K)$, $C_G(F(G)) \subseteq F(G)$ and $O_{\pi'}(G) = 1$. Then we have

- (i) $J_2(K) \text{ char } G$
- (ii) If $|F(G)|$ is odd and G has an Abelian S_2 -subgroup, then $A_2(K) = A_\infty(K)$ and $J_\infty(K) = J_\infty(G)$.

Proof. Assume (i) is false for G . Let α be an automorphism of G , and choose $g \in G$ such that $K^\alpha = K^g$. If $J_2(K) \triangleleft G$, then

$$(J_2(K))^\alpha = J_2(K^\alpha) = J_2(K^g) = (J_2(K))^g = J_2(K).$$

Therefore $J_2(K) \text{ char } G$. Hence $J_2(K) \not\triangleleft G$.

Let L be the largest normal subgroup of G which normalizes $J_2(K)$. Then $K \cap L$ is an S_π -subgroup of L by [11], lemma 7.2 p. 444. Since $J_2(K \cap L) \text{ char } K \cap L$, it follows therefore, by a generalization of the Frattini argument that $G = LN$, where $N = N_G(J_2(K \cap L))$. If $J_2(K) \subseteq K \cap L$, then $J_2(K) = J_2(K \cap L)$. In this case $N = N_G(J_2(K))$. But then $G = LN \subseteq N_G(J_2(K))$ and $J_2(K) \triangleleft G$, a contradiction. Thus, we may assume that $J_2(K) \not\subseteq L \cap K$. By Proposition 1, $F(G) \subseteq A$ for every $A \in A_2(K) \subseteq A_\infty(K)$. In particular, $[F(G), A, A] = 1$. Since G is π -stable, it follows from the definition that $AC_G(F(G))/C_G(F(G)) \subseteq O_\pi(G/C_G F(G))$. By definition of L and by hypothesis $C_G(F(G)) \subseteq F(G) \subseteq L$. Hence $AL/L \subseteq O_\pi(G/L)$ for every $A \in A_2(K) \subseteq A_\infty(K)$.

However, we claim that $O_\pi(G/L) = 1$. Indeed, set $O_\pi(G/L) =$

T/L . $K \cap T$ is an S_π -subgroup of T , therefore $(K \cap T)L/L$ is an S_π -subgroup of T/L by [11], Lemma 7.2 p. 444. Therefore $(K \cap T)L = T$. But $K \cap T \subseteq N_G(J_2(K))$, thus $K \cap T \subseteq L$, whence $T = L$, and $O_\pi(G/L) = 1$. Therefore $A \subseteq L$ for every $A \in A_2(K) \subseteq A_x(K)$. Therefore $J_2(K) \subseteq L \cap K$, a contradiction. Therefore $J_2(K) \text{ char } G$, proving (i).

Clearly $F(G) \subseteq F(K)$ and $|F(K)|$ is odd, since $C_G(F(G)) \subseteq F(G)$. Hence by hypothesis $C_G(F(K)) \subseteq F(K)$. Proposition 1 yields that $F(K) \subseteq A$ for every $A \in A_x(K)$. Therefore $A_x(K) = A_2(K)$ as a consequence of [13], Theorem 1. Proposition 1 implies that $F(G) \subseteq A$ for every $A \in A_x(G)$. Therefore $Z(A) \subseteq C_G(F(G)) = Z(F(G))$. By hypothesis $F(G)$ is a π -group. Hence A is a π -group for every $A \in A_x(G)$. Since G is a D_π^N -group and $J_x(K) \text{ char } G$ by part (i), we obtain:

$$J_x(G) = \langle J_x(K^X) / X \in G \rangle = J_x(K),$$

proving (ii).

We now obtain:

Proof of Theorem 3. (i) We use induction on the order of G . Let $T = O_p(G)$, $H = O_{pp}(G)$, $G^* = AH$, and $K^* = A(K \cap H)$. Then $K \cap H$ is an S_π -subgroup of H and K^* is an S_π -subgroup of G^* .

Suppose that $G^* \subset G$. Since $A \subseteq K^*$, $A \in A_x(K^*)$. By induction, $O_p(A) \subseteq O_p(G^*)$. Hence

$$[H, O_p(A)] \subseteq H \cap O_p(G^*) \subseteq O_p(H) = T.$$

Therefore, $O_p(A)T/T \subseteq C_{G/T}(H/T) \subseteq H/T$, by [8], p. 228. Consequently, $O_p(A) \subseteq H$. So,

$$O_p(A) \subseteq H \cap O_p(G^*) = O_p(H) = T.$$

On the other hand, $T = O_p(G) \subseteq O_p(A)$ by Proposition 1. Therefore, $O_p(A) = O_p(G)$, as desired.

Suppose that $G^* = G$. Then $A_p T$ is an S_p -subgroup of G . By Proposition 1, $T = O_p(G) \subseteq O_p(A)$. Therefore $O_p(A)$ is an S_p -subgroup of G . It is well known that G is p -strongly solvable for every $p \in \pi - \{2\}$. By definition G is p -stable for $p = 2$. Hence G is p -stable for every $p \in \pi$. Therefore $G = O_{p', p, p}(G)$ by Proposition 2. Hence $O_p(A) \subseteq O_{p', p}(G)$. By Proposition 1, $AF(O_{p', p}(G))$ is nilpotent. Hence $O_p(A) \subseteq C_G(F(O_{p', p}(G)))$. By [3], Lemma 4, $O_p(A) \subseteq O_p(G)$, for every $p \in \pi$. Therefore $O_p(A) = O_p(G)$, as desired. In particular $O_p(A) = O_p(K)$, for every $p \in \pi$, proving (i).

(ii) Define $\sigma = \pi(F(G))$. It is well known that G is a D_σ^N -group. By [1], Corollary 4.8, $C_G(F(G)) \subseteq F(G)$. Proposition 1 yields that $F(G) \subseteq A$. Therefore $Z(A) \subseteq C_G(F(G)) \subseteq F(G)$. Hence A is a σ -group. By definition $O_\sigma(G) = 1$. Since by hypothesis G is p -strongly solvable, for every $p \in \sigma$, G is σ -stable by [2], Lemma 3.4. Let R be an S_σ -subgroup of G . Then $J_\infty(R) = J_\infty(K) = J_\infty(G)$ by Proposition 2. So $J_\infty(K) = J_\infty(G) = F(G) = F(K) = A$, by part (i).

The author knows of no counterexample to the conjecture that if G is a group of odd order then $O_p(A_2) \subseteq O_p(G)$, where $p \geq 5$ and $A_2 \in A_2(G)$.

Theorem 3 has three corollaries.

COROLLARY 1. *If the Sylow subgroups of a solvable group G are all Abelian or if G is of odd order and the Sylow subgroups of G are of class at most 2, then $F(G) \in A_\infty(G)$.*

COROLLARY 2. *If P is an S_p -subgroup of a group G , p odd, $\text{cl}(P) \leq 2$, and if $N_G(P)$ has a normal p -complement, then so does G .*

Proof. Following the method of [8], Theorem 8.3.1 and using Theorem 3 we obtain Corollary 3. One must take $N_G(P)$ instead of $N_G(ZJ(P))$ in the above mentioned theorem and its proof.

Corollary 2 is a known result. It can be obtained by [11], Theorem 8.1 p. 447.

We shall say that G is a ψ -group if $A_\infty(G)$ and Carter subgroups of G coincide.

COROLLARY 3. *Let G be a group of odd order. Assume that G and every subgroup of G is a ψ -group, then G is nilpotent.*

Proof. Let G be a minimal counterexample. By induction every proper subgroup of G is nilpotent. Therefore the Sylow subgroups of G are of class at most 2 by [11], Theorem 5.2, p. 281. Hence, Theorem 3 implies that $\langle A_\infty(G) \rangle = F(G)$ is a Carter subgroup of G . Therefore G is nilpotent as desired.

REFERENCES

1. Z. Arad, *Products of fitting classes of finite groups*, Israel J. Math., **12** (1972), 61-69.
2. ———, *A characteristic subgroup of π -stable groups*, Canad. J. Math., **XXVI**, 6 (1974), 1509-1514.
3. Z. Arad and G. Glauberman, *A characteristic subgroup of a group of odd order*, Pacific J. Math., **56** (1975), 305-319.

4. A. Bialostocki, *On products of two nilpotent subgroups of a finite group*, Israel J. Math., **20** (1975), 178–188.
5. R. W. Carter, *On a class of finite solvable groups*, Proc. London Math. Soc., (3) **9** (1959), 623–640.
6. W. Feit and J. G. Thompson, *Solvability of groups of odd order*, Pacific J. Math., **13** (1963), 775–1029.
7. G. Glauberman, *On Burnside's other $p^a q^b$ Theorem*, Pacific J. Math., **56** (1975), 469–476.
8. D. Gorenstein, *Finite Groups*, Harper and Row, New York, (1968).
9. M. Hall, *The Theory of Groups*, New York: Macmillan Company (1959).
10. P. Hall, *On the system normalizers of a solvable group*, Proc. London Math. Soc., (2) **43** (1937), 507–508.
11. B. Huppert, *Endlich Gruppen I*, Springer-Verlag, New York (1967).
12. N. Ito, *Über der Kleinsten p -Durchschnitt auflösbarer Gruppen*, Arch. Math., **9** (1958), 27–32.
13. A. Mann, *Injectors and normal subgroups of finite groups*, Israel J. Math., **9**, No. 4, (1971), 554–558.
14. W. R. Scott, *Group Theory*, Englewood Cliffs (New-Jersey) Prentice-Hall, Inc. (1964).

Received September 26, 1974 and in revised form October 30, 1975.

BAR-ILAN UNIVERSITY

Pacific Journal of Mathematics

Vol. 62, No. 1

January, 1976

Mieczyslaw Altman, <i>Contractor directions, directional contractors and directional contractions for solving equations</i>	1
Michael Peter Anderson, <i>Subgroups of finite index in profinite groups</i>	19
Zvi Arad, <i>Abelian and nilpotent subgroups of maximal order of groups of odd order</i>	29
John David Baidon and Ruth Silverman, <i>On starshaped sets and Helly-type theorems</i>	37
John W. Baker and R. C. Lacher, <i>Some mappings which do not admit an averaging operator</i>	43
Joseph Barback, <i>Composite numbers and prime regressive isols</i>	49
David M. Boyd, <i>Composition operators on $H^p(A)$</i>	55
Maurice Chacron, <i>Co-radical extension of PI rings</i>	61
Fred D. Crary, <i>Some new engulfing theorems</i>	65
Victor Dannon and Dany Leviatan, <i>A representation theorem for convolution transform with determining function in L^p</i>	81
Mahlon M. Day, <i>Lumpy subsets in left-amenable locally compact semigroups</i>	87
Michael A. Gauger, <i>Some remarks on the center of the universal enveloping algebra of a classical simple Lie algebra</i>	93
David K. Haley, <i>Equational compactness and compact topologies in rings satisfying A.C.C</i>	99
Raymond Heitmann, <i>Generating ideals in Prüfer domains</i>	117
Gerald Norman Hile, <i>Entire solutions of linear elliptic equations with Laplacian principal part</i>	127
Richard Oscar Hill, <i>Moore-Postnikov towers for fibrations in which $\pi_1(\text{fiber})$ is non-abelian</i>	141
John Rast Hubbard, <i>Approximation of compact homogeneous maps</i>	149
Russell L. Merris, <i>Relations among generalized matrix functions</i>	153
V. S. Ramamurthi and Edgar Andrews Rutter, <i>On cotorsion radicals</i>	163
Ralph Tyrrell Rockafellar and Roger Jean-Baptiste Robert Wets, <i>Stochastic convex programming: basic duality</i>	173
Alban J. Roques, <i>Local evolution systems in general Banach spaces</i>	197
I. Bert Russak, <i>An indirect sufficiency proof for problems with bounded state variables</i>	219
Richard Alexander Sanerib, Jr., <i>Ultrafilters and the basis property</i>	255
H. A. Seid, <i>The decomposition of multiplication operators on L_p-spaces</i>	265
Franklin D. Tall, <i>The density topology</i>	275
John Campbell Wells, <i>Invariant manifolds on non-linear operators</i>	285
James Chin-Sze Wong, <i>A characterization of topological left thick subsets in locally compact left amenable semigroups</i>	295