

Pacific Journal of Mathematics

**SOME REMARKS ON THE CENTER OF THE UNIVERSAL
ENVELOPING ALGEBRA OF A CLASSICAL SIMPLE LIE
ALGEBRA**

MICHAEL A. GAUGER

SOME REMARKS ON THE CENTER OF THE UNIVERSAL ENVELOPING ALGEBRA OF A CLASSICAL SIMPLE LIE ALGEBRA

MICHAEL A. GAUGER

This paper is concerned with explicitly producing generating sets of the centers of the universal enveloping algebras of classical simple Lie algebras.

Let L be a finite-dimensional simple Lie algebra over an algebraically closed field K of characteristic zero, let U be its universal enveloping algebra, and let Z be the center of U . If l is the dimension of a Cartan subalgebra H of L , then it is known that Z is a polynomial ring in l independent variables. In this paper a set of l algebraically independent generators of Z is produced rather explicitly for the classical algebras of type A, B, C, D by casewise considerations.

It is straightforward to show that generating Z is equivalent to generating the L -invariants I_L^* in the symmetric algebra S_L of L^* . In addition, there is a homomorphism from S_L onto S_H which embeds I_L^* into the Weyl-invariants I_W . Due to Chevalley this embedding is also a surjection. For the classical simple Lie algebras the action of the Weyl group W on S_H is describable in a sufficiently convenient fashion so as to permit easy construction of generators of I_W . It is shown here that certain generating sets of I_W can be explicitly lifted back to I_L^* via trace functions on the first fundamental representation of L . As a result of this construction of the generators of I_W and the lifting process, the following well-known results are proven rather directly for the classical algebras:

1. $I_L^* \cong I_W$ (Chevalley), and
2. Z and I_W are polynomial rings in l algebraically independent variables.

The center Z of U plays a fundamental role in the finite-dimensional representation theory of L . Since any irreducible representation is determined up to isomorphism by its character, if z_1, \dots, z_l were generators of Z and if M and N were non-isomorphic irreducible L -modules, then for some i one must have $(z_i)_M \neq (z_i)_N$ (due to Schur's lemma they are scalars). The central element $(z_i - (z_i)_N) / ((z_i)_M - (z_i)_N)$ would act as one on M and zero on N . For any list of pairwise non-isomorphic irreducible L -modules one could thus find a central element acting as one on one of them, and as zero on the rest. Such elements could be used to isolate the isotypic components in a reducible

representation of L . Hence there is good reason to produce generators of Z as explicitly as possible.

Section 1 is concerned with showing that generating Z is equivalent to generating I_L^* and leads up to §§2–5 where the Chevalley isomorphism $I_L^* \cong I_w$ is proven by explicitly lifting generating sets of I_w back to I_L^* .

1. Generation of Z . There are well known actions of L on the symmetric algebras S_L and S_{L^*} by graded derivations extending the adjoint representation of L and its contragredient, and if W is the Weyl group of L with respect to the Cartan subalgebra H , it acts on S_H by graded automorphisms. The standard symmetrization map $\eta: S_L \rightarrow U$ given by $(x_1 \cdots x_r)^\eta = (1/r!) \sum_{\alpha \in S} x_{\alpha(1)} \cdots x_{\alpha(r)}$ for a monomial of degree r in S_L , induces a linear isomorphism between the L -invariants I_L in S_L and the L -invariants Z in U since it is an L -module isomorphism. While this induced map between I_L and Z is not an algebra isomorphism it is known to have the following redeeming qualities:

LEMMA. *Suppose S is a finite set of homogeneous invariant elements in S_L generating I_L . Then S^η generates Z , and if S is algebraically independent so is S^η .*

Proof. Let U have its usual filtration and let U_p be the subspace of all elements of filter less than or equal to p . Observe that due to the Poincaré-Birkhoff-Witt theorem, if x_1, \dots, x_r are homogeneous elements of S_L of degrees d_1, \dots, d_r and $d = \sum_i d_i$, then $(x_1 \cdots x_r)^\eta = x_1^\eta \cdots x_r^\eta + t$ where t is in U_{d-1} .

(i) Since L acts by graded derivations, I_L is homogeneous. Recalling that η induces a linear isomorphism between I_L and Z , proceed by induction on the filter of a central element to show it is in the subalgebra of U generated by S^η . Let $S = \{x_1, \dots, x_r\}$. Now η takes constants to constants so it suffices to check the induction step. Every element in Z is a linear combination of images of homogeneous elements in I_L , so it suffices to show that if $x_{i_1} \cdots x_{i_k}$ is a monomial in I_L then $(x_{i_1} \cdots x_{i_k})^\eta$ is in the subalgebra generated by S^η . The remarks in the first paragraph complete the proof.

(ii) Set $y_i = x_i^\eta$. Suppose the y_i are algebraically dependent and let p be a nonzero polynomial in $K[Y_1, \dots, Y_r]$ such that $p(y_1, \dots, y_r) = 0$. Write $p = q + t$ where q is the homogeneous part of p of highest total degree d . Since η takes $q(x_1, \dots, x_r)$ onto $q(y_1, \dots, y_r)$ plus an element $u(y_1, \dots, y_r)$ whose filter is less than d , there is a polynomial h of degree less than d such that η takes $h(x_1, \dots, x_r)$ onto $t(y_1, \dots, y_r) - u(y_1, \dots, y_r)$. Since η is an isomorphism $(q + h)(x_1, \dots, x_r) = 0$. This contradicts the independence of the x_i since $q + h \neq 0$.

Since the Killing form of L is nondegenerate there is an induced

isomorphism between L and L^* which extends to an L -module algebra isomorphism between S_L and S_{L^*} . Hence there is an induced algebra isomorphism between I_L and I_L^* . Viewing S_{L^*} as the ring of polynomial functions on L , one gets by restriction to H an epimorphism $\rho: S_{L^*} \rightarrow S_H$ which injects I_L^* into I_W ([2], 126). The remainder is concerned with producing algebraically independent generating sets of I_W and exhibiting how they lift back to I_L^* . Chevalley's isomorphism ($I_L^* \cong I_W$) is thus proven as well as the theorems that Z and I_W are polynomial rings in l independent variables.

2. Simple algebras of type A. Let L be simple of type A_l . View L as the Lie algebra of trace zero endomorphisms of $V = K^{l+1}$, and identify L with its matrices with respect to standard basis vectors e_1, \dots, e_{l+1} . Let H be the Cartan subalgebra of diagonal matrices of trace zero and let $\epsilon_1, \dots, \epsilon_{l+1}$ be functionals on H given by $(e_i)h = \epsilon_i(h)e_i$ for h in H . Then the ϵ_j generate H^* , $\sum_j \epsilon_j = 0$, and W acts as the symmetric group on the ϵ_j . ([3], 136 and [1] 205–207, 250–251). Let A be an $l+1$ -dimensional auxiliary space with basis $\bar{\epsilon}_1, \dots, \bar{\epsilon}_{l+1}$ on which W acts as the symmetric group. There is a W -epimorphism $A \rightarrow H^*$ taking $\bar{\epsilon}_i$ to ϵ_i which extends to a W -epimorphism $S_A \rightarrow S_{H^*}$. Hence there is an induced epimorphism $\bar{I}_W \rightarrow I_W$ where \bar{I}_W is the set of W -invariants in S_A . Now \bar{I}_W is generated by the (algebraically independent) elementary symmetric functions in $\bar{\epsilon}_1, \dots, \bar{\epsilon}_{l+1}$. The kernel of $\bar{I}_W \rightarrow I_W$ is easily seen to be generated by $\sum_j \bar{\epsilon}_j$, thus I_W is generated by the algebraically independent elementary symmetric functions s_2, \dots, s_{l+1} in $\epsilon_1, \dots, \epsilon_{l+1}$ — the analysis being identical to the situation $K[X_1, \dots, X_{l+1}] \rightarrow K[X_1, \dots, X_l]$ where X_{l+1} goes to zero. Unfortunately the symmetric functions do not lift easily. Due to Newton's identities, however, $I_W = K[p_2, \dots, p_{l+1}]$ where $p_i = \epsilon_1^i + \dots + \epsilon_{l+1}^i$ and the p_i do lift easily. They are algebraically independent since they generate a ring known to have transcendence degree equal to l . Now let F_i in I_L^* be given by $F_i(x) = \text{tr}(x_V)^i$. Then $F_i^? = p_i$ and $\rho: I_L^* \rightarrow I_W$ is surjective. Under the isomorphisms $Z \cong I_L \cong I_L^*$ the element z_k of Z corresponding to F_k is given by

$$(1) \quad z_k = \sum_{i_1, \dots, i_k=1}^n \text{tr}(u_{i_1} \cdots u_{i_k})_V u^{i_1} \cdots u^{i_k}$$

where $\{u_i\}, \{u^i\}$ are dual bases of L with respect to its Killing form. By Lemma 1 and the discussion in §1 $Z = K[z_2, \dots, z_{l+1}]$ and the z_k are algebraically independent.

3. Simple algebras of type B. Let L be a simple algebra of type B_l . Let V be a $(2l+1)$ -dimensional space with basis e_1, \dots, e_{2l+1} ,

and define a non-degenerate symmetric form on V by $B(e_1, e_1) = 1 = B(e_i, e_{i+l}) = B(e_{i+l}, e_i)$ $i = 2, \dots, l+1$ and $B(e_j, e_k) = 0$ otherwise. View L as the Lie algebra of all endomorphisms of V which are skew with respect to this form and identify L with its matrices with respect to the e_i . Let H be the Cartan subalgebra of diagonal matrices in L , and let $\epsilon_1, \dots, \epsilon_l$ be functionals on H given by $(e_i)h = \epsilon_i(h)e_i$ for h in H ([3], 138). Then $\{\epsilon_k\}_k$ is a basis of H^* and W is the semidirect product of the symmetric group S_l on $\epsilon_1, \dots, \epsilon_l$ with $(\mathbf{Z}/2\mathbf{Z})^l$ acting by $\epsilon_i \rightarrow (\pm 1)_i \epsilon_i$. Thus I_W consists of symmetric functions in $\epsilon_1^2, \dots, \epsilon_l^2$ ([2], 202 and 252). By Newton's identities $I_W = k[p_1, \dots, p_l]$ where $p_i = \epsilon_1^{2i} + \dots + \epsilon_l^{2i}$. Since $\epsilon_1^2, \dots, \epsilon_l^2$ are algebraically independent, so are the p_i . Let F_i in I_L^* be given by $F_i(x) = \text{tr}(x_V)^{2i}$. Then $F_i^\rho = 2p_i$ and ρ is onto. $Z = K[z_2, z_4, \dots, z_{2l}]$ where the z_{2k} are as in (1).

4. Simple algebras of type C. Let L be simple of type C_l . Let V be a $2l$ -dimensional space with basis e_1, \dots, e_{2l} , and define a nondegenerate skew form on V by $B(e_i, e_{i+l}) = 1 = -B(e_{i+l}, e_i)$ $i = 1, \dots, l$ and $B(e_j, e_k) = 0$ otherwise. View L as the Lie algebra of endomorphisms which are skew with respect to this form, and identify L with its matrices with respect to the e_i . Let H be the Cartan subalgebra of diagonal matrices in L , and let $\epsilon_1, \dots, \epsilon_l$ be functionals on H given by $(e_i)h = \epsilon_i(h)e_i$ when h is in H ([3], 139). Then $\epsilon_1, \dots, \epsilon_l$ is a basis of H^* , W acts just as in the preceding case, and I_W consists of symmetric functions in $\epsilon_1^2, \dots, \epsilon_l^2$ ([2], 204 & 254). As before one sees ρ is onto and $Z = K[z_2, \dots, z_{2l}]$ where z_{2k} is as in (1).

5. Simple algebras of type D. Let L be simple of type D_l . Let V be a $2l$ -dimensional space with basis e_1, \dots, e_{2l} , and define a nondegenerate symmetric form on V by $B(e_i, e_{i+l}) = 1 = B(e_{i+l}, e_i)$ $i = 1, \dots, l$ and $B(e_j, e_k) = 0$ otherwise. View L as the Lie algebra of endomorphisms of V which are skew with respect to this form and identify L with its matrices with respect to the e_j . Let H be the Cartan subalgebra of diagonal matrices in L and let $\epsilon_1, \dots, \epsilon_l$ be functionals on H given by $(e_i)h = \epsilon_i(h)e_i$ when h is in H ([3], 140). Then $\epsilon_1, \dots, \epsilon_l$ is a basis of H^* and W is the semi-direct product of the symmetric group S_l acting as before with $(\mathbf{Z}/2\mathbf{Z})^{l-1}$ acting by $\epsilon_i \rightarrow (\pm 1)_i \epsilon_i$ where $\prod_i (\pm 1)_i = 1$ ([2], 208 and 256). Thus I_W consists of polynomials in the elementary symmetric functions in $\epsilon_1^2, \dots, \epsilon_l^2$ and the function $\epsilon_1 \dots \epsilon_l$. Let s_k be the k th elementary symmetric function in the ϵ_i^2 and let $t = \epsilon_1 \dots \epsilon_l$. Since $s_l = t^2$, one has $I_W = K[s_1, \dots, s_{l-1}, t]$. If s_1, \dots, s_{l-1}, t were algebraically dependent, by an even-odd degree argument there would be a relation in which every monomial has t to an even power, or every monomial has t to an odd power. If the relation is of the second type multiply it by t to make it of the first type. But a relation of the first type is impossible

since the elementary symmetric function in the ϵ_i^2 are algebraically independent. Thus I_w is a polynomial ring in l independent variables. By Newton's identities $I_w = K[p_1, \dots, p_{l-1}, t]$ where $p_i = \epsilon_1^{2i} + \dots + \epsilon_l^{2i}$. These generators are also algebraically independent since there are l of them in a ring known to have transcendence degree equal to l . As before $2p_i$ lifts back to I_L^* as $\text{tr}(\)_V^{2i}$, and it is easy to check that $t = \epsilon_1 \cdots \epsilon_l$ lifts back to I_L^* as $\text{pf}(\)_V$ — the pfaffian. Thus ρ is onto and $Z = K[z_2, z_4, \dots, z_{2l-2}, w]$ where the z_{2k} are as in (1) and w corresponds to $\text{pf}(\)_V$ under $Z \cong I_L \cong I_L^*$.

REMARK. Dual bases of L with respect to its Killing Form can be explicitly constructed ([3], 246; $h^i = h_{\lambda_i}$ where the λ_i are the fundamental weights). According to part VI of Planche I–IV ([1], 250–258) the coefficients q_{ij} of the equations $\lambda_i = \sum_j q_{ij} \alpha_j$ ($\alpha_1, \dots, \alpha_l$ a simple root system) are known, thus enabling one to express h^i explicitly as a \mathbf{Q} -linear combination of the h_i .

REFERENCES

1. N. Bourbaki, *Elements de Mathématique*, XXXIV Groupes et algèbres de Lie. Ch. 6: Systèmes de racines, Actualités Sci. Indust., no. 1337, Hermann, Paris, 1968.
2. J. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag, N.Y. 1972.
3. N. Jacobson, *Lie Algebras*, Interscience Tracts in Pure and Applied Math., no. 10, Interscience, New York, 1962.

Received May 15, 1974. The research presented here was supported in part by the National Science Foundation under grant GP-43814.

UNIVERSITY OF MASSACHUSETTS

Pacific Journal of Mathematics

Vol. 62, No. 1

January, 1976

Mieczyslaw Altman, <i>Contractor directions, directional contractors and directional contractions for solving equations</i>	1
Michael Peter Anderson, <i>Subgroups of finite index in profinite groups</i>	19
Zvi Arad, <i>Abelian and nilpotent subgroups of maximal order of groups of odd order</i>	29
John David Baidon and Ruth Silverman, <i>On starshaped sets and Helly-type theorems</i>	37
John W. Baker and R. C. Lacher, <i>Some mappings which do not admit an averaging operator</i>	43
Joseph Barback, <i>Composite numbers and prime regressive isols</i>	49
David M. Boyd, <i>Composition operators on $H^p(A)$</i>	55
Maurice Chacron, <i>Co-radical extension of PI rings</i>	61
Fred D. Crary, <i>Some new engulfing theorems</i>	65
Victor Dannon and Dany Leviatan, <i>A representation theorem for convolution transform with determining function in L^p</i>	81
Mahlon M. Day, <i>Lumpy subsets in left-amenable locally compact semigroups</i>	87
Michael A. Gauger, <i>Some remarks on the center of the universal enveloping algebra of a classical simple Lie algebra</i>	93
David K. Haley, <i>Equational compactness and compact topologies in rings satisfying A.C.C</i>	99
Raymond Heitmann, <i>Generating ideals in Prüfer domains</i>	117
Gerald Norman Hile, <i>Entire solutions of linear elliptic equations with Laplacian principal part</i>	127
Richard Oscar Hill, <i>Moore-Postnikov towers for fibrations in which $\pi_1(\text{fiber})$ is non-abelian</i>	141
John Rast Hubbard, <i>Approximation of compact homogeneous maps</i>	149
Russell L. Merris, <i>Relations among generalized matrix functions</i>	153
V. S. Ramamurthi and Edgar Andrews Rutter, <i>On cotorsion radicals</i>	163
Ralph Tyrrell Rockafellar and Roger Jean-Baptiste Robert Wets, <i>Stochastic convex programming: basic duality</i>	173
Alban J. Roques, <i>Local evolution systems in general Banach spaces</i>	197
I. Bert Russak, <i>An indirect sufficiency proof for problems with bounded state variables</i>	219
Richard Alexander Sanerib, Jr., <i>Ultrafilters and the basis property</i>	255
H. A. Seid, <i>The decomposition of multiplication operators on L_p-spaces</i>	265
Franklin D. Tall, <i>The density topology</i>	275
John Campbell Wells, <i>Invariant manifolds on non-linear operators</i>	285
James Chin-Sze Wong, <i>A characterization of topological left thick subsets in locally compact left amenable semigroups</i>	295