Pacific Journal of Mathematics

SUBSPACES OF SYMMETRIC MATRICES CONTAINING MATRICES WITH A MULTIPLE FIRST EIGENVALUE

SHMUEL FRIEDLAND AND RAPHAEL LOEWY

SUBSPACES OF SYMMETRIC MATRICES CONTAINING MATRICES WITH A MULTIPLE FIRST EIGENVALUE

S. FRIEDLAND AND R. LOEWY

Let \mathcal{U} be an (r-1)(2n-r+2)/2 dimensional subspace of $n \times n$ real valued symmetric matrices. Then \mathcal{U} contains a nonzero matrix whose greatest eigenvalue is at least of multiplicity r, if $2 \leq r \leq n-1$. This bound is best possible. We apply this result to prove the Bohnenblust generalization of Calabi's theorem. We extend these results to hermitian matrices.

1. Introduction. Let \mathcal{W}_n be the n(n+1)/2 dimensional vector space of all real valued $n \times n$ symmetric matrices. Let A belong to \mathcal{W}_n . Arrange the eigenvalues of A in decreasing order

(1.1)
$$\lambda_1(A) \geq \lambda_2(A) \geq \cdots \geq \lambda_n(A).$$

We say that $\lambda_1(A)$ is of multiplicity r if

(1.2a)
$$\lambda_1(A) = \cdots = \lambda_r(A),$$

(1.2b)
$$\lambda_r(A) > \lambda_{r+1}(A).$$

Let \mathcal{U} be a subspace of \mathcal{W}_n of dimension k. We consider the question of how large k has to be so that \mathcal{U} must contain a nonzero matrix A which satisfies (1.2a) for a given r. The nontrivial case would be

$$(1.3) 2 \leq r \leq n-1.$$

Clearly for r = n we must have k = n(n + 1)/2 as \mathcal{U} will contain the identity matrix *I*.

We now state our main result:

THEOREM 1. Let \mathcal{U} be a k dimensional subspace in the space \mathcal{W}_n of $n \times n$ real valued matrices. Assume that an integer r satisfies the inequalities (1.3).

If

(1.4)
$$k \ge \kappa(r)$$

where

(1.5)
$$\kappa(r) = (r-1)(2n-r+2)/2, \qquad r = 1, 2, \cdots, n$$

then \mathcal{U} contains a nonzero matrix A such that the greatest eigenvalue of A is at least of multiplicity r. The lower bound $\kappa(r)$ is best possible for $2 \leq r \leq n-1$.

Theorem 1 is proved in §2. In §3 we prove that Theorem 1 is equivalent to the following result due to Bohnenblust (cf. [1] and [4]). We denote as usual by (x, y) the inner product of the vectors x and y in \mathbb{R}^n , which is the underlying vector space for \mathcal{W}_n .

THEOREM 2 (Bohnenblust). Let \mathcal{V} be a subspace of dimension k in \mathcal{W}_n and let $1 \leq r \leq n-1$. Assume that \mathcal{V} has the following property:

(1.6)
$$\sum_{i=1}^{r} (Ax_i, x_i) = 0 \text{ for every } A \text{ in } \mathcal{V}$$

implies that $x_i = 0$ for $i = 1, \dots, r$. If

(1.7)
$$k < f(r+1) - \delta_{n,r+1},$$

where

(1.8)
$$f(r) = r(r+1)/2,$$

then \mathcal{V} contains a positive definite matrix.

In case r = 1, Bohnenblust's result reduces to the following theorem, known as the *Calabi theorem* [2]: Let $n \ge 3$ and suppose that S_1 and S_2 are $n \times n$ symmetric matrices such that $(S_1x, x) = (S_2x, x) = 0$ implies x = 0. Then there exist real α_1 and α_2 such that $\alpha_1S_1 + \alpha_2S_2$ is positive definite.

Bohnenblust defines a subspace \mathcal{V} with the property:

(1.9)
$$\sum_{i=1}^{r} (Ax_i, x_i) = 0$$
 for every $A \neq 0$ in \mathcal{V} implies $x_1 = x_2 = \cdots = x_r = 0$

to be *jointly definite of degree r*. Thus, the equivalence of Theorems 1 and 2 relates the notion of a subspace which is jointly definite of degree r with that of a subspace containing a nonzero matrix whose largest eigenvalue has multiplicity r.

Finally, in §4 we prove that if we let \mathcal{W}_n be the n^2 dimensional *real* space of all $n \times n$ hermitian matrices then Theorems 1 and 2 remain correct if $\kappa(r)$ and f(r) are defined as follows

390

(1.10)
$$\kappa(r) = (r-1)(2n-r+1),$$

(1.11)
$$f(r) = r^2$$
.

2. *Proof of Theorem* 1. We first establish a weaker form of Theorem 1 which will be needed for the proof of Theorem 1.

LEMMA 1. Let $1 \leq r \leq n$. Let \mathcal{U} be a k-dimensional subspace of \mathcal{W}_n and assume that

$$(2.1) k \ge 1 + \kappa(r).$$

Then there exists A in U such that

(2.2)
$$\lambda_1(A) = \cdots = \lambda_r(A) = 1.$$

Proof. For r = 1 (2.2) trivially holds. For r = n (2.2) is also obvious as $1 + \kappa (n) = n(n + 1)/2$. Suppose that the lemma holds for r = p. Next we construct A which satisfies (2.2) for r = p + 1. Let B^* satisfy

(2.3)
$$\lambda_1(B^*) = \cdots = \lambda_p (B^*) = 1, \qquad (p \ge 1).$$

The existence of B^* follows from our assumptions. Assume that

(2.4)
$$1 > \lambda_{p+1}(B^*).$$

Otherwise B^* would satisfy (2.2) for r = p + 1. Let

$$(2.5) B^*\xi_i = \lambda_i (B^*)\xi_i ; (\xi_i, \xi_j) = \delta_{ij}, i, j = 1, \cdots, n.$$

Suppose that A_1, \dots, A_k form a basis for \mathcal{U} . Consider the system

(2.6)
$$\sum_{j=1}^{k} \alpha_j A_j \xi_i = 0, \qquad i = 1, \cdots, p.$$

We claim that (2.6) is equivalent to $\kappa(p+1) = \kappa(r)$ scalar equations. Indeed, we can assume $[\xi_1, \dots, \xi_n]$ to be the standard basis in \mathbb{R}^n . Then each A_i is represented by an appropriate $n \times n$ symmetric matrix

(2.7)
$$A_i = (a_{\mu\nu}^i), \qquad i = 1, \cdots, k.$$

So (2.6) is equivalent to

(2.8a)
$$\sum_{j=1}^{k} \alpha_{j} a_{\mu\mu} = 0, \qquad \mu = 1, \cdots, p,$$

(2.8b)
$$\sum_{j=1}^{k} \alpha_{j} \dot{a}_{\mu\nu}^{j} = 0, \quad \mu = 1, \cdots, p; \ \nu = \mu + 1, \cdots, n.$$

Clearly (2.8a) and (2.8b) are a system of $\kappa (p+1) = p(2n-p+1)/2$ linear equations in the unknowns $\alpha_1, \dots, \alpha_k$. As $k \ge 1 + \kappa (p+1)$ we have a nontrivial solution of (2.6). Hence there exists $C \ne 0$ in \mathcal{U} such that

$$(2.9) C\xi_i = 0, i = 1, \cdots, p.$$

We can assume that

$$(2.10) \qquad \qquad \lambda_1(C) > 0.$$

(Otherwise take -C). Consider the matrix

(2.11)
$$C(\alpha) = B^* + \alpha C.$$

Clearly, (2.3), (2.4) and (2.9) imply for $|\alpha|$ small enough

(2.12a)
$$\lambda_1(C(\alpha)) = \cdots = \lambda_p(C(\alpha)) = 1,$$

(2.12b) $1 > \lambda_{p+1}(C(\alpha)).$

We claim that there exists α^* such that

(2.13)
$$\lambda_1(C(\alpha^*)) = \cdots = \lambda_{p+1}(C(\alpha^*)) = 1.$$

Otherwise we must have for all $\alpha > 0$ the conditions (2.12). But for a large positive α we have that $\lambda_1(C(\alpha)) = \alpha \lambda_1(C) + O(1)$. This contradicts (2.12a). Thus (2.13) holds. End of proof.

Thus, Theorem 1 shows that if we relax the condition that the largest eigenvalue of $A \neq 0$ of multiplicity r would be distinct from zero then for $2 \leq r \leq n-1$ the bound (2.1) can be reduced by 1. We will show later that the bound $\kappa(r) + 1$ is sharp.

LEMMA 2. Let $2 \leq r \leq n$. Let \mathcal{U} be a k-dimensional subspace of \mathcal{W}_n and suppose that $k \geq \kappa(r)$. Assume that for any nonzero A in \mathcal{U} we have

(2.14)
$$\lambda_1(A) > \lambda_r(A).$$

392

Let $\eta_1, \eta_2, \dots, \eta_{r-1}$ be a set of r-1 arbitrary orthonormal vectors. Consider the system

$$(2.15) A\eta_i = \lambda\eta_i, \quad i = 1, 2, \cdots, r-1, \quad and \quad A \in \mathcal{U}.$$

Then there exists a nonzero matrix A_0 in \mathcal{U} and a scalar λ_0 such that

$$(2.16) A_0 \eta_i = \lambda_0 \eta_i, i = 1, 2, \cdots, r-1,$$

and

(2.17)
$$\lambda_0 = \lambda_1(A_0) = \cdots = \lambda_{r-1}(A_0).$$

Moreover, for any pair A and λ , where A belongs to \mathcal{U} , that satisfies (2.15), there exists α such that

 $A = \alpha A_0$ and $\lambda = \alpha \lambda_0$.

Proof. From Lemma 1 we deduce the existence of $B^* \neq 0$ in \mathcal{U} such that $\lambda_1(B^*) = \lambda_{r-1}(B^*) = 1$. Let ξ_1, \dots, ξ_{r-1} be r-1 orthonormal vectors corresponding to 1. We first prove the lemma in case that $\eta_i = \xi_i$, $i = 1, \dots, r-1$. Suppose that there exists a matrix C in \mathcal{U} , linearly independent of B^* , such that $C\xi_i = \mu\xi_i$, $i = 1, \dots, r-1$. We may assume that $\mu = 0$, for otherwise replace C by $C - \mu B^*$. As in the proof of Lemma 1 we define $C(\alpha) = B^* + \alpha C$ and may conclude that there exists α^* such that $\lambda_1(C(\alpha^*)) = \lambda_r(C(\alpha^*))$ holds. This contradicts (2.14). Thus $C = \beta B^*$ and since $\mu = 0$ we must have that $\beta = 0$. So for $\eta_i = \xi_i$, $i = 1, \dots, r-1$ the lemma is proved.

Now let $\eta_1, \dots, \eta_{r-1}$ be r-1 arbitrary orthonormal vectors. Since r-1 < n it is easy to show that there exists a system $\xi_1(t), \dots, \xi_{r-1}(t)$ of r-1 orthonormal vectors for $0 \le t \le 1$ which depends continuously on t and

(2.18)
$$\xi_i(0) = \xi_i, \quad \xi_i(1) = \eta_i, \qquad i = 1, \cdots, r-1.$$

For any t, $0 \le t \le 1$, consider now the system

(2.19)
$$A\xi_i(t) = \lambda\xi_i(t), \quad i = 1, \dots, r-1, \text{ and } A \in \mathcal{U}.$$

As was shown in the proof of Lemma 1, this system is equivalent to $\kappa(r)$ linear equations. The number of variables is k + 1, namely $\alpha_1, \dots, \alpha_k, \lambda$ where $A = \sum_{i=1}^k \alpha_i A_i$ and k is the dimension of $\mathcal{U}(A_1, A_2, \dots, A_k$ form a basis for \mathcal{U}). The assumption $k \ge \kappa(r)$ implies the existence of a nontrivial solution of (2.19). Clearly, if A = 0 then $\lambda = 0$, so we always have a nontrivial solution with respect to $\alpha_1, \dots, \alpha_k$. For t = 0 it follows from (2.18) that the system (2.19) has rank $\kappa(r)$, whence $k = \kappa(r)$. Thus for $0 \le t \le \epsilon$ ($\epsilon > 0$) we would always have, up to scalar multiples, exactly one nontrivial solution A(t) in \mathcal{U} such that

(2.20)
$$A(t)\xi_{i}(t) = \lambda(t)\xi_{i}(t), \qquad i = 1, \cdots, r-1.$$

We can choose A(t) to be dependent continuously on t as long as the rank of the system (2.19) is $\kappa(r)$. Without any restriction we may assume that ||A(t)|| = 1 for some matrix norm on \mathcal{W}_n . Since $\lambda(0) = \lambda_1(A(0)) = \cdots = \lambda_{r-1}(A(0))$, the continuity of A(t) for $0 \le t \le \epsilon$ and the assumption (2.14) imply

(2.21)
$$\lambda_1(A(t)) = \lambda(t)$$

for $0 \le t \le \epsilon$. Suppose to the contrary that (2.15) has at least two linearly independent solutions. Let $0 < t_0 \le 1$ be the first time that the system (2.19) has two linearly independent solutions. Thus A(t) is continuous for $0 \le t < t_0$. Now (2.21) together with the assumption ||A(t)|| = 1 implies the existence of $B \ne 0$ in \mathcal{U} such that

$$(2.22) B\xi_i(t_0) = \lambda_0\xi_i(t_0), i = 1, \cdots, r-1,$$

and $\lambda_0 = \lambda_1(B) = \cdots = \lambda_{r-1}(B)$. The condition (2.14) implies that $\lambda_1(B) > \lambda_r(B)$. By assumption we must have a solution C in \mathcal{U} , linearly independent of B, such that

(2.23)
$$C\xi_i(t_0) = \mu\xi_i(t_0), \qquad i = 1, \cdots, r-1.$$

If $\mu = 0$ then, as in the proof of Lemma 1, we deduce that there exists α^* such that $\lambda_1(C(\alpha^*)) = \lambda_r(C(\alpha^*))$, where $C(\alpha) = B + \alpha C$. If $\mu \neq 0$ let $B_1 = C(\alpha_1)$ where α_1 is chosen to be small enough such that $\lambda_1(B_1) > \lambda_r(B_1)$ and $\lambda_1(B_1) \neq 0$. Then as in the proof of Lemma 1 we may assume that $\mu = 0$ and we again have the equality $\lambda_1(C(\alpha^*)) = \lambda_r(C(\alpha^*))$. This contradicts (2.14). The proof is complete.

Proof of Theorem 1. Let $2 \le r \le n-1$. Assume to the contrary that any $A \ne 0$ in \mathcal{U} satisfies the inequality (2.14). We then deduce the existence of a nonzero matrix in \mathcal{U} such that

(2.24)
$$\lambda_1(C) > \lambda_2(C) = \cdots = \lambda_r(C) > \lambda_n(C).$$

For r = 2 the condition (2.14) implies (2.24) for any $C \neq 0$. Let $3 \leq r \leq n-1$. Consider again the matrix B^* which satisfies $\lambda_1(B^*) = \cdots = \lambda_{r-1}(B^*) = 1$. Let ξ_1, \dots, ξ_{r-1} be r-1 corresponding orthonormal

eigenvectors. Let \mathcal{U}' be a $\kappa(r) - 1$ dimensional subspace of \mathcal{U} which does not contain B^* . Consider the equation

(2.25)
$$C\xi_i = 0, \quad i = 2, \cdots, r-1 \text{ and } C \in \mathcal{U}'.$$

Since U' is $\kappa(r) - 1$ dimensional, (2.25) is equivalent to a linear system of $\kappa(r-1)$ equations in $\kappa(r) - 1$ unknowns. Since we assumed that $3 \le r \le n-1$ it follows that $\kappa(r) - 1 > \kappa(r-1)$, whence there exists a nonzero solution C of (2.25).

If $\lambda_2(C) = \cdots = \lambda_{n-1}(C) = 0$ then (2.24) clearly holds. Hence we may assume that $\lambda_1(C) \ge \lambda_2(C) > 0$, and let $C(\alpha) = B^* + \alpha C$. It follows from (2.25) that $\lambda_1(B^*)$ is an eigenvalue of $C(\alpha)$ of multiplicity r - 2 at least, for any α . But for α sufficiently large $\lambda_1(C(\alpha)) > \lambda_1(B^*)$ and $\lambda_2(C(\alpha)) > \lambda_1(B^*)$. Define

$$T = \{ \alpha : \alpha \ge 0, \lambda_1(C(\alpha)) > \lambda_1(B^*) \text{ and } \lambda_2(C(\alpha)) > \lambda_1(B^*) \}.$$

T is not empty, so define $\gamma = \inf\{\alpha : \alpha \in T\}$. We must have $\gamma > 0$, because of (2.14). The matrix $C(\gamma)$ satisfies (2.24).

Finally, we show that (2.14) leads to a contradiction. Let C be a matrix that satisfies (2.24). Let $\eta_1, \eta_2, \dots, \eta_{r-1}$ be r-1 orthonormal eigenvectors corresponding to $\lambda_2(C) = \dots = \lambda_r(C)$. By Lemma 2, there exists a matrix A in $\mathcal{U}, A \neq 0$, such that $\lambda_1(A) = \lambda_{r-1}(A)$ and $A\eta_i = \lambda_1(A)\eta_i$, $i = 1, 2, \dots, r-1$. Moreover, by Lemma 2 $C = \alpha A$ for some $\alpha \neq 0$. But this contradicts (2.24). This contradiction proves that there exists a nonzero matrix in \mathcal{U} satisfying the condition $\lambda_1(A) = \dots = \lambda_r(A)$.

We now show that the bound $\kappa(r)$ is sharp. Consider the subspace \mathscr{U} of $n \times n$ symmetric matrices $A = (a_y)$ of the form

(2.26)
$$a_{ij} = 0, \quad i, j = 1, \cdots, n - r + 1,$$

(2.27)
$$\sum_{i=n-r+2}^{n} a_{ii} = 0.$$

It is clear that the dimension of this subspace is $\kappa(r) - 1$. We claim that there exists no $A \neq 0$ in \mathcal{U} which satisfies $\lambda_1(A) = \lambda_r(A)$. Suppose to the contrary that such A exists. As $\operatorname{tr}(A) = 0$ and $A \neq 0$ we must have that $\lambda_1(A) > 0$. Consider the matrix $B = \lambda_1(A)I - A$. The assumption $\lambda_1(A) = \lambda_r(A)$ implies that the rank of B does not exceed n - r. From the conditions (2.26) we deduce that the principal minor $B(\frac{1}{1,\dots,n-r+1}) = \lambda_1(A)^{n-r+1} \neq 0$. So the rank of B is at least n - r + 1. From the contradiction above we deduce the non-existence of $A \neq 0$ in \mathcal{U} satisfying $\lambda_1(A) = \lambda_r(A)$. The proof of the theorem is completed. REMARK 1. By modifying the example given in the proof of Theorem 1 we demonstrate that the bound $\kappa(r) + 1$ which was given in Lemma 1 is sharp. Consider the $\kappa(r)$ dimensional subspace \mathcal{U} given by the condition (2.26). Let $A \neq 0$ and $\lambda_1(A) = \lambda_r(A)$. The existence of such A follows from Theorem 1. Now let $B = \lambda_1(A)I - A$. Thus the rank of B does not exceed n - r. So $B(1, \dots, n-r+1) = \lambda_1(A)^{n-r+1} = 0$.

Theorem 1 shows that the situation described in Lemma 2 can only hold for r = n. Thus we have

COROLLARY 1. Let \mathcal{U} be a subspace of \mathcal{W}_n of co-dimension 1 (dim $\mathcal{U} = n(n+1)/2 - 1$). Assume that \mathcal{U} does not contain the identity matrix I. Then for any given n-1 orthonormal vectors $\eta_1, \dots, \eta_{n-1}$ there exists a unique nonzero matrix A in \mathcal{U} (up to a multiplication by positive scalar) such that

(2.28)
$$\lambda_1(A) = \cdots = \lambda_{n-1}(A) > \lambda_n(A)$$

and the corresponding eigenspace for the eigenvalue $\lambda_1(A)$ is spanned by $\eta_1, \dots, \eta_{n-1}$.

3. The equivalence of Theorems 1 and 2. We regard W_n as a real inner product space with the standard inner product (A, B) = tr(AB). Let

$$(3.1) B\xi_i = \lambda_i (B)\xi_i, (\xi_j, \xi_j) = \delta_{ij}, i, j = 1, \cdots, n.$$

Then by choosing $[\xi_1, \dots, \xi_n]$ as a basis in \mathbb{R}^n we obtain

(3.2)
$$\operatorname{tr}(AB) = \sum_{i=1}^{n} \lambda_{i}(B)(A\xi_{i}, \xi_{i}).$$

We need in the sequel the following well known lemma (cf. [3]).

LEMMA 3. Let \mathcal{U} be a subspace and \mathcal{K} be a pointed closed convex cone in \mathbb{R}^n . Let \mathcal{U}^{\perp} be the orthogonal complement of \mathcal{U} and \mathcal{K}^* the dual of \mathcal{K} in \mathbb{R}^n . Then the following are equivalent

(a)
$$\mathcal{U} \cap \mathcal{H} = \{0\}.$$

(b) $\mathcal{U}^{\perp} \cap interior \ \mathcal{K}^* \neq \emptyset$.

Now let \mathcal{K} be the cone of positive semidefinite matrices in \mathcal{W}_n . It is a well known fact that $\mathcal{K}^* = \mathcal{K}$. Finally we remark that the functions $\kappa(r)$ and f(r) defined by (1.5) and (1.8), respectively, satisfy the identity

(3.3)
$$\kappa(r) + f(n-r+1) = \dim \mathcal{W}_n, \qquad r = 1, \cdots, n.$$

(In case that \mathcal{W}_n is the space of $n \times n$ hermitian matrices we use the Definitions (1.10) and (1.11).)

Theorem 1 implies Theorem 2. Suppose that the subspace \mathcal{V} of \mathcal{W}_n satisfies the assumptions of Theorem 2. By Lemma 3 it suffices to prove that

$$(3.4) \qquad \qquad \mathcal{V}^{\perp} \cap \mathcal{K} = \{0\}.$$

Suppose this is not the case. It follows from (1.6) and (3.2) that \mathcal{V}^{\perp} contains no nonzero positive semidefinite matrix of rank *r* or less. Let d = dimension of \mathcal{V}^{\perp} . It follows from (1.7) and (3.3) that

$$(3.5) \quad d = \frac{n(n+1)}{2} - k > \frac{n(n+1)}{2} - f(r+1) + \delta_{n,r+1} = \kappa(n-r) + \delta_{n,r+1}.$$

Since $1 \le r \le n-1$ we have $1 \le n-r \le n-1$.

Suppose first that \mathcal{V}^{\perp} contains a positive definite matrix. Since the assumptions and the conclusion of Theorem 2 remain valid under a congruence transformation, we may assume that $I \in \mathcal{V}^{\perp}$. If $r \leq n-2$ then (3.5) and Theorem 1 imply that there exists a nonzero matrix in \mathcal{V}^{\perp} such that $\lambda_1(A) = \lambda_{n-r}(A) > \lambda_n(A)$. Hence there exists a nonzero positive semidefinite matrix in \mathcal{V}^{\perp} of the form $\alpha A + \beta I$ which has rank r or less, contrary to our assumption. If r = n - 1 then $d \geq 2$, by (3.5). Hence there exists A in \mathcal{V}^{\perp} which is linearly independent of I. The matrix $\lambda_1(A)I - A$ is a nonzero positive semidefinite matrix of rank n-1 or less, contrary to our assumption.

It remains to consider the case that \mathcal{V}^{\perp} contains no positive definite matrix. Let A_1 be a nonzero positive semidefinite in \mathcal{V}^{\perp} of minimal rank q. Then $q \ge r+1$. Hence we may assume that $1 \le r \le n-2$. We may also assume that

$$A_1 = \begin{bmatrix} I_q & 0 \\ 0 & 0 \end{bmatrix}.$$

Let A_1, A_2, \dots, A_d be a basis for \mathscr{V}^{\perp} . Partition these matrices in the form

$$A_i = [A_i^{(1)}, A_i^{(2)}], \qquad i = 1, 2, \cdots, d,$$

where $A_i^{(1)}$ is of size $n \times q$. We claim that the matrices $A_2^{(2)}, \dots, A_d^{(2)}$ are linearly dependent. Indeed, consider

$$\sum_{i=2}^{d} \alpha_i A_i^{(2)} = 0.$$

This leads to a linear system of $n(n+1)/2 - q(q+1)/2 = \kappa(n+1-q)$ equations in d-1 unknowns. By (3.5) $d-1 \ge \kappa(n-r)$, so we get a nontrivial solution with the only possible exception being q = r+1 and $d-1 = \kappa(n-r)$. But in the latter case, if $A_2^{(2)}, \dots, A_d^{(2)}$ are linearly independent, we may form a new basis for \mathcal{V}^{\perp} that contains among its matrices the matrix A_1 and the matrices B_1, B_2, \dots, B_{n-q} , where

$$\boldsymbol{B}_{i} = \begin{bmatrix} \boldsymbol{B}_{11}^{i} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{E}_{ii} \end{bmatrix}, \qquad i = 1, 2, \cdots, n-q.$$

Here E_{ii} is the matrix of order $n - q \times n - q$ all of whose entries are zero except the *i*, *i* entry which is 1. We can now form a positive definite matrix as a linear combination of A_1, B_1, \dots, B_{n-q} , contrary to assumption. Hence $A_2^{(2)}, \dots, A_d^{(2)}$ are linearly dependent.

Hence there exists a matrix B, $B = \sum_{i=2}^{d} \alpha_i A_i$, such that $b_{ij} = 0$ whenever i > q or j > q. Clearly, there exists a linear combination of A_1 and B which is nonzero and positive semidefinite of rank q - 1 or less. This contradicts the definition of q. Hence (3.4) is satisfied, completing the proof.

Theorem 2 implies Theorem 1. Assume that $2 \le r \le n-1$ and that \mathscr{U} satisfies the assumptions of Theorem 1. Suppose that \mathscr{U} contains no nonzero matrix A such that $\lambda_1(A) = \lambda_r(A)$. Then $I \not\in \mathscr{U}$ and let $\mathscr{U}_1 =$ linear space spanned by \mathscr{U} and I. Clearly dim $\mathscr{U}_1 \ge \kappa(r) + 1$. Let $\mathscr{V} = \mathscr{U}_1^{\perp}$, so $\mathscr{U}_1 = \mathscr{V}^{\perp}$. The subspace \mathscr{U}_1 contains no nonzero positive semidefinite matrix of rank n-r or less. Now (3.3) implies that dim $\mathscr{V} < f(n-r+1)$. Since $n-r \le n-2$ we have that $\delta_{n,n+1-r} = 0$, so the subspace \mathscr{V} satisfies the assumptions of Theorem 2. It follows that \mathscr{V} contains a positive definite matrix. However, since I is in \mathscr{U}_1 , from the fact that $\mathscr{V} = \mathscr{U}_1^{\perp}$ it follows that for any A in \mathscr{V} we must have that tr(AI) = tr(A) = 0. Thus \mathscr{V} could not contain a positive definite matrix. This contradiction implies the existence of $A \ne 0$ in \mathscr{U} such that $\lambda_1(A) = \lambda_r(A)$.

4. Extensions and remarks. We now reformulate Theorems 1 and 2 in the case where \mathcal{W}_n is the n^2 dimensional real space of $n \times n$ complex valued hermitian matrices.

THEOREM 3. Let \mathcal{U} be a k dimensional subspace in the space \mathcal{W}_n of $n \times n$ complex valued hermitian matrices. Assume that an integer r satisfies the inequalities $2 \leq r \leq n-1$. If $k \geq \kappa(r)$, where $\kappa(r) = (r-1)(2n-r+1)$, then \mathcal{U} contains a nonzero matrix such that the greatest eigenvalue of A is at least of multiplicity r. The lower bound $\kappa(r)$ is best possible for $2 \leq r \leq n-1$.

Proof. The proof of this theorem is identical with the proof of Theorem 1 except for the following detail. Let ξ_1, \dots, ξ_{r-1} be r-1 orthonormal vectors. Consider the system

(4.1)
$$A\xi_i = \lambda\xi_i, \qquad j = 1, \cdots, r-1,$$

where A belongs to \mathcal{U} . We claim that this system is equivalent to $\kappa(r)$ real valued equations. Indeed, if we complete the set ξ_1, \dots, ξ_{r-1} to a basis of orthonormal vectors $[\xi_1, \dots, \xi_n]$ then, assuming this to be the standard basis, we obtain instead of (4.1):

and

(4.3)
$$a_{\mu\nu} = 0, \quad \mu = 1, \cdots, r-1; \quad \nu = \mu + 1, \cdots, n.$$

Since $A = (a_{ij})$, is hermitian, $a_{\mu\mu}$ is real. So (4.2) is equivalent to r-1 equations. Since $a_{\mu\nu}$ for $\mu \neq \nu$ is complex valued, (4.3) is equivalent to (r-1)(2n-r) real equations. This fact explains the change of the value of $\kappa(r)$ in case that \mathcal{W}_n is the space of hermitian matrices. End of proof.

Finally, we restate Bohnenblust's theorem for the hermitian case.

THEOREM 4 (Bohnenblust). Let \mathcal{V} be a subspace of dimension k in \mathcal{W}_n and let $1 \leq r \leq n-1$. Assume that for any A in \mathcal{V} the equality (1.6) implies that $x_i = 0$ for $i = 1, \dots, r$. If the inequality (1.7) holds where $f(r) = r^2$, then \mathcal{V} contains a positive definite matrix.

ACKNOWLEDGEMENT. After the completion of this manuscript, we have received Bohnenblust's unpublished original manuscript. His proof of Theorems 2 and 4 differs from our approach.

REFERENCES

1. F. Bohnenblust, Joint positiveness of matrices, unpublished manuscript.

2. E. Calabi, Linear systems of real quadratic forms, Proc. Amer. Math. Soc., 15 (1964), 844-846.

3. A. Ben Israel, *Complex Linear Inequalities, Inequalities III*, edited by O. Shisha, Academic Press, New York and London, 1972.

4. O Taussky, Positive Definite Matrices, Inequalities I, edited by O. Shisha, Academic Press, New York, 1967.

Received September 4, 1975. The first author was supported in part by N.S.F. Grant M.P.S. 72-05055 A02.

The'Institute for Advanced Study and The University of Tennessee

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California Los Angeles, California 90024 J. Dugundл

Department of Mathematics University of Southern California Los Angeles, California 90007

D. GILBARG AND J. MILGRAM Stanford University Stanford, California 94305

University of Washington Seattle, Washington 98105

ASSOCIATE EDITORS

E. F. BECKENBACH

R. A. BEAUMONT

B. H. NEUMANN

F. Wolf

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF HAWAII UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

> Copyright © 1976 Pacific Journal of Mathematics All Rights Reserved

Pacific Journal of Mathematics Vol. 62, No. 2 February, 1976

Christopher Allday, <i>The stratification of compact connected Lie group</i> <i>actions by subtori</i>
actions by subtori
-
Martin Bartelt, <i>Commutants of multipliers and translation operators</i> 329
Herbert Stanley Bear, Jr., Ordered Gleason parts
James Robert Boone, On irreducible spaces. II
James Robert Boone, On the cardinality relationships between discrete
collections and open covers
L. S. Dube, On finite Hankel transformation of generalized functions 365
Michael Freedman, Uniqueness theorems for taut submanifolds 379
Shmuel Friedland and Raphael Loewy, Subspaces of symmetric matrices
containing matrices with a multiple first eigenvalue
Theodore William Gamelin, Uniform algebras spanned by Hartogs
series
James Guyker, On partial isometries with no isometric part
Shigeru Hasegawa and Ryōtarō Satō, A general ratio ergodic theorem for
<i>semigroups</i>
Nigel Kalton and G. V. Wood, <i>Homomorphisms of group algebras with norm</i> less than $\sqrt{2}$
Thomas Laffey, On the structure of algebraic algebras 461
Will Y. K. Lee, On a correctness class of the Bessel type differential operator S. 473
Robert D Little Complex vector fields and divisible Chern classes 483
Kenneth Louden Maximal quotient rings of ring extensions 480
Dieter Lutz Scalar spectral operators ordered l^{ρ} direct sums and the
counterexample of Kakutani-McCarthy 407
Ralph Tyrrell Rockafellar and Roger Jean-Baptiste Robert Wets Stochastic
convex programming: singular multipliers and extended duality
Edward Barry Saff and Richard Steven Varga, Geometric overconvergence of
rational functions in unbounded domains
Ioel Linn Schiff Isomorphisms between harmonic and P-harmonic Hardy
spaces on Riemann surfaces 551
Virinda Mohan Sehgal and S. P. Singh <i>On a fixed point theorem of</i>
Krasnoselskii for locally convex spaces
Lewis Shilane. Filtered spaces admitting spectral sequence operations 569
Michel Smith. Generating large indecomposable continua
John Yuan, On the convolution algebras of H-invariant measures