

Pacific Journal of Mathematics

WEAK TYPE MULTIPLIERS FOR HANKEL TRANSFORMS

WILLIAM C. CONNETT, V AND ALAN SCHWARTZ

WEAK TYPE MULTIPLIERS FOR HANKEL TRANSFORMS

WILLIAM C. CONNETT AND ALAN L. SCHWARTZ

The main result of this paper is that weak type multiplier theorems for Jacobi expansions yield weak type multiplier theorems for Hankel transforms.

In recent papers the authors studied multiplier theorems for ultraspherical and Jacobi expansions. An interesting paper of Igari suggested a new approach to multiplier theorems that used asymptotics instead of the elaborate machinery used earlier.

This paper extends the method of Igari to give the first weak type multiplier theorem for Hankel transforms. This extension is important in itself and because this method together with the authors' results for Jacobi multipliers will generalize to the "radial" functions associated with the other compact symmetric spaces.

Let $\{P_n^{(\alpha, \beta)}(x)\}$ be the Jacobi polynomials with indices (α, β) . The functions $\{P_n^{(\alpha, \beta)}(\cos \theta)\}$ are orthogonal with respect to the measure $d\mu(\theta) = (\sin \theta/2)^{2\alpha+1} (\cos \theta/2)^{2\beta+1} d\theta$. For measurable f on $[0, \pi]$ define

$$\|f\|_p = \left\{ \int_0^\pi |f(\theta)|^p d\mu(\theta) \right\}^{1/p}$$

and

$$\hat{f}(n) = \int_0^\pi f(\theta) P_n^{(\alpha, \beta)}(\cos \theta) d\mu(\theta)$$

so that if

$$h_n^{-1} = \int_0^\pi [P_n^{(\alpha, \beta)}(\cos \theta)]^2 d\mu(\theta),$$

then

$$f(\theta) \sim \sum_{n=0}^{\infty} \hat{f}(n) h_n P_n^{(\alpha, \beta)}(\cos \theta)$$

where equality holds at least for finite series.

The multiplier transformation defined by the function $\phi(x)$ is denoted by T_ϕ , where at least formally,

$$T_\phi f(\theta) \sim \sum_{n=0}^{\infty} \phi(n) \hat{f}(n) h_n P_n^{(\alpha, \beta)}(\cos \theta).$$

The operator T_ϕ is said to be of strong type p , $1 < p < \infty$ if

$$\|T_\phi\|_p = \|\phi(n)\|_p = \sup \|T_\phi f\|_p \quad (\|f\|_p = 1).$$

The operator is said to be of weak type p if

$$\langle\langle T_\phi \rangle\rangle_p = \langle\langle \phi(n) \rangle\rangle_p = \sup \lambda [\mu\{x: |T_\phi f(x)| > \lambda\}]^{1/p} \quad (\lambda > 0, \|f\|_p = 1).$$

The same objects of harmonic analysis are also needed for the Hankel transforms. Let $d\eta(x) = x^{2\alpha+1}dx$ $\mathcal{J}_\alpha(x) = x^{-\alpha}J_\alpha(x)$ and briefly define:

$$\|g\|_p = \left(\int_0^\infty |g(x)|^p d\eta(x) \right)^{1/p},$$

$$\hat{g}(x) = \int_0^\infty g(y) \mathcal{J}_\alpha(xy) d\eta(y)$$

(at least for g continuous with compact support),

$$g(x) \sim \int_0^\infty \hat{g}(y) \mathcal{J}_\alpha(xy) d\eta(y),$$

$$U_\phi g \sim \int_0^\infty \phi(y) \hat{g}(y) \mathcal{J}_\alpha(xy) d\eta(y),$$

$\|U_\phi\|_p$ for the strong type p norm of the operator,

$\langle U_\phi \rangle_p$ for the weak type p norm.

The idea of Igari [4] is to transform strong type multiplier theorems for Jacobi expansions into strong type multiplier theorems for Hankel transforms. His result is

THEOREM (Igari). *Let $\alpha, \beta > -1$ and assume ϕ is continuous on $(0, \infty)$ except on a null set. If $1 \leq p < \infty$, then*

$$\|\phi(x)\|_p = \liminf_{\varepsilon \rightarrow 0} \|\phi(\varepsilon n)\|_p.$$

Igari's techniques are adapted here to prove a similar theorem for weak type multipliers.

THEOREM 1. *Under the same hypotheses as above*

$$\langle \phi(x) \rangle_p = \liminf_{\varepsilon \rightarrow 0} \langle\langle \phi(\varepsilon n) \rangle\rangle_p.$$

The theorem will be proved following a

LEMMA. *Let $E \subset R$ satisfy $\eta(E) < \infty$ and let $f_n(x) \rightarrow f(x)$ a.e. in E as $n \rightarrow \infty$. Then if $s > 0$*

$$\eta\{x \in E: |f(x)| > s\} \leq \liminf_{n \rightarrow \infty} \eta\{x \in E: |f_n(x)| > s\}.$$

The lemma can be proved by applying Fatou's lemma to the characteristic function of $\{x \in E: |f_n(x)| > s\}$.

Proof of Theorem 1. It will be sufficient to prove the theorem for compactly supported infinitely differentiable functions, so let g be one such with support in $(0, M)$.

Let $g_\lambda(\theta) = g(\lambda\theta)$ for λ so large that $\lambda\pi > M$, and let

$$G(\tau, \lambda) = \sum_{n=0}^{\infty} \phi\left(\frac{n}{\lambda}\right) g_\lambda(n) P_n^{(\alpha, \beta)}\left(\cos \frac{\tau}{\lambda}\right),$$

$$G^N(\tau, \lambda) = \sum_{n=0}^{N[\lambda]} \phi\left(\frac{n}{\lambda}\right) g_\lambda(n) P_n^{(\alpha, \beta)}\left(\cos \frac{\tau}{\lambda}\right),$$

$$H^N(\tau, \lambda) = G(\tau, \lambda) - G^N(\tau, \lambda),$$

$$G(\tau) = \int_0^\infty \phi(y) \hat{g}(y) \mathcal{F}_\alpha(\tau y) d\eta(y) = U_\phi g(\tau).$$

A careful reading of Igari's argument shows that there is a constant B independent of λ, N and K such that once λ is sufficiently large

$$(1) \quad \int_0^K |H^N(\tau, \lambda)|^2 d\eta(\tau) \leq B/N^2.$$

He shows that $G^N(\tau, \lambda)$ converges everywhere to a function $G^N(\tau)$ and that a subsequence of $G^N(\tau)$ converges a.e. in $(0, \infty)$ to $G(\tau)$.

Let $\varepsilon > 0, \delta > 0, K > 0$, and $s > 0$ and define

$$M_\lambda = \left\langle \left\langle \phi\left(\frac{n}{\lambda}\right) \right\rangle \right\rangle_p^p \|g_\lambda\|_p^2 s^{-p}.$$

If $\phi(n/\lambda)$ defines a weak type p operator on Jacobi series then

$$\mu\{\phi: |G(\lambda\phi, \lambda)| > s\} \leq M_\lambda.$$

Then if λ is sufficiently large

$$\eta\{\tau \leq K: |G(\tau, \lambda)| > s\} \leq 2^{2\alpha+1} \lambda^{2\alpha+2} M_\lambda (1 + \varepsilon).$$

Let $N^2 > B/\varepsilon\delta^2$ so that relation (1) implies

$$\eta\{\tau \leq K: |H^N(\tau, \lambda)| > \delta\} \leq \varepsilon,$$

and $G^N(\tau, \lambda) = G(\tau, \lambda) - H^N(\tau, \lambda)$ so that

$$\{\tau \leq K: |G^N(\tau, \lambda)| > s + \delta\} \subset \{\tau \leq K: |G(\tau, \lambda)| > s\} \cup \{\tau \leq K: |G(\tau, \lambda)| \leq s \text{ and } |H^N(\tau, \lambda)| > \delta\},$$

thus

$$(2) \quad \eta\{\tau \leq K: |G^N(\tau, \lambda)| > s + \delta\} \leq 2^{2\alpha+1} \lambda^{2\alpha+2} M_\lambda (1 + \varepsilon) + \varepsilon.$$

Now choose $\lambda_j \rightarrow \infty$ such that

$$\lim_{j \rightarrow \infty} \left\langle \left\langle \phi \left(\frac{n}{\lambda_j} \right) \right\rangle \right\rangle_p^p = \liminf_{\lambda \rightarrow \infty} \left\langle \left\langle \phi \left(\frac{n}{\lambda} \right) \right\rangle \right\rangle_p^p$$

and observe that $\lim_{\lambda \rightarrow \infty} 2^{\alpha+1} \lambda^{2\alpha+2} \|g_\lambda\|_p^p = \|g\|_p^p$ so

$$\lim 2^{2\alpha+1} \lambda_j^{2\alpha+2} M_{\lambda_j} = M = \liminf_{\lambda \rightarrow \infty} \left\langle \left\langle \phi \left(\frac{n}{\lambda} \right) \right\rangle \right\rangle_p^p \|g\|_p^p s^{-p},$$

thus if λ is replaced by λ_j in (2) and then j increased, the lemma implies

$$(3) \quad \eta\{\tau \leq K: |G^N(\tau)| > s + \delta\} \leq M(1 + \varepsilon) + \varepsilon.$$

If N_j is now chosen so that $G^{N_j}(\tau) \rightarrow G(\tau)$ a.e. in $(0, \infty)$, then relation (3) and the lemma yield

$$\eta\{\tau \leq K: |G(\tau)| > s + \delta\} \leq M(1 + \varepsilon) + \varepsilon.$$

Now, first the ε , then the δ , and finally the K can be removed by standard arguments to yield

$$\eta\{\tau: |U_\phi g(\tau)| > s\} \leq \liminf_{\lambda \rightarrow \infty} \left\langle \left\langle \phi \left(\frac{n}{\lambda} \right) \right\rangle \right\rangle_p^p \|g\|_p^p s^{-p}$$

and the Theorem is proved.

To transfer strong type multiplier theorems from Jacobi series to Hankel transforms Igari's theorem should be used, for weak type results Theorem 1 is the tool; for instance ($[\alpha]$ is the greatest integer not exceeding α).

THEOREM 2. *Let $\alpha \geq \beta > -1/2$ and $m = [\alpha] + 2$; assume that ϕ is m times continuously differentiable in $(0, \infty)$ and let*

$$[A_m(\phi)]^2 = \sup_x |\phi(x)|^2 + \sup_M M^{-1} \int_M^{2M} |x^m \phi^{(m)}(x)|^2 x^{-1} dx,$$

then U_ϕ is of weak type 1 and

$$\langle U_\phi \rangle \leq CA_m(\phi)$$

where C does not depend on ϕ .

Proof. Define $\Delta^0 \phi(n) = \phi(n)$ and $\Delta^k \phi(n) = \Delta^{k-1} \phi(n+1) - \Delta^k \phi(n)$ for $k = 1, 2, \dots$ and let $\phi_\varepsilon(x) = \phi(\varepsilon x)$.

It can be shown (see Bonami and Clerc [1], Theorem 4.12) that

$$|\Delta^m \phi_\varepsilon(n)| \leq \varepsilon^{m-1} \int_{\varepsilon n}^{\varepsilon(n+m)} |\phi^{(m)}(x)| dx.$$

Consequently by the Schwarz inequality and the Mean Value Theorem

for integrals there is constant C independent of n and ϕ such that

$$\frac{1}{n} |n^m \Delta^m \phi_\varepsilon(n)|^2 \leq m \int_{\varepsilon n}^{\varepsilon(n+m)} |x^m \phi^{(m)}(x)|^2 x^{-1} dx.$$

Thus if

$$[B_m(\phi)]^2 = \sup_n |\phi(n)|^2 + \sup_M M^{-1} \sum_{n=M}^{2M} |n^m \Delta^m \phi(n)|^2,$$

then there is a constant C_1 independent of ϕ such that

$$(4) \quad B_m(\phi_\varepsilon) \leq C_1 A_m(\phi).$$

The main result of [3] actually shows that if $\alpha \geq \beta > -1/2$ and $m = [\alpha] + 2$ there is a constant C_2 such that

$$\langle\langle T_\phi \rangle\rangle_1 \leq C_2 B_m(\phi).$$

Thus the result follows from Theorem 1 and (4).

REFERENCES

1. A. Bonami and J.-L. Clerc, *Sommes de Cesàro et multiplicateurs des développements en harmoniques sphériques*, Trans. Amer. Math. Soc., **183** (1973), 223-263.
2. W. C. Connett and A. L. Schwartz, *A multiplier theorem for ultraspherical series*, Studia Mathematica, **L** (1973), 51-70.
3. ———, *A multiplier theorem for Jacobi series*, Studia Mathematica, **LII** (1975), 243-261.
4. S. Igari, *On the multipliers of Hankel transform*, Tôhoku, Math. J., Second Series, **24** (1972), 201-206.

Received March 15, 1974 and in revised form December 4, 1975. Supported in part by AFOSR Grant 71-2047.

UNIVERSITY OF MISSOURI—ST. LOUIS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 63, No. 1

March, 1976

Ralph Artino, <i>Gevrey classes and hypoelliptic boundary value problems</i>	1
B. Aupetit, <i>Caractérisation spectrale des algèbres de Banach commutatives</i>	23
Leon Bernstein, <i>Fundamental units and cycles in the period of real quadratic number fields. I</i>	37
Leon Bernstein, <i>Fundamental units and cycles in the period of real quadratic number fields. II</i>	63
Robert F. Brown, <i>Fixed points of automorphisms of compact Lie groups</i>	79
Thomas Ashland Chapman, <i>Concordances of noncompact Hilbert cube manifolds</i>	89
William C. Connett, V and Alan Schwartz, <i>Weak type multipliers for Hankel transforms</i>	125
John Wayne Davenport, <i>Multipliers on a Banach algebra with a bounded approximate identity</i>	131
Gustave Adam Efroymsen, <i>Substitution in Nash functions</i>	137
John Sollion Hsia, <i>Representations by spinor genera</i>	147
William George Kitto and Daniel Eliot Wulbert, <i>Korovkin approximations in L_p-spaces</i>	153
Eric P. Kronstadt, <i>Interpolating sequences for functions satisfying a Lipschitz condition</i>	169
Gary Douglas Jones and Samuel Murray Rankin, III, <i>Oscillation properties of certain self-adjoint differential equations of the fourth order</i>	179
Takaši Kusano and Hiroshi Onose, <i>Nonoscillation theorems for differential equations with deviating argument</i>	185
David C. Lantz, <i>Preservation of local properties and chain conditions in commutative group rings</i>	193
Charles W. Neville, <i>Banach spaces with a restricted Hahn-Banach extension property</i>	201
Norman Oler, <i>Spaces of discrete subsets of a locally compact group</i>	213
Robert Olin, <i>Functional relationships between a subnormal operator and its minimal normal extension</i>	221
Thomas Thornton Read, <i>Bounds and quantitative comparison theorems for nonoscillatory second order differential equations</i>	231
Robert Horace Redfield, <i>Archimedean and basic elements in completely distributive lattice-ordered groups</i>	247
Jeffery William Sanders, <i>Weighted Sidon sets</i>	255
Aaron R. Todd, <i>Continuous linear images of pseudo-complete linear topological spaces</i>	281
J. Jerry Uhl, Jr., <i>Norm attaining operators on $L^1[0, 1]$ and the Radon-Nikodým property</i>	293
William Jennings Wickless, <i>Abelian groups in which every endomorphism is a left multiplication</i>	301