

Pacific Journal of Mathematics

**ABELIAN GROUPS IN WHICH EVERY ENDOMORPHISM IS A
LEFT MULTIPLICATION**

WILLIAM JENNINGS WICKLESS

ABELIAN GROUPS IN WHICH EVERY ENDOMORPHISM IS A LEFT MULTIPLICATION

W. J. WICKLESS

Let $\langle G+ \rangle$ be an abelian group. With each multiplication on G (binary operation $*$ such that $\langle G+ * \rangle$ is a ring) and each $g \in G$ is associated the endomorphism g_l^* of left multiplication by g . Let $L(G) = \{g_l^* \mid g \in G, * \in \text{Mult } G\}$. Abelian groups G such that $L(G) = E(G)$ are studied. Such groups G are characterized if G is torsion, reduced algebraically compact, completely decomposable, or almost completely decomposable of rank two. A partial results is obtained for mixed groups.

Let $\langle G+ \rangle$ be an abelian group. With each multiplication on G (binary operation $*$ such that $\langle G+ * \rangle$ is a ring) and each $g \in G$ is associated the endomorphism g_l^* of left multiplication by g given by $g_l^*(x) = g * x, x \in G$. Let $L(G)$ be the set of all such endomorphisms, i.e., $L(G) = \{g_l^* \mid g \in G, * \in \text{Mult}(G)\}$. In general all one can say is that $L(G)$ is a subset of the endomorphism ring $E(G)$. In this paper we consider abelian groups G such that every endomorphism is a left multiplication.

DEFINITION 1. An abelian group G is multiplicatively faithful iff $L(G) = E(G)$.

We mostly follow the notations in [2]. Specifically: all groups are abelian, rings are not necessarily associative, \otimes denotes the tensor product over Z and $g \otimes_*$ the natural map $x \rightarrow g \otimes x$ from G into $G \otimes G$, $o(x)$ is the order of an element x , $Z(d)$ is the cyclic group of order d and $Z(d)^*$ is the multiplicative group of units in $Z(d)$. For a prime p , we write Z_p for the localization of Z at p and \hat{Z}_p for the ring (or group) of p -adic integers. We use $t(A)[t(x)]$ for the type of a rank one torsion free group A [element x] and $h(x)$ for the height sequence. Finally, $\langle S \rangle[\langle S \rangle_*]$ [is the subgroup [pure subgroup] generated by S .

We begin by listing some simple results.

A. Let $\theta_g: \text{Hom}(G \otimes G, G) \rightarrow E(G)$ be given by $\theta_g(\Delta) = \Delta \circ (g \otimes_*)$, $\Delta \in \text{Hom}(G \otimes G, G)$, $g \in G$. Then G is multiplicatively faithful iff $\bigcup_{g \in G} \text{Image } \theta_g = E(G)$.

Proof. Mult G , the group of all multiplications on G , is isomorphic

to $\text{Hom}(G \otimes G, G)$. Under this identification $\Delta \circ (g \otimes _) = g_1$.

B. G is multiplicatively faithful iff for each $\theta \in E(G)$, there exists $u \in G, \sigma \in \text{Mult } G$ such that the following diagram commutes:

$$\begin{array}{ccc} G & \xrightarrow{u \otimes _} & G \otimes G \\ \searrow \theta & & \swarrow \sigma \\ & & G \end{array}$$

Proof. Obvious.

C. A divisible group is multiplicatively faithful iff it is torsion free. More generally, if $G = D \oplus R, D$ the maximal divisible subgroup of G with D torsion free, then $L(G) = E(G)$ iff $L(R) = E(R)$.

Proof. This follows directly from (B) and elementary properties of the tensor product.

D. If Z is a direct summand of G , then $L(G) = E(G)$. More generally, if A is a ring, $1 \in A$, and H is a unital A module, then $A \oplus H$ is multiplicatively faithful.

Proof. Let $\theta \in E(A \oplus H)$. Set $u = 1 \in A$, and define $\sigma \in \text{Mult } G$ by $\sigma(\sum a_i \otimes x_i \oplus y) = \sum a_i \theta(x_i); a_i \in A, x_i \in A \oplus H, y \in H \otimes (A \oplus H)$. Then

$$\begin{array}{ccc} A \oplus H & \xrightarrow{1 \otimes _} & (A \oplus H) \otimes (A \oplus H) \\ \searrow \theta & & \swarrow \sigma \\ & & A \oplus H \end{array}$$

commutes.

E. Let $R(G)$ be the set of all right multiplications by elements of G for all rings on G . Then $L(G) = E(G)$ iff $R(G) = E(G)$.

Proof. This follows from considering opposite rings.

Multiplicatively faithful torsion groups are easily characterized.

THEOREM 1. *Let G be a torsion group. Then G is multiplicatively faithful iff G is bounded.*

Proof. If $L(G) = E(G)$, then there exists $u \in G, \sigma \in \text{Mult } G$ such that $\sigma \circ (u \otimes _) = 1_G$, where 1_G is the identity endomorphism. It follows

that $nG = (0)$, where $n = o(u)$. If $nG = (0)$, $n \in Z^+$, we can write $G = Z(n) \oplus H$. (D) applies to give $L(G) = E(G)$.

We next consider mixed groups, and characterize the multiplicatively faithful ones in one special case.

THEOREM 2. *Let G be mixed with maximal torsion subgroup $T = \bigoplus_p T_p$. Suppose that $T_p \neq (0)$ for only a finite number of primes p , and also that G/T is homogeneous completely decomposable. Then $L(G) = E(G)$ iff (1) $G = T \oplus F$, (2) each rank 1 summand of G/T has idempotent type, (3) $p(G/T) = G/T$ implies T_p is bounded.*

Proof. Suppose (1), (2) and (3) hold for G as above. Let $T = T_1 \oplus T_2$, where T_1 is the sum of the bounded and T_2 the sum of the unbounded p components of T . Since T_1 is bounded, write $T_1 = Z(n) \oplus X$ with X a unital $Z(n)$ module. $F \cong G/T$ is homogeneous, completely decomposable and nonzero. Say $F = A \oplus B$ where A is torsion free of rank one and $B = \bigoplus_{\alpha \in I} (A)_\alpha$. ($I = \emptyset$ is allowed.) Since $t(A)$ is idempotent, A is (may be regarded as) a subring with identity of Q ([2], Th. 121.1). Moreover, since $pA = A$ only when $(T_2)_p = (0)$, $B \oplus T_2$ may be made into a unital A module in the natural way. Thus, $X \oplus B \oplus T_2$ is a unital $Z(n) \oplus A$ module and (D) applies to show G is multiplicatively faithful.

Conversely, let $L(G) = E(G)$ for G satisfying the conditions of our theorem. Let $u \in G$ be such that $u_i^* = 1_G$, $*$ some multiplication on G . If $u \in pG$, clearly $T_p = (0)$.

Now consider a prime p such that $u + T \in p(G/T)$. Since $(u + T)_i$ induces the identity endomorphism on G/T , it follows immediately that $u + T \in p^n(G/T)$ for all $n \in Z^+$. Write $u = pg + t = pg + t_1 + t_2$, where $o(t_1) = p^k$, $(o(t_2), p) = 1$. If $t_1 = 0$, then $u \in pG$ and $T_p = (0)$. If $t_1 \neq 0$, then, for all $x \in T_p$,

$$x = u * x = (pg + t_1 + t_2) * x = p(g * x) + t_1 * x.$$

(Since $(o(t_2), p) = 1$ and $x \in T_p$, $t_2 * x = 0$.) But $o[p(g * x)] < o(x)$, $o(t_1 * x) \leq o(x)$, so $o(x) = o(t_1 * x) \leq o(t_1)$. Thus T_p is bounded.

Thus, for each p such that $u + T \in p(G/T)$, we have $u + T \in p^n(G/T)$ for all $n \in Z^+$, and T_p is bounded. Since $t(u + T)$ is the type of each rank 1 summand of G/T —(recall G/T is homogeneous)—(2) and (3) hold. Let T_1, T_2 be as before. Since T_1 is bounded, $G = T_1 \oplus H$ with $T_2 \subseteq H$.

To establish (1), we must show that T_2 is a direct summand of H . Write H/T_2 as a direct sum of isomorphic rank one groups, $H/T_2 = \bigoplus A_i$, and let $A_i = \langle a_i + T_2 \rangle_*$ where $h(a_i + T_2) = (m_{ij})$, $m_{ij} = 0$ or ∞ for all i, j . Since $p(H/T_2) = H/T_2 \rightarrow (T_2)_p = (0)$, the following

implication holds: $\alpha_i + T_2 \in p(H/T_2) \rightarrow \alpha_i \in pH$. From this one easily obtains $H = T_2 \oplus F$, where $F = \langle \{\alpha_i\} \rangle_*$.

REMARK. The condition $T_p \neq (0)$ for only finitely many p is necessary for the theorem. Let $G = \prod_p Z(p)$. Then $T(G) = \bigoplus_p Z(p)$ is not a direct summand of G . However, $G/T(G)$ is homogeneous completely decomposable (torsion free divisible) and—as we shall see in Theorem 3 — $L(G) = E(G)$.

We next characterize reduced algebraically compact multiplicatively faithful groups. If G is reduced algebraically compact, then $G = \prod_p G_p$, where each G_p is a complete module over \hat{Z}_p . Since each G_p is fully invariant in G ($qG_p = G_p$ for all $q \neq p$) and since $\text{Hom}(G_p \otimes G_q, G_r) = (0)$ unless $p = q = r$, it follows that $L(G) = E(G)$ iff $L(G_p) = E(G_p)$ for all p . Each G_p may be written as a completion: $G_p = (B_p^0 \oplus B_p)^\wedge$, where $B_p^0 = \bigoplus_{\alpha \in I} (\hat{Z}_p)_\alpha$, $B_p = \bigoplus_{\beta \in J} Z(p^{k_\beta})$, $0 < k_\beta < \infty$. (See [2], § 40 for details.)

THEOREM 3. *Let G be reduced algebraically compact. Then G is multiplicatively faithful iff, for each p , either $B_p^0 \neq (0)$ or G_p is bounded.*

Proof. If G_p is bounded, then $L(G_p) = E(G_p)$ by Theorem 1. If $B_p^0 \neq (0)$, write $B_p^0 = \hat{Z}_p \oplus B'$. Then $G_p = (\hat{Z}_p \oplus B' \oplus B_p)^\wedge$. Since \hat{Z}_p is algebraically compact and pure in G_p ([2], Th. 41.7, 41.9), we have $G_p = \hat{Z}_p \oplus G'$. Since G_p is a unital \hat{Z}_p module, (D) gives $L(G_p) = E(G_p)$.

Conversely, suppose G is reduced, algebraically compact and multiplicatively faithful. Then $L(G_p) = E(G_p)$ for all p . If for some p $B_p^0 = (0)$, then $B_p = \bigoplus_{\beta \in J} Z(p^{k_\beta}) \subseteq T \subseteq G_p \subseteq \prod_{\beta \in J} Z(p^{k_\beta})$, where T is the torsion subgroup of the direct product. ($T \subseteq \hat{B}_p = G_p$.) Now, G_p/T is torsion free divisible, thus homogeneous completely decomposable. Moreover, T is a p -group, and $L(G_p) = E(G_p)$. Theorem 2 applies to give a splitting $G_p = T \oplus F$. Since $G_p = \hat{T}$, $F = (0)$. Thus, G_p is a reduced algebraically compact torsion group, and is, therefore, bounded ([2], Cor. 40.3).

For the rest of the paper, we consider torsion free groups. First, we do the completely decomposable case.

THEOREM 4. *Let $G = \bigoplus_{\lambda \in \Lambda} A_\lambda$, where each A_λ is torsion free rank one. Then $L(G) = E(G)$ iff there exist subsets A, \dots, A_n of the index set Λ and rank one groups $A_{\lambda_1}, \dots, A_{\lambda_n}$, $\lambda_i \in A_i$, with (1) $A = \bigcup_{i=1}^n A_i$ and (2) $t(A_{\lambda_i}) + t(A_{\lambda'}) \leq t(A_{\lambda'})$ for all $\lambda' \in A_i$, $i = 1, \dots, n$.*

Proof. Suppose $A_1, \dots, A_n; A_{\lambda_1}, \dots, A_{\lambda_n}$ exist satisfying the above

conditions. Without loss of generality, assume A_1, \dots, A_n are disjoint. Put $\lambda' = \lambda_i$ in (2) to see that each $t(A_{\lambda_i})$ is idempotent. Thus, each A_{λ_i} can be made into a rank one ring with identity. Let $G_i = \bigoplus_{\lambda \in A_i} G_\lambda$. Due to (2), each G_i can be regarded (in the natural way) as a unital A_{λ_i} module. So we have $G = \bigoplus_{i=1}^n G_i$ is a unital A module with $A = \bigoplus_{i=1}^n A_{\lambda_i}$ (ring direct sum). Since A is a (group) direct summand of G , (D) applies.

Now suppose $G = \bigoplus_{\lambda \in A} A_\lambda$ with $L(G) = E(G)$. Choose $u \in G$, $\sigma \in \text{Mult } G$ such that $\sigma \circ (u \otimes -) = 1_G$. Write $u = \sum_{i=1}^n a_{\lambda_i}$, $a_{\lambda_i} \in A_{\lambda_i}$. Then, for all $\lambda \in A$, $\pi \sigma (\bigoplus_{i=1}^n A_{\lambda_i} \otimes A_\lambda) = A_\lambda$ when π is the projection from G onto A_λ . Thus, for each λ , there exists at least one i , $1 \leq i \leq n$, with $t(A_{\lambda_i} \otimes A_\lambda) = t(A_{\lambda_i}) + t(A_\lambda) \leq t(A_\lambda)$. The desired partition of A now easily can be constructed.

Let G be an almost completely decomposable rank two torsion free group, i.e., $G \cong A \oplus B \cong dG$ for some $d \in \mathbb{Z}^+$ and rank one subgroups A, B of G . We will obtain a numerical condition to show when such a G is multiplicatively faithful. We may assume $t(A)$ and $t(B)$ are incomparable. (If $t(A)$ and $t(B)$ are comparable, then $G \cong A \oplus B$ by Theorem 9.6 of [1]. If $G \cong A \oplus B$, Theorem 4 gives a complete description of when G is multiplicatively faithful.)

Let $A = \langle a \rangle_*$, $B = \langle b \rangle_*$ and let d be the minimal positive integer with $dG \subseteq A \oplus B$. It is easy to show that $G = \langle A \oplus B, a + nb/d \rangle \subseteq Q \oplus Q$ where n is an integer with $(n, d) = 1$. ($G/A \oplus B \cong \mathbb{Z}(d)$.)

Let $h_p(x)$ be the p -component of the height sequence of x and let $\Pi_A = \{p \mid h_p(a) = \infty\}$, $\Pi_B = \{p \mid h_p(b) = \infty\}$. It is also easy to show that $p \in \Pi_A \cup \Pi_B \rightarrow (p, d) = 1$. Let S be the multiplicative subgroup of $\mathbb{Z}(d)^*$ generated by $\Pi_A \cup \Pi_B$.

THEOREM 5. *Let $G = \langle A \oplus B, a + nb/d \rangle$ be as above. Then $L(G) = E(G)$ iff $t(A)$ and $t(B)$ are idempotent and $n \in S$.*

Proof. Suppose $L(G) = E(G)$. If either A or B — A say—had nil type, then $AG = GA = (0)$ for any multiplication on G . (Recall that $t(A), t(B)$ are incomparable.) Thus, 1_G could not be represented as a left multiplication for any ring on G . Since $L(G) = E(G)$ we must have $t(A), t(B)$ idempotent.

Since $t(A), t(B)$ are idempotent we can assume, without loss of generality, that $h_p(a) = 0$, $p \notin \Pi_A$, $h_p(b) = 0$, $p \notin \Pi_B$. Choose $\sigma \in \text{Mult}(G)$, $x = \alpha a + \beta b \in G$, $\alpha, \beta \in Q$, such that the following is a commutative diagram:

$$\begin{array}{ccc}
 G & \xrightarrow{x \otimes -} & G \otimes G \\
 & \searrow & \swarrow \sigma \\
 & & G
 \end{array}$$

Let $\overline{\Pi}_A = \{m \in Z \mid m = p_1^{e_1} \cdots p_k^{e_k}, p_i \in \Pi_A\}$ and define $\overline{\Pi}_B$ similarly. Since $t(A), t(B)$ are incomparable, we have $\sigma(a \otimes b) = \sigma(b \otimes a) = 0$; $\sigma(a \otimes a) = (c/h)a, h \in \overline{\Pi}_A$; $\sigma(b \otimes b) = (e/k)b, k \in \overline{\Pi}_B$. Let $y = a + nb/d$. Then

$$\sigma(y \otimes y) = \frac{1}{d^2} \left[\frac{c}{h} a + \frac{n^2 e}{k} b \right] \in G.$$

Since d is relatively prime both to n^2 and to anything in $\overline{\Pi}_A \cup \overline{\Pi}_B$, we must have $c = c'd, e = e'd$

$$\frac{1}{d} \left[\frac{c'}{h} a + \frac{n^2 e'}{k} b \right] \in G.$$

But $1/d[a + nb] \in G$. A short computation yields: $n^2 e'/k - nc'/h \equiv 0 \pmod{d}$. Since $(n, d) = 1$, we have $ne'h - c'k \equiv 0 \pmod{d}$.

Now $\sigma[x \otimes a] = \sigma[(\alpha a + \beta b) \otimes a] = \alpha \sigma(a \otimes a) = \alpha(c'd/h)a = 1_a(a) = a$, so $\alpha = h/c'd$. Similarly, $\beta = k/e'd$. Since $\alpha a + \beta b \in G$, we must have $c' \in \overline{\Pi}_A, e' \in \overline{\Pi}_B$. But then $n \equiv c'k/e'h \pmod{d}$, so $n \in S$. This shows the two conditions of our theorem are necessary for $L(G) = E(G)$.

Conversely, suppose $t(A), t(B)$ are idempotent and $n \in S$. Let a, b be as before. Let $\lambda \in E(G)$. Since $t(A), t(B)$ are incomparable, $\lambda(a) = (m/h)a, \lambda(b) = (t/k)b; h \in \overline{\Pi}_A, k \in \overline{\Pi}_B$. Now $\lambda(y) = 1/d[(m/h)a + (nt/k)b] \in G$, so we must have $mk - th \equiv 0 \pmod{d}$.

Since $n \in S$, it is easy to choose $c, c_1 \in \overline{\Pi}_A, e, e_1 \in \overline{\Pi}_B$ such that $nec_1 \equiv ce_1 \pmod{d}$.

Let σ be defined by $\sigma(a \otimes a) = (dc/c_1)a, \sigma(b \otimes b) = (de/e_1)b, \sigma(a \otimes b) = \sigma(b \otimes a) = 0$. To show $\sigma[G \otimes G] \subseteq G$, it is enough to check that $\sigma(y \otimes a), \sigma(a \otimes y), \sigma(y \otimes b), \sigma(b \otimes y)$ and $\sigma(y \otimes y)$ are all in G . All of these elements are obviously in G except the last one, and

$$\sigma(y \otimes y) = \frac{1}{d^2} \left[\frac{dc}{c_1} a + \frac{n^2 de}{e_1} b \right] = \frac{1}{d} \left[\frac{c}{c_1} a + \frac{n^2 e}{e_1} b \right].$$

This is in G iff $n(c/c_1) \equiv n^2(e/e_1) \pmod{d}$, which is true by choice c, c_1, e, e_1 . Thus, $\sigma \in \text{Mult } G$.

Now let

$$g = \frac{1}{d} \left[\frac{c_1 m}{he} a + \frac{e_1 t}{ke} b \right].$$

It follows directly that $\sigma \circ (g \otimes -) = \lambda$. (One need only check this identity on the independent set $\{a, b\}$.) It remains to show that $g \in G$. Now $g \in G$ iff $n[c_1 m/hc] \equiv e_1 t/ke \pmod{d}$. This congruence is easy to derive from $nec_1 \equiv ce_1 \pmod{d}$ and $mk \equiv th \pmod{d}$, both of which are given. Thus, $g \in G, g_i^2 = \lambda$, and G is multiplicatively faithful.

The above theorem can be used to construct an example which shows that multiplicative faithfulness is not a quasi-isomorphism invariant for torsion free groups. Let $A = \{(m/3^k)a \mid m, k \in \mathbb{Z}\}$, $B = \{(m/(11)^k)b \mid m, k \in \mathbb{Z}\}$, and let $G = \langle A \oplus B, a + 2b/61 \rangle$. Then $\Pi_A = \{3\}$, $\Pi_B = \{11\}$ and $2 \notin \langle \Pi_A \cup \Pi_B \rangle \subseteq \mathbb{Z}(61)^*$. G is not multiplicatively faithful by Theorem 5. $A \oplus B$ is multiplicatively faithful by Theorem 4. G is quasi-isomorphic to $A \oplus B$, since $G \cong A \oplus B \cong 61G$.

We give a name to a common occurrence for torsion free groups.

DEFINITION 2. Let p be a prime and A a rank one subgroup of a torsion free group G . A is called p -dense in G iff $p(G/A) = G/A$ and G is p -reduced.

THEOREM 6. Let A be p -dense in G for some prime p . Let $0 \neq a \in A$ and let $\Delta, \Gamma \in \text{Mult } G$ be such that $a_\Delta^A = a_\Gamma^A$. Then $\Delta = \Gamma$.

Proof. Since A is p -dense, $\text{Hom}(G/A \otimes G, G) = (0)$. But then also $\text{Hom}(G/\langle a \rangle \otimes G, G) = (0)$, since $A/\langle a \rangle \otimes G$ is the torsion subgroup of $G/\langle a \rangle \otimes G$ and G is torsion free.

The exact sequence: $0 \rightarrow G \xrightarrow{a \otimes -} G \otimes G \rightarrow G/\langle a \rangle \otimes G \rightarrow 0$ yields: $0 \rightarrow \text{Hom}(G/\langle a \rangle \otimes G, G) \rightarrow \text{Mult } G \xrightarrow{\theta} E(G)$, where θ is given by $\theta(\Delta) = \Delta \circ (a \otimes -) = a_\Delta^A \in E(G)$. Since $\text{Hom}(G/\langle a \rangle \otimes G, G) = (0)$, θ is 1 - 1.

REFERENCES

1. R. A. Beaumont and R. S. Pierce, *Torsion Free Groups of Rank Two*, Amer. Math. Soc., Mem. 38, Amer. Math. Soc., Providence, R. I., 1961.
2. L. Fuchs, *Infinite Abelian Groups*, v. I-II, Academic Press, New York, 1970, 1973.

Received October 3, 1975 and in revised form November 5, 1975.

THE UNIVERSITY OF CONNECTICUT

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 63, No. 1

March, 1976

Ralph Artino, <i>Gevrey classes and hypoelliptic boundary value problems</i>	1
B. Aupetit, <i>Caractérisation spectrale des algèbres de Banach commutatives</i>	23
Leon Bernstein, <i>Fundamental units and cycles in the period of real quadratic number fields. I</i>	37
Leon Bernstein, <i>Fundamental units and cycles in the period of real quadratic number fields. II</i>	63
Robert F. Brown, <i>Fixed points of automorphisms of compact Lie groups</i>	79
Thomas Ashland Chapman, <i>Concordances of noncompact Hilbert cube manifolds</i>	89
William C. Connett, V and Alan Schwartz, <i>Weak type multipliers for Hankel transforms</i>	125
John Wayne Davenport, <i>Multipliers on a Banach algebra with a bounded approximate identity</i>	131
Gustave Adam Efroymsen, <i>Substitution in Nash functions</i>	137
John Sollion Hsia, <i>Representations by spinor genera</i>	147
William George Kitto and Daniel Eliot Wulbert, <i>Korovkin approximations in L_p-spaces</i>	153
Eric P. Kronstadt, <i>Interpolating sequences for functions satisfying a Lipschitz condition</i>	169
Gary Douglas Jones and Samuel Murray Rankin, III, <i>Oscillation properties of certain self-adjoint differential equations of the fourth order</i>	179
Takaši Kusano and Hiroshi Onose, <i>Nonoscillation theorems for differential equations with deviating argument</i>	185
David C. Lantz, <i>Preservation of local properties and chain conditions in commutative group rings</i>	193
Charles W. Neville, <i>Banach spaces with a restricted Hahn-Banach extension property</i>	201
Norman Oler, <i>Spaces of discrete subsets of a locally compact group</i>	213
Robert Olin, <i>Functional relationships between a subnormal operator and its minimal normal extension</i>	221
Thomas Thornton Read, <i>Bounds and quantitative comparison theorems for nonoscillatory second order differential equations</i>	231
Robert Horace Redfield, <i>Archimedean and basic elements in completely distributive lattice-ordered groups</i>	247
Jeffery William Sanders, <i>Weighted Sidon sets</i>	255
Aaron R. Todd, <i>Continuous linear images of pseudo-complete linear topological spaces</i>	281
J. Jerry Uhl, Jr., <i>Norm attaining operators on $L^1[0, 1]$ and the Radon-Nikodým property</i>	293
William Jennings Wickless, <i>Abelian groups in which every endomorphism is a left multiplication</i>	301