

Pacific Journal of Mathematics

ABSOLUTE NÖRLUND SUMMABILITY FACTORS FOR FOURIER SERIES

G. D. DIKSHIT

ABSOLUTE NÖRLUND SUMMABILITY FACTORS FOR FOURIER SERIES

G. D. DIKSHIT

In this paper a general theorem on the absolute Nörlund summability factors of a Lebesgue-Fourier series at a given point has been established. The theorem exhibits the potency of a Nörlund method as a tool to study absolute summability and the absolute convergence of Fourier series. Several existing results in the field are deduced as special cases. This also shows some sort of continuity amongst these theorems which otherwise seem apparently to be disconnected.

1. Let $\{p_n\}$ be a sequence of constants such that

$$P_n = p_0 + p_1 + \cdots + p_n \neq 0, \quad \text{for } n = 0, 1, 2, \dots.$$

Given a series $\sum u_n$ we define $\{t_n\}$ of its Nörlund means by

$$t_n = \frac{1}{P_n} \sum_0^n P_k u_{n-k}.$$

The series $\sum u_n$ is said to be summable $|N, p_n|$ and we write $\sum u_n \in |N, p_n|$, if the sequence $\{t_n\} \in bv$, that is $\sum |t_n - t_{n+1}|$ is convergent.

In the special case when $p_n = \Gamma(n+k)/\Gamma(k)\Gamma(n+1)$, $k > 0$, the summability $|N, p_n|$ reduces to the familiar Cesàro summability $|C, k|$ and when $p_n = 1/(n+1)$, it is the same as the absolute harmonic summability.

2. Let f be a periodic function with period 2π and let $f \in L(-\pi, \pi)$. Let the Fourier series of f at a point x be given by

$$f(x) \sim \frac{1}{2}a_0 + \sum_1^\infty (a_n \cos nx + b_n \sin nx) \equiv \sum_0^\infty A_n(x).$$

Throughout the paper we use the following notations:

$$\phi(t) = \frac{1}{2}\{f(x+t) + f(x-t)\},$$

$$\Phi_0(t) = \phi(t),$$

$$\Phi_\alpha(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-u)^{\alpha-1} \phi(u) du, \quad \alpha > 0,$$

$$\phi_\alpha(t) = \Gamma(\alpha+1)t^{-\alpha}\Phi_\alpha(t), \quad \alpha \geq 0,$$

$$G(n, t) = P_n^{-1} \sum_0^{n-1} p_k \varepsilon_{n-k} e^{i(n-k)t},$$

$$\begin{aligned}
 g(n, t) &= \operatorname{Im} (G(n, t)) , \\
 H(n, t) &= \frac{P_\tau}{P_n} \left(|\varepsilon_{n-m}| + \frac{1}{n} \sum_{k=m}^{n+1} |\varepsilon_k| \right) + \frac{p_m}{t P_n} , \\
 &\quad \text{where } \tau = [1/t] \text{ and } m = [(1/2)n] , \\
 J(n, t) &= \frac{1}{\Gamma(1-\alpha)} \int_t^\pi (w-t)^{-\alpha} \frac{d}{dw} g(n, w) dw, \quad 0 \leq \alpha < 1 , \\
 L(n, t) &= \frac{1}{\Gamma(\alpha+1)} \int_t^\pi u^\alpha \frac{d}{du} J(n, u) du, \quad 0 \leq \alpha < 1 , \\
 V(n, t) &= \frac{1}{\Gamma(\alpha+1)} \int_0^t u^\alpha \frac{d}{du} J(n, u) du, \quad 0 \leq \alpha < 1 .
 \end{aligned}$$

K, K_1, K_2, \dots , denote absolute constants not necessarily the same at different occurrences.

B denotes the class of bounded sequences and

\mathcal{M} denotes a class of positive and nonincreasing sequences:

$$\mathcal{M} = \left\{ s_n : s_n > 0, \frac{s_{n+1}}{s_n} \leq \frac{s_{n+2}}{s_{n+1}} \leq 1 \right\} .$$

3. Since the publication of the classical theorems of Bosanquet ([3], [4]) on absolute Cesàro summability of a Fourier series in 1936, various results have been worked out on absolute Cesàro summability, absolute harmonic summability and absolute Nörlund summability of Fourier series and series related with it. The purpose of this paper is to furnish a general theorem on the absolute Nörlund summability of a Fourier series from which we deduce several known and unknown results.

We establish the following theorem:

THEOREM. *Let α satisfy $0 \leq \alpha < 1$ and let $\{p_n\} \in \mathcal{M}$ and $\{\varepsilon_n\} \in bv$ be such that*

$$(i) \quad \sum_n^\infty |\Delta \varepsilon_k| = O(|\varepsilon_n|) \text{ and}$$

$$(ii) \quad \left\{ \frac{P_n}{n^\alpha} \sum_n^\infty \frac{|\varepsilon_k|}{k^{1-\alpha} P_k} \right\} \in B.$$

If $\phi_\alpha(t) \in BV(0, \pi)$, then $\sum A_n(x) \varepsilon_n \in |N, p_n|$.

4. We use the results in the following lemmas towards the proof of our theorem.

LEMMA 1 (McFadden [13]). *If $\{p_n\}$ is a nonnegative nonincreasing sequence then for $0 < t \leq \pi$ and for any n, a and b*

$$\left| \sum_a^b p_k e^{i(n-k)t} \right| \leq KP_{[1/t]} .$$

LEMMA 2. Let $\{p_n\}$ be a nonnegative nonincreasing sequence and $\{\varepsilon_n\} \in bv$ be such that $\sum_n^{2n} |\Delta\varepsilon_k| = O(|\varepsilon_n|)$. Then for $1/n < t \leq \pi$, $\tau = [1/t]$ and $m = [(1/2)n]$,

(i)
$$G(n, t) = O\left(\frac{P_\tau |\varepsilon_{n-m}|}{P_n}\right) + O\left(\frac{p_m}{tP_n}\right);$$

(ii)
$$\frac{d}{dt}G(n, t) = O(nH(n, t));$$

and

(iii)
$$L(n, t) = O(n^{\alpha t} H(n, t)) .$$

Proof. (i) Let

$$\begin{aligned} P_n G(n, t) &= \left(\sum_0^m + \sum_{m+1}^{n-1} \right) p_k \varepsilon_{n-k} e^{i(n-k)t} \\ &= S_1 + S_2 . \end{aligned}$$

Then by Lemma 1

$$\begin{aligned} S_1 &= O\left(P_\tau \left\{ |\varepsilon_{n-m}| + \sum_0^{m-1} |\Delta\varepsilon_{n-k}| \right\}\right) \\ &= O(P_\tau |\varepsilon_{n-m}|) , \end{aligned}$$

and

$$\begin{aligned} S_2 &= \sum_1^{n-m-1} p_{n-k} \varepsilon_k e^{ikt} \\ &= O\left(\frac{1}{t}\right) \left\{ p_{m+1} + \sum_1^{n-m-2} |\Delta(\varepsilon_k p_{n-k})| \right\} \\ &= O\left(\frac{1}{t}\right) \left\{ p_{m+1} + \sum_1^{n-m-2} |\varepsilon_k| (p_{n-k-1} - p_{n-k}) + \sum_1^{n-m-2} p_{n-k-1} |\Delta\varepsilon_k| \right\} \\ &= O\left(\frac{p_m}{t}\right) , \end{aligned}$$

by hypotheses.

(ii) Let

$$\begin{aligned} P_n \frac{d}{dt}G(n, t) &= \left(\sum_0^m + \sum_{m+1}^{n-1} \right) p_k \varepsilon_{n-k} (n-k) i e^{i(n-k)t} \\ &= S_3 + S_4 . \end{aligned}$$

Then, proceeding as above in (i)

$$\begin{aligned}
 S_3 &= O(P_\tau) \left\{ (n-m) |\varepsilon_{n-m}| + \sum_0^{m-1} |\Delta((n-k)\varepsilon_{n-k})| \right\} \\
 &= O(P_\tau) \left\{ n |\varepsilon_{n-m}| + \sum_{n-m}^{n-1} k |\Delta \varepsilon_k| + \sum_{n-m}^{n-1} |\varepsilon_{k+1}| \right\} \\
 &= O(P_\tau) \left\{ n |\varepsilon_{n-m}| + \sum_{n-m}^n |\varepsilon_k| \right\}; \\
 S_4 &= \sum_1^{n-m-1} k \varepsilon_k p_{n-k} i e^{\varepsilon_k t} \\
 &= O(1/t) \left\{ (n-m-1) |\varepsilon_{n-m-1}| p_{m+1} + \sum_1^{n-m-2} |\Delta(k \varepsilon_k p_{n-k})| \right\} \\
 &= O(1/t) \left\{ n p_m + \sum_1^{n-m-2} k |\Delta(\varepsilon_k p_{n-k})| + \sum_1^{n-m-1} p_{n-k} |\varepsilon_k| \right\} \\
 &= O(n/t) \left\{ p_m + \sum_1^{n-m-2} |\Delta(\varepsilon_k p_{n-k})| \right\} \\
 &= O(n p_m / t).
 \end{aligned}$$

(iii) We have

$$\begin{aligned}
 \Gamma(1-\alpha) J(n, t) &= \left(\int_t^{\varepsilon^{t+1/n}} + \int_{\varepsilon^{t+1/n}}^\pi \right) (u-t)^{-\alpha} \frac{d}{du} g(n, u) du \\
 &= I_1 + I_2, \quad \text{say.} \\
 I_1 &= O(n H(n, t)) \int_t^{\varepsilon^{t+1/n}} (u-t)^{-\alpha} du, \quad \text{by (ii),} \\
 &= O(n^\alpha H(n, t)),
 \end{aligned}$$

and

$$\begin{aligned}
 I_2 &= O(n^\alpha) | [g(n, u)]_{\varepsilon^{t+1/n}}^{\pi'} |, \quad t + \frac{1}{n} < \pi' \leq \pi, \\
 &= O\left(\frac{n^\alpha}{P_n}\right) \left\{ P_\tau |\varepsilon_{n-m}| + \frac{p_m}{t} \right\}, \quad \text{by (i).}
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \Gamma(\alpha+1) |L(n, t)| &= |[u^\alpha J(n, u)]_t^\pi - \alpha \int_t^\pi u^{\alpha-1} J(n, u) du| \\
 &= \left| t^\alpha J(n, t) + \frac{\alpha}{\Gamma(1-\alpha)} \int_t^\pi u^{\alpha-1} \int_u^\pi (w-u)^{-\alpha} \frac{d}{dw} g(n, w) dw du \right| \\
 &= \left| t^\alpha J(n, t) + \frac{\alpha}{\Gamma(1-\alpha)} \int_t^\pi \frac{d}{dw} g(n, w) dw \int_t^w (w-u)^{-\alpha} u^{\alpha-1} du \right| \\
 &= \left| t^\alpha J(n, t) + \frac{\alpha}{\Gamma(1-\alpha)} \int_t^\pi \frac{d}{dw} g(n, w) dw \int_{t/w}^1 u^{\alpha-1} (1-u)^{-\alpha} du \right| \\
 &\leq K_1 t^\alpha |J(n, t)| + K_2 |g(n, t')|, \quad t \leq t' < \pi.
 \end{aligned}$$

$$\Rightarrow L(n, t) = O(n^\alpha t^\alpha H(n, t)).$$

LEMMA 3 (Das [6]). *Let $\{p_n\} \in \mathcal{M}$. Then $\sum u_n \in |N, p_n|$, if and only if*

$$\sum_1^\infty \frac{1}{nP_n} \left| \sum_1^n p_{n-k} k u_k \right| < \infty .$$

5. *Proof of the Theorem.* Let

$$T_n \equiv T_n(x) = \frac{1}{P_n} \sum_1^n p_{n-k} k A_k(x) \varepsilon_k .$$

Then by Lemma 3, it is sufficient to show that

$$\sum \frac{|T_n|}{n} < \infty .$$

As

$$\begin{aligned} kA_k(x) &= \frac{2}{\pi} \int_0^\pi \phi(t) \frac{d}{dt} \sin ktdt , \\ \frac{\pi}{2} T_n &= \frac{1}{P_n} \sum_1^n p_{n-k} \varepsilon_k \int_0^\pi \phi(t) \frac{d}{dt} \sin ktdt \\ &= \int_0^\pi \phi(t) \frac{d}{dt} g(n, t) dt \\ &= \frac{1}{\Gamma(1-\alpha)} \int_0^\pi \frac{d}{dt} g(n, t) \int_0^t (t-u)^{-\alpha} d\Phi_\alpha(u) dt, \quad 0 < \alpha < 1 , \\ &= \frac{1}{\Gamma(1-\alpha)} \int_0^\pi d\Phi_\alpha(u) \int_u^\pi (t-u)^{-\alpha} \frac{d}{dt} g(n, t) dt \\ &= - \int_0^\pi \Phi_\alpha(u) \frac{d}{du} J(n, u) du, \quad 0 \leq \alpha < 1 , \\ &= - [\phi_\alpha(u) V(n, d)]_0^\pi + \int_0^\pi V(n, u) d\phi_\alpha(u) \\ &= -\phi_\alpha(\pi) V(n, \pi) + \int_0^\pi V(n, u) d\phi_\alpha(u) . \end{aligned}$$

If in particular we choose $\phi(t) = 1$, then $\phi_\alpha(t) = 1$ and $T_n = 0$ for every n . Hence

$$V(n, \pi) = 0 .$$

Thus

$$\frac{\pi}{2} T_n = \int_0^\pi V(n, u) d\phi_\alpha(u) .$$

As $\phi_\alpha(t) \in BV(0, \pi)$, to complete the proof it is sufficient to show that uniformly in $t, 0 < t \leq \pi$

$$\sum \frac{|V(n, t)|}{n} \leq K.$$

Since $V(n, t) + L(n, t) = 0$, we have

$$\sum \frac{|V(n, t)|}{n} \leq \sum_1^\tau \frac{|V(n, t)|}{n} + \sum_{\tau+1}^\infty \frac{|L(n, t)|}{n}.$$

As

$$|g(n, t)| \leq \frac{1}{P_n} \sum_0^{n-1} p_k |\varepsilon_{n-k}| \leq K,$$

and

$$\left| \frac{d}{dt} g(n, t) \right| \leq \frac{1}{P_n} \sum_0^{n-1} p_k (n - k) |\varepsilon_{n-k}| \leq Kn,$$

we have

$$\begin{aligned} \Gamma(1 - \alpha)J(n, t) &= \left(\int_t^{t+1/n} + \int_{t+1/n}^\pi \right) (u - t)^{-\alpha} \frac{d}{du} g(n, u) du \\ &= \text{etc.} \\ &= O(n^\alpha). \end{aligned}$$

Hence, for $0 < \alpha < 1$,

$$\Gamma(1 + \alpha)|V(n, t)| = \left| [u^\alpha J(n, u)]_0^t - \alpha \int_0^t u^{\alpha-1} J(n, u) du \right| \leq Kn^{\alpha t},$$

and

$$\sum_1^\tau \frac{|V(n, t)|}{n} \leq Kt^\alpha \sum_1^\tau n^{\alpha-1} \leq K.$$

For $\alpha = 0$, we note that

$$|V(n, t)| = |g(n, t)| \leq \frac{1}{P_n} \sum_0^{n-1} p_k |\varepsilon_{n-k}| (n - k)t \leq nt,$$

and thus again

$$\sum_1^\tau \frac{|V(n, t)|}{n} \leq K.$$

Therefore, after Lemma 2(iii), it is sufficient to show that

$$t^\alpha \sum_{\tau+1}^\infty n^{\alpha-1} |H(n, t)| \leq K,$$

uniformly in $t, 0 < t \leq \pi$. However

$$\begin{aligned}
 & t^\alpha \sum_{\tau+1}^{\infty} n^{\alpha-1} |H(n, t)| \\
 &= P_\tau t^\alpha \sum_{\tau+1}^{\infty} \frac{n^{\alpha-1}}{P_n} |\varepsilon_{n-m}| + P_\tau t^\alpha \sum_{\tau+1}^{\infty} \frac{n^{\alpha-2}}{P_n} \sum_{n-m}^{n+1} |\varepsilon_k| + t^{\alpha-1} \sum_{\tau+1}^{\infty} \frac{n^{\alpha-1} p_m}{P_n} \\
 &\leq K_1 + P_\tau t^\alpha \sum_{[(\tau+1)/2]}^{\infty} |\varepsilon_k| \sum_{n=k-1}^{2k} \frac{n^{\alpha-2}}{P_n} + K_2 t^{\alpha-1} \sum_{\tau+1}^{\infty} n^{\alpha-2} \\
 &\leq K_1 + K_2 P_\tau t^\alpha \sum_{[(\tau+1)/2]}^{\infty} \frac{|\varepsilon_k|}{k^{1-\alpha} P_k} \\
 &\leq K.
 \end{aligned}$$

This completes the proof of the theorem.

6. COROLLARIES.

6.1. Taking $p_n = \Gamma(n + \beta)/\Gamma(\beta)\Gamma(n + 1), 1 \geq \beta > 0$, we get

COROLLARY 1. *Let α satisfy $0 \leq \alpha < 1$, and let $\{\varepsilon_n\} \in bv$ be such that*

- (i) $\sum_n^\infty |\Delta \varepsilon_k| = O(|\varepsilon_n|)$ and
- (ii) $\left\{ n^{\beta-\alpha} \sum_n^\infty \frac{|\varepsilon_k|}{k^{1+\beta-\alpha}} \right\} \in B$.

If $\phi_\alpha(t) \in BV(0, \pi)$, then $\sum A_n(x)\varepsilon_n \in |C, \beta|, \beta \geq \alpha$.

The case $\{\varepsilon_n\} \equiv \{1\}$, and $\beta > \alpha$, furnishes corresponding results due to Bosanquet ([3], [4]). Taking $0 < \alpha = \beta < 1$ and specializing $\{\varepsilon_n\}$ to $\{(\log(n + 1))^{-1-\varepsilon}\}, \varepsilon > 0$, we get a result due to Cheng ([5]). Again in the case $\alpha = \beta$, taking $\{\varepsilon_n\}$ to be a convex sequence (see Zygmund [18], p. 93, for the definition and certain properties as needed) we obtain a result due to Prasad and Bhatt ([14]). The case $\beta = \alpha = 0$ is covered in Corollary 3 below.

6.2. The case $p_n = 1/(n + 1)$ furnishes the following result on absolute harmonic summability:

COROLLARY 2. *Let α satisfy $0 \leq \alpha < 1$, and let $\{\varepsilon_n\} \in bv$ be such that*

- (i) $\sum_n^\infty |\Delta \varepsilon_k| = O(|\varepsilon_n|)$ and
- (ii) $\left\{ \frac{\log n}{n^\alpha} \sum_n^\infty \frac{|\varepsilon_k|}{k^{1-\alpha} \log k} \right\} \in B$.

If $\phi_\alpha(t) \in BV(0, \pi)$, then $\sum A_n(x)\varepsilon_n \in |N, 1/n + 1|$.

The case $\alpha = 0$, includes a well known theorem due to Varshney ([17], Varshney has proved the result for $\{\varepsilon_n\} = \{1/\log(n+2)\}$). Specialising $\{\varepsilon_n\}$ to be $\{\log(n+1)\lambda_n/n^\alpha\}$, where $\{\lambda_n\}$ is a convex sequence, we get the result due to Bhatt [1].

6.3. It is now known (see Dikshit [8]) that if $\{p_n\} \in \mathcal{M}$ and $\{P_n\} \in B$ then the method $|N, p_n|$ is ineffective, in the sense that it sums only absolutely convergent series. Thus the extra hypothesis that $\{P_n\} \in B$ in the theorem yields the following result on absolute convergence factors for Fourier series.

COROLLARY 3. *Let α satisfy $0 \leq \alpha < 1$, and let $\{\varepsilon_n\} \in bv$ be such that*

$$(i) \quad \sum_n^\infty |\Delta \varepsilon_k| = O(|\varepsilon_n|) \text{ and}$$

$$(ii) \quad \left\{ \frac{1}{n^\alpha} \sum_n^\infty \frac{|\varepsilon_k|}{k^{1-\alpha}} \right\} \in B.$$

If $\phi_\alpha(t) \in BV(0, \pi)$, then $\sum A_n(x)\varepsilon_n$ is absolutely convergent.

Results in somewhat weaker form are eventually known in as much as they could be deduced from the theorem of Cheng [5], Prasad and Bhatt [14], or the Corollary 2, with an application of a result of Kogbetliantz ([11]).

6.4. The case $\{\varepsilon_n\} \equiv \{1\}$ yields the following:

COROLLARY 4. *Let $0 \leq \alpha < 1$ and let $\{p_n\} \in \mathcal{M}$ and be such that*

$$\left\{ \frac{P_n}{n^\alpha} \sum_n^\infty \frac{1}{k^{1-\alpha} P_k} \right\} \in B.$$

If $\phi_\alpha(t) \in BV(0, \pi)$, then $\sum A_n(x) \in |N, p_n|$.

A more general result in this direction is also known and is given elsewhere (Dikshit [7], [9]; see also Lal [12]).

6.5. Writing $\{\varepsilon_n\} = \{P_n \lambda_n / n^\alpha\}$ we obtain:

COROLLARY 5. *Let α satisfy $0 \leq \alpha < 1$, and let $\{p_n\} \in \mathcal{M}$ and $\{\lambda_n\}$ be a sequence such that*

$$(i) \quad \left\{ \frac{P_n \lambda_n}{n^\alpha} \right\} \in bv,$$

$$(ii) \quad \sum_n^\infty \left| \Delta \left(\frac{P_k \lambda_k}{k^\alpha} \right) \right| = O \left(\frac{P_n |\lambda_n|}{n^\alpha} \right) \text{ and}$$

$$(iii) \left\{ \frac{P_n}{n^\alpha} \sum_n \frac{|\lambda_k|}{k} \right\} \in B.$$

If $\phi_\alpha(t) \in BV(0, \pi)$, then $\sum A_n(x) P_n \lambda_n / n^\alpha \in |N, p_n|$.

It is worthwhile to compare the result of this Corollary for $\alpha = 0$ with one due to T. Singh [16] and L. B. Singh [15] and for $0 < \alpha < 1$, with those due to Nand Kishore [10] and Bhatt [2].

The author would like to thank the referee for the very valuable suggestions.

REFERENCES

1. S. N. Bhatt, *On the absolute summability factors of Fourier series*, Annali di Matematica pura ed applicata (4), **72** (1966), 253-266.
2. ———, *A note on the absolute Nörlund summability of a factored Fourier series*, Indian J. Math., **11** (1969), 37-42.
3. L. S. Bosanquet, *Note on the absolute summability (C) of a Fourier series*, J. London Math. Soc., **11** (1936), 11-15.
4. ———, *The absolute Cesàro summability of a Fourier series*, Proc. London Math. Soc., (2), **41** (1936), 517-528.
5. M. T. Cheng, *Summability factors of Fourier series at a given point*, Duke Math. J., **15** (1948), 29-36.
6. G. Das, *Tauberian theorems for absolute Nörlund summability*, Proc. London Math. Soc., (3), **19** (1969), 357-384.
7. G. D. Dikshit, *On the absolute Nörlund summability of a Fourier series*, Indian J. Math., **9** (1967), 331-342.
8. ———, *A note on Riesz and Nörlund means*, Rendiconti del Circolo Mat. di Palermo (2), **18** (1969), 49-61.
9. ———, *Absolute Nörlund summability of Fourier series*, Mathematical Chronicle, **2** (1973), 117-124.
10. Nand Kishore, *Absolute Nörlund summability of a factored Fourier series*, Indian J. Math., **9** (1967), 123-136.
11. E. Kogbetliantz, *Sur les séries absolument sommable par la méthode des moyennes arithmétiques*, Bull. des. Sc. Math., (2), **49** (1925), 234-256.
12. S. N. Lal, *On the absolute Nörlund summability of a Fourier series*, Indian J. Math., **13** (1971), 51-67.
13. L. McFadden, *Absolute Nörlund summability*, Duke Math. J., **9** (1942), 168-207.
14. B. N. Prasad and S. N. Bhatt, *The summability factors of a Fourier series*, Duke Math. J., **24** (1957), 103-117.
15. L. B. Singh, *Absolute Nörlund summability of a factored Fourier series*, Indian J. Math., **12** (1970), 189-194.
16. T. Singh, *Absolute Nörlund summability of a factored Fourier series*, Indian J. Math., **9** (1967), 227-236.
17. O. P. Varshney, *On the absolute harmonic summability of a series related to a Fourier series*, Proc. American Math. Soc., **10** (1959), 784-789.
18. A. Zygmund, *Trigonometric series*, Vol. 1, Cambridge University Press, New York, 1959.

Received October 8, 1975 and in revised form January 9, 1976.

UNIVERSITY OF AUCKLAND, NEW ZEALAND

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Joseph Anthony Ball and Arthur R. Lubin, <i>On a class of contractive perturbations of restricted shifts</i>	309
Joseph Becker and William C. Brown, <i>On extending higher derivations generated by cup products to the integral closure</i>	325
Andreas Blass, <i>Exact functors and measurable cardinals</i>	335
Joseph Eugene Collison, <i>A variance property for arithmetic functions</i>	347
Craig McCormack Cordes, <i>Quadratic forms over nonformally real fields with a finite number of quaternion algebras</i>	357
Freddy Delbaen, <i>Weakly compact sets in H^1</i>	367
G. D. Dikshit, <i>Absolute Nörlund summability factors for Fourier series</i>	371
Edward Richard Fadell, <i>Nielsen numbers as a homotopy type invariant</i>	381
Josip Globevnik, <i>Analytic extensions of vector-valued functions</i>	389
Robert Gold, <i>Genera in normal extensions</i>	397
Solomon Wolf Golomb, <i>Formulas for the next prime</i>	401
Robert L. Griess, Jr., <i>The splitting of extensions of $SL(3, 3)$ by the vector space F_3^3</i>	405
Thomas Alan Keagy, <i>Matrix transformations and absolute summability</i>	411
Kazuo Kishi, <i>Analytic maps of the open unit disk onto a Gleason part</i>	417
Kwangil Koh, Jiang Luh and Mohan S. Putcha, <i>On the associativity and commutativity of algebras over commutative rings</i>	423
James C. Lillo, <i>Asymptotic behavior of solutions of retarded differential difference equations</i>	431
John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues</i>	445
Anna Maria Mantero, <i>Sets of uniqueness and multiplicity for L^p</i>	467
J. F. McClendon, <i>Embedding metric families</i>	481
L. Robbiano and Giuseppe Valla, <i>Primary powers of a prime ideal</i>	491
Wolfgang Ruess, <i>Generalized inductive limit topologies and barrelledness properties</i>	499
Judith D. Sally, <i>Bounds for numbers of generators of Cohen-Macaulay ideals</i>	517
Helga Schirmer, <i>Mappings of polyhedra with prescribed fixed points and fixed point indices</i>	521
Cho Wei Sit, <i>Quotients of complete multipartite graphs</i>	531
S. Sznajder and Zbigniew Zielezny, <i>Solvability of convolution equations in \mathcal{H}'_p, $p > 1$</i>	539
Mitchell Herbert Taibleson, <i>The existence of natural field structures for finite dimensional vector spaces over local fields</i>	545
William Yslas Vélez, <i>A characterization of completely regular fields</i>	553
P. S. Venkatesan, <i>On right unipotent semigroups</i>	555
Kenneth S. Williams, <i>A rational octic reciprocity law</i>	563
Robert Ross Wilson, <i>Lattice orderings on the real field</i>	571
Harvey Eli Wolff, <i>V-localizations and V-monads. II</i>	579