

# Pacific Journal of Mathematics

**FORMULAS FOR THE NEXT PRIME**

SOLOMON WOLF GOLOMB

## FORMULAS FOR THE NEXT PRIME

SOLOMON W. GOLOMB

In 1971, J. M. Gandhi showed that if the first  $n$  primes,  $p_1, p_2, \dots, p_n$  are known, then the next prime,  $p_{n+1}$ , is given "explicitly" by the formula:

$$(1) \quad 1 < b^t \left( \sum_{d|P_n} \frac{\mu(d)}{b^d - 1} - \frac{1}{b} \right) < b,$$

where  $b$  is any positive integer  $\geq 2$ , where  $P_n = p_1 p_2 \cdots p_n$ , where  $\mu(d)$  is the Möbius function, and where the unique integer value of  $t$  which satisfies the indicated inequalities is in fact  $p_{n+1}$ .

In this paper, we obtain of the following formulas for  $p_{n+1}$ :

$$(2) \quad p_{n+1} = \lim_{s \rightarrow \infty} \{P_n(s)\zeta(s) - 1\}^{-1/s}$$

$$(3) \quad p_{n+1} = \lim_{s \rightarrow \infty} \{P_n(s) - \zeta^{-1}(s)\}^{-1/s}$$

$$(4) \quad p_{n+1} = \lim_{s \rightarrow \infty} \{\zeta(s) - Q_n(s)\}^{-1/s}$$

and

$$(5) \quad p_{n+1} = \lim_{s \rightarrow \infty} \{1 - \zeta^{-1}(s)Q_n(s)\}^{-1/s}.$$

Here  $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$  for (real)  $s > 1$  is the Riemann Zeta Function, with  $\zeta^{-1}(s) = \sum_{n=1}^{\infty} \mu(n)/n^s$ ;  $P_n(s) = \prod_{p_i|P_n} (1 - p_i^{-s})$ , and  $Q_n(s) = \{P_n(s)\}^{-1} = \sum_{n=1}^{\infty} n^{-s}$ , where the *prime* indicates that summation is extended over those values of  $n$  having no prime factors exceeding  $p_n$ .

The approach to be followed here involves the derivation of a more general formula, based on the notion of probability distributions on the positive integers, from which both the Gandhi formula and the new formulas listed above follow as special cases.

2. Probability formulas for the integers. Let  $\alpha(n)$  be a probability function on the positive integers. That is,  $\alpha(n) \geq 0$  for all  $n = 1, 2, 3, \dots$ , and  $\sum_{n=1}^{\infty} \alpha(n) = 1$ .

Let  $\beta(m) = \sum_{n=1}^{\infty} \alpha(mn)$ . In the probability distribution  $D$  determined by  $\{\alpha(n)\}$ ,  $\beta(m)$  is the probability that a randomly chosen integer is a multiple of  $m$ . Next, let  $\gamma(k) = \sum_{d|k} \mu(d)\beta(d)$ . Then  $\gamma(k)$  is the probability (in  $D$ ) that a randomly chosen integer is relatively prime to  $k$ , because

$$\gamma(k) = 1 - \sum_{p_i|k} \beta(p_i) + \sum_{p_i p_j|k} \beta(p_i p_j) - + \cdots.$$

Let  $P_n = p_1 p_2 \cdots p_n$  be the product of the first  $n$  primes. Then

$\gamma(P_n) = \sum_{d|P_n} \mu(d)\beta(d)$  by the definition of  $\gamma(k)$ ; but also

$$(6) \quad \gamma(P_n) = \alpha(1) + \alpha(p_{n+1}) + \cdots = \sum_{j=1}^{\infty} \alpha(j),$$

where  $\sum''$  indicates summation over all positive integers divisible by none of the first  $n$  primes.

**THEOREM 1.** *Suppose  $1 \leq n_1 < n_2 < n_3 < \cdots$  is any subsequence of the positive integers, and there exists an operator  $T$  such that*

$$T\left(\sum_{i=1}^{\infty} \alpha(n_i)\right) = n_1$$

for all such subsequences. Then

$$(7) \quad T(\gamma(P_n) - \alpha(1)) = p_{n+1}$$

is a "formula" for the next prime,  $p_{n+1}$ .

*Proof.* Since  $\gamma(P_n) - \alpha(1) = \sum''_{j>1} \alpha(j) = \alpha(p_{n+1}) + \cdots$ , we have  $T(\gamma(P_n) - \alpha(1)) = p_{n+1}$  by the hypothesis concerning the operator  $T$ .

Another general result is given by:

**THEOREM 2.**

$$(8) \quad \gamma(0) = \lim_{n \rightarrow \infty} \gamma(P_n) = \sum_{d=1}^{\infty} \mu(d)\beta(d) = \alpha(1).$$

*Proof.* This follows directly from

$$\gamma(P_n) = \sum_{d|P_n} \mu(d)\beta(d) = \sum_{j=1}^{\infty} \alpha(j).$$

As we shall see, Theorem 2 is a generalization of Euler's product formula for the Zeta Function.

3. **Some special cases.** Suppose  $\alpha(n) = (b-1)b^{-n}$  for  $n = 1, 2, 3, \cdots$  where  $b > 1$  is a positive integer. This is a geometric distribution on the positive integers. Then

$$\beta(m) = \sum_{n=1}^{\infty} \alpha(mn) = \frac{b-1}{b^m-1},$$

and

$$\gamma(k) = \sum_{d|k} \mu(d)\beta(d) = (b-1) \sum_{d|k} \frac{\mu(d)}{b^d-1}.$$

In particular,  $\gamma(P_n) = (b - 1) \sum_{d|P_n} \mu(d)/(b^d - 1)$ , and

$$\gamma(P_n) - \alpha(1) = (b - 1) \left( \sum_{d|P_n} \frac{\mu(d)}{b^d - 1} - \frac{1}{b} \right) = (b - 1) \left\{ \frac{1}{b^{p_{n+1}}} + \dots \right\},$$

and to recover  $p_{n+1}$  it suffices to divide by  $b - 1$ , and then multiply by the smallest power  $b^t$  of  $b$ ,  $t$  an integer, such that

$$b^t \left( \sum_{d|P_n} \frac{\mu(d)}{b^d - 1} - \frac{1}{b} \right) > 1.$$

This is Gandhi's Formula (1). (For other derivations, see [1] and [2].)

Alternatively, let  $\alpha(n) = n^{-s}/\zeta(s)$  for a fixed real value of  $s$ ,  $s > 1$ . (Note that  $\sum_{n=1}^{\infty} \alpha(n) = (\sum_{n=1}^{\infty} n^{-s})/\zeta(s) = 1$ .) Then  $\beta(m) = \sum_{n=1}^{\infty} \alpha(mn) = m^{-s}$ , and  $\gamma(k) = \sum_{d|k} \mu(d)d^{-s} = \sum_{p|k} (1 - p^{-s})$ . Specifically,  $\gamma(P_n) = \sum_{d|P_n} (\mu(d))/d^s = \prod_{i=1}^n (1 - p_i^{-s}) = P_n(s)$ . Note also that  $\alpha(1) = 1/\zeta(s)$ . Thus  $\gamma(P_n) - \alpha(1) = (p_{n+1})^{-s} + \dots$ , and an appropriate operator  $T$  to recover the term  $p_{n+1}$ , in the sense of Theorem 1, is  $T = \lim_{s \rightarrow \infty} ( )^{-1/s}$ . Thus

$$(3) \quad P_{n+1} = \lim_{s \rightarrow \infty} \{P_n(s) - \zeta^{-1}(s)\}^{-1/s}.$$

Each of the formulas (2), (3), (4), (5) can be given a direct interpretation. Thus

$$(9) \quad P_n(s)\zeta(s) - 1 = \sum_1 a^{-s}$$

$$(10) \quad P_n(s) - \zeta^{-1}(s) = - \sum_2 \mu(a)a^{-s}$$

$$(11) \quad \zeta(s) - Q_n(s) = \sum_2 a^{-s}$$

$$(12) \quad 1 - \zeta^{-1}(s)Q_n(s) = - \sum_1 \mu(a)a^{-s}$$

where  $\sum_1$  indicates summation over those integers  $a > 1$  all of whose prime factors exceed  $p_n$ ; where  $\sum_2$  indicates summation over those integers  $a > 1$  having at least one prime factor exceeding  $p_n$ ; and where  $\mu(a)$  is the Möbius function. In all four of these expressions, the first surviving term in  $p_{n+1}^{-s}$ , which is recovered by the inversion operator  $T$  to yield the formulas (2), (3), (4), and (5).

For the case  $\alpha(n) = n^{-s}/\zeta(s)$ , Theorem 2 yields the identity

$$(13) \quad \gamma(0) = \prod_{i=1}^{\infty} (1 - p_i^{-s}) = \sum_{d=1}^{\infty} \mu(d) \cdot d^{-s} = 1/\sum_{n=1}^{\infty} n^{-s}$$

which includes the Euler Product Formula for the Zeta Function (cf. [3]).

The reader is invited to find other distributions on the positive integers for which Theorems 1 and 2 yield interesting formulas. A simpler proof of Gandhi's formula was given by the present author in 1974.

#### REFERENCES

1. J. M. Gandhi, Formulae for the  $N$ th prime, Proc. Washington State University Conference on Number Theory, (1971), 96-107.
2. S. W. Golomb, *A direct interpretation of Gandhi's formula*, Amer. Math. Monthly, **81** (1974), 752-754.
3. ———, *A class of probability distributions on the integers*, J. Number Theory, **2** (1970), 189-192.

Received August 13, 1975, and in revised form December 19, 1975. This research was supported in part by the U.S. Army Research Office, under Contract DA-ARO-D-31-124-73-G167.

UNIVERSITY OF SOUTHERN CALIFORNIA

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RICHARD ARENS (Managing Editor)  
University of California  
Los Angeles, California 90024

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. A. BEAUMONT  
University of Washington  
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM  
Stanford University  
Stanford, California 94305

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),  
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

# Pacific Journal of Mathematics

Vol. 63, No. 2

April, 1976

Joseph Anthony Ball and Arthur R. Lubin, <i>On a class of contractive perturbations of restricted shifts</i> .....	309
Joseph Becker and William C. Brown, <i>On extending higher derivations generated by cup products to the integral closure</i> .....	325
Andreas Blass, <i>Exact functors and measurable cardinals</i> .....	335
Joseph Eugene Collison, <i>A variance property for arithmetic functions</i> .....	347
Craig McCormack Cordes, <i>Quadratic forms over nonformally real fields with a finite number of quaternion algebras</i> .....	357
Freddy Delbaen, <i>Weakly compact sets in <math>H^1</math></i> .....	367
G. D. Dikshit, <i>Absolute Nörlund summability factors for Fourier series</i> .....	371
Edward Richard Fadell, <i>Nielsen numbers as a homotopy type invariant</i> .....	381
Josip Globevnik, <i>Analytic extensions of vector-valued functions</i> .....	389
Robert Gold, <i>Genera in normal extensions</i> .....	397
Solomon Wolf Golomb, <i>Formulas for the next prime</i> .....	401
Robert L. Griess, Jr., <i>The splitting of extensions of <math>SL(3, 3)</math> by the vector space <math>F_3^3</math></i> .....	405
Thomas Alan Keagy, <i>Matrix transformations and absolute summability</i> .....	411
Kazuo Kishi, <i>Analytic maps of the open unit disk onto a Gleason part</i> .....	417
Kwangil Koh, Jiang Luh and Mohan S. Putcha, <i>On the associativity and commutativity of algebras over commutative rings</i> .....	423
James C. Lillo, <i>Asymptotic behavior of solutions of retarded differential difference equations</i> .....	431
John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues</i> .....	445
Anna Maria Mantero, <i>Sets of uniqueness and multiplicity for <math>L^p</math></i> .....	467
J. F. McClendon, <i>Embedding metric families</i> .....	481
L. Robbiano and Giuseppe Valla, <i>Primary powers of a prime ideal</i> .....	491
Wolfgang Ruess, <i>Generalized inductive limit topologies and barrelledness properties</i> .....	499
Judith D. Sally, <i>Bounds for numbers of generators of Cohen-Macaulay ideals</i> .....	517
Helga Schirmer, <i>Mappings of polyhedra with prescribed fixed points and fixed point indices</i> .....	521
Cho Wei Sit, <i>Quotients of complete multipartite graphs</i> .....	531
S. Sznajder and Zbigniew Zielezny, <i>Solvability of convolution equations in <math>\mathfrak{K}'_p</math>, <math>p &gt; 1</math></i> .....	539
Mitchell Herbert Taibleson, <i>The existence of natural field structures for finite dimensional vector spaces over local fields</i> .....	545
William Yslas Vélez, <i>A characterization of completely regular fields</i> .....	553
P. S. Venkatesan, <i>On right unipotent semigroups</i> .....	555
Kenneth S. Williams, <i>A rational octic reciprocity law</i> .....	563
Robert Ross Wilson, <i>Lattice orderings on the real field</i> .....	571
Harvey Eli Wolff, <i>V-localizations and V-monads. II</i> .....	579