

Pacific Journal of Mathematics

**MAPPINGS OF POLYHEDRA WITH PRESCRIBED FIXED
POINTS AND FIXED POINT INDICES**

HELGA SCHIRMER

MAPPINGS OF POLYHEDRA WITH PRESCRIBED FIXED POINTS AND FIXED POINT INDICES

HELGA SCHIRMER

The following problem is studied: If points c_k of a polyhedron and integers i_k are given, when does there exist a selfmap within a given homotopy class which has the c_k as its fixed points and the i_k as its fixed point indices? Necessary and sufficient conditions for the existence of such selfmaps are established if the selfmap is a deformation and the polyhedron is of type W , and if the selfmap is arbitrary and the polyhedron is of type S . It is further shown that there always exists a selfmap of an n -sphere ($n \geq 2$) which has arbitrarily prescribed locations and indices of its fixed points. The proofs are based on Shi Gen-Hua's construction of selfmaps with a minimum number of fixed points.

1. Introduction. It is known that an arbitrarily given closed and nonempty subset of a polyhedron of type W can always be the fixed point set of a suitable selfmap, and even of a deformation [2]. We now ask what happens if not only the locations of the fixed points, but also their indices are prescribed. More precisely, we deal with the following problem:

If the points c_k of a polyhedron and the integers i_k , where $k = 1, 2, \dots, m$, are given, when does there exist a selfmap within a given homotopy class which has the c_k as its fixed points and the i_k as its fixed point indices?

The problem is an extension of the well-known one concerning the existence of maps with a minimum number of fixed points, whose most general solution to date is due to Shi Gen-Hua [4]. We use Shi's results and methods to a considerable degree.

We first show that the number, location and indices of the fixed points of a deformation of a polyhedron can be arbitrarily prescribed with the only (obvious) condition that the sum of their fixed point indices equals the Euler characteristic of the polyhedron (Theorem 1). In the case of arbitrary selfmaps the—necessary and sufficient—conditions which the fixed point indices must satisfy are naturally more complicated, and express the fact that the number and the indices of the essential fixed point classes of a map are homotopy invariant (Theorem 2). As in Shi's work [4] the assumptions which are made about the polyhedron are more restrictive in the case of

arbitrary selfmaps than in the case of deformations, as the polyhedron has to be of type S and not only of type W . (See § 2 for definitions.)

Polyhedra of type S or W cannot be one-dimensional, but selfmaps with arbitrarily given fixed point sets (but not fixed point indices) were in [2] also constructed for all one-dimensional connected polyhedra. So one would also like to find selfmaps of one-dimensional polyhedra with prescribed locations and indices of their fixed point sets, but this cannot be done if only the conditions of Theorems 1 and 2 are assumed. The situation in the one-dimensional case seems to be much more complicated, and only a special case (where the one-dimensional polyhedron is acyclic and all fixed points are attractive or repulsive) has been settled so far [3].

Another open, and probably difficult, question arises if not only maps within a specific homotopy class, but all selfmaps of a polyhedron are considered. Only a very special case, namely the one where the polyhedron is a sphere, is considered here. We show that there always exists a selfmap of an n -sphere (where $n \geq 2$) which has arbitrarily prescribed locations and indices of its fixed points.

The beginning of the paper contains two lemmas which deal with the splitting and moving of fixed points with given indices. Together with Shi's results they permit quick proofs of the later theorems. Shi's work has been included in the recent book by R. F. Brown [1], and all references are made to this book in order to have them easily accessible. For the same reason Brown's book is used as a reference for facts about fixed point indices and fixed point classes.

2. Splitting and moving of fixed points. In this preliminary paragraph we develop, in the form of two lemmas, the tools for the proofs of the results in this paper. Some definitions are needed first.

We denote by $|K|$ a polyhedron which is the realization of a finite simplicial complex K , by σ an open simplex of $|K|$, and by $\bar{\sigma}$ its closure. The *carrier* $\kappa(x)$ of a point $x \in |K|$ is the unique simplex for which $x \in \kappa(x)$. The *star* $\text{st } \sigma$ of a simplex σ consists of all simplices which have σ as a face. σ is called a *maximal* simplex if $\sigma = \text{st } \sigma$.

We use $\Phi(f)$ to denote the fixed point set of the map $f: |K| \rightarrow |K|$. The point $c \in \Phi(f)$ is an isolated fixed point if there exists an open set U with $c \in U$ and $\bar{U} \cap \Phi(f) = c$. If $|K|$ is connected and if the *fixed point index of f on U* , called $i(|K|, f, U)$, is defined as on

p. 65 of [1], then it follows from the additivity axiom (see [1], p. 52) that $i(|K|, f, U)$ is independent of the choice of the open set U as long as $c \in U$ and $\bar{U} \cap \Phi(f) = c$. For an isolated fixed point c we therefore write $i(|K|, f, c)$ for the *index of f at c* , where $i(|K|, f, c) = i(|K|, f, U)$ for any such set U . For a point c which is contained in a maximal simplex this definition coincides with the one on p. 122 of [1].

We are now ready for the two lemmas. The first states that any isolated fixed point in a maximal simplex can be split into an arbitrary number of fixed points as long as the Lefschetz number is not changed. It is a counterpart to Lemma 6 on p. 133 of [1], where fixed points are united.

LEMMA 1 (Splitting of fixed points). *Let f be a selfmap of the polyhedron $|K|$ and let the fixed point a of f be contained in a maximal simplex σ of dimension at least two for which $\Phi(f) \cap \bar{\sigma} = a$. Then there exists, for every set of m integers i_1, i_2, \dots, i_m with $\sum_{k=1}^m i_k = i(|K|, f, a)$, a selfmap f' of $|K|$ which is homotopic to f , equals f on $|K| \setminus \sigma$, and has on σ exactly m fixed points $a_k (k = 1, 2, \dots, m)$ with indices $i(|K|, f', a_k) = i_k$.*

Proof. Choose an $\varepsilon > 0$ for which $\bar{U}(a, 5\varepsilon) = \{x \in |K| \mid d(a, x) \leq 5\varepsilon\} \subset \sigma$, where d is the metric of $|K|$. As f is continuous, there exists a $\delta > 0$ with $f(\bar{U}(a, \delta)) \subset \bar{U}(a, \varepsilon)$, and we can choose it so that $0 < \delta < \varepsilon$. Let B denote the p -ball $\bar{U}(a, \delta)$, and select points a_1, a_2, \dots, a_m in $B \setminus a$ so that $d(a, a_k)$ is constant for all $k = 1, 2, \dots, m$, and so that there exists a $\rho > 0$ with $\delta - d(a, a_k) < \rho < d(a, a_k)$ for which the $\bar{U}(a_k, \rho)$ are pairwise disjoint. Put $A_k = B \cap \bar{U}(a_k, \rho)$. Then each A_k is homeomorphic to a p -ball, $a \notin A_k$, and A_k intersects the boundary ∂B of B .

In order to construct a map with fixed points a_k of index i_k we use the notation of [1], pp. 120-121. Hence $|K|$ is imbedded into Euclidean space R^n , and

$$h_k: R^n, \sigma, a_k \longrightarrow R^n, R^p, 0$$

(which corresponds to h_1 on p. 120 of [1]) is an isometry which transforms a_k into the origin 0. A map $d'_k: \partial A_k \cap \partial B \rightarrow R^p \setminus 0$ is defined by $d'_k(y) = h_k(y) - h_k f(y)$ for all $y \in \partial A_k \cap \partial B$. As

$$\begin{aligned} |d'_k(y)| &\leq |h_k(y)| + |h_k \circ f(y)| \\ &\leq \rho + \varepsilon + \delta < 3\varepsilon, \end{aligned}$$

d'_k is a map of the form $d'_k: \partial A_k \cap \partial B \rightarrow \bar{U}(0, 3\varepsilon) \setminus 0$.

Now take a p -ball B_k with centre a_k and with $B_k \subset A_k$, and choose generators $\bar{\mu}_p \in H^{p-1}(R^p \setminus 0)$ and $\alpha_p \in H^{p-1}(\partial B_k)$ as on p. 121 of [1]. Denote by $i: U(0, 3\varepsilon) \setminus 0 \rightarrow R^p \setminus 0$ the inclusion and by $\tilde{h}_k: \partial B_k \rightarrow \partial A_k$ the homeomorphism obtained by linear projection from the centre a_k . As $\partial A_k \cap \partial B$ is contractible and as $(\partial A_k, \partial A_k \cap \partial B)$ has the homotopy extension property with respect to $\bar{U}(0, 3\varepsilon) \setminus 0$, there exists an extension $d_k: \partial A_k \rightarrow \bar{U}(0, 3\varepsilon) \setminus 0$ of d'_k for which

$$\tilde{h}_k^* d_k^* i^*(\bar{\mu}_p) = i_k \cdot \alpha_p,$$

where i^* , d_k^* and \tilde{h}_k^* are the induced homomorphisms of the appropriate cohomology groups. We define a map g'_k on ∂A_k by

$$g'_k(y) = h_k^{-1}(h_k(y) - d_k(y)).$$

Then

$$\begin{aligned} |h_k \circ g'_k(y)| &\leq |h_k(y)| + |d_k(y)| \\ &\leq \delta + 3\varepsilon < 4\varepsilon, \end{aligned}$$

therefore

$$\begin{aligned} d(a, g'_k(y)) &\leq d(a, a_k) + d(a_k, g'_k(y)) \\ &\leq \delta + 4\varepsilon < 5\varepsilon, \end{aligned}$$

and $g'_k(y) \in \sigma$. As A_k is homeomorphic to a p -ball, we can extend the map $g'_k: \partial A_k \rightarrow \sigma$ linearly from a_k to a map $g_k: A_k \rightarrow \sigma$ which has a_k as its only fixed point. Note that then $g_k(y) = f(y)$ if $y \in \partial B$. We define a map $g': \bigcup_k A_k \cup \partial B \rightarrow \sigma$ by

$$g'(x) = \begin{cases} g_k(x) & \text{if } x \in A_k, \text{ where } k = 1, 2, \dots, m, \\ f(x) & \text{if } x \in \partial B. \end{cases}$$

As the closure of the subset $(B \setminus \bigcup_k A_k)$ of B is homeomorphic to a p -ball with centre a , and as g' is defined on its boundary, we can extend g' to a map $g: B \rightarrow \sigma$ such that $g(x) = g'(x)$ if $x \in \bigcup_k A_k \cup \partial B$, and that $\{a, a_1, a_2, \dots, a_m\}$ is its fixed point set. Then the map $f_1: |K| \rightarrow |K|$ given by

$$f_1(x) = \begin{cases} f(x) & \text{if } x \in |K| \setminus B, \\ g(x) & \text{if } x \in B \end{cases}$$

has the fixed point set $\{a, a_1, a_2, \dots, a_m\}$ on σ , and is homotopic to f .

It follows from the construction of f_1 with the help of the d_k , and from pp. 120-122 of [1], that $i(|K|, f_1, a_k) = i_k$. In consequence of the homotopy axiom (see [1], p. 52) we have

$$i(|K|, f_1, a) + \sum_{k=1}^m i(|K|, f_1, a_k) = i(|K|, f, a),$$

and therefore $i(|K|, f_1, a) = 0$. Select an ε' with $0 < \varepsilon' < d(a, a_k)$ and $\bar{U}(a, 2\varepsilon') \subset \sigma$, and a δ' with $0 < \delta' < \varepsilon'$ and $f_1(\bar{U}(a, \delta')) \subset \bar{U}(a, \varepsilon')$. According to Theorem 4 on p. 123 of [1] there exists a map $f': |K| \rightarrow |K|$ with $f'(x) = f_1(x)$ for $x \in |K| \setminus \bar{U}(a, \delta')$, with $d(f_1, f') < \varepsilon'$ and which is fixed point free on $\bar{U}(a, \delta')$. As $f'(\bar{U}(a, \delta')) \subset \bar{U}(a, 2\varepsilon') \subset \sigma$, f' is homotopic to f_1 and hence to f . The set of fixed points of f' on σ is $\{a_1, a_2, \dots, a_m\}$ and the index $i(|K|, f', a_k)$ equals i_k .

The second lemma will show that isolated fixed points can be moved to arbitrarily prescribed points if $|K|$ satisfies a connectedness condition. More precisely we require that $|K|$ is of type W ([1], p. 143), i.e. that every maximal simplex of $|K|$ is of dimension at least two and that for every two maximal simplices σ, σ' of $|K|$ there exist maximal simplices $\sigma_1, \sigma_2, \dots, \sigma_r$ with $\sigma = \sigma_1, \sigma_r = \sigma'$ and $\bar{\sigma}_i \cap \bar{\sigma}_{i+1}$ of dimension at least one for $i = 1, 2, \dots, r - 1$. Lemma 2 is an extension of Lemma 6 on p. 135 of [1], as we do not assume that $\kappa(c)$ is a maximal simplex.

LEMMA 2 (Moving of fixed points). *Let $|K|$ be a polyhedron of type W , let f be a selfmap of $|K|$ with fixed point set $\Phi(f) = \{a_1, a_2, \dots, a_m\}$ and let a_1 be contained in a maximal simplex. Then there exists, for any $c \notin \Phi(f)$, a selfmap f' which is homotopic to f and for which*

$$\Phi(f') = \{c, a_2, a_3, \dots, a_m\} \text{ and } i(|K|, f', c) = i(|K|, f, a_1).$$

Proof. Let σ and σ' be two maximal simplices with $a_1 \in \sigma$ and $c \in \sigma'$. As $|K|$ is of type W , there exists a chain $\sigma_1 = \sigma, \sigma_2, \dots, \sigma_r = \sigma'$ of maximal simplices such that $\bar{\sigma}_i \cap \bar{\sigma}_{i+1}$ has dimension at least one for $i = 1, 2, \dots, r - 1$. By repeated use of Lemma 6 on p. 135 of [1] we can find a map g_1 which is homotopic to f and has a fixed point set $\Phi(g_1) = \{b, a_2, a_3, \dots, a_m\}$, where $b \in \sigma'$. We can also require that the line-segment $[b, c]$ contains no points of $\Phi(f)$, and that $g_1 = f$ in a neighbourhood of $\{a_2, a_3, \dots, a_m\}$. Next we obtain a map $g_2: |K| \rightarrow |K|$ by the same process (too complicated to describe briefly) which is used in the proof of Lemma 6 on p. 135 of [1] to obtain a map denoted there by f_2 , and therefore arrive at a map g_2 which is homotopic to g_1 , agrees with g_1 in a neighbourhood of $\{a_2, a_3, \dots, a_m\}$, has the same fixed points as g_1 , but has the property that $\bar{\kappa}(x) \cap \bar{\kappa}(g_2(x)) \neq \emptyset$ for all $x \in [b, c]$. Hence we can choose an $\eta > 0$ so that the following three conditions are satisfied:

- (i) $\bar{U}([b, c], \eta) \subset \text{st } \kappa(c)$,
- (ii) $\Phi(g_2) \cap \bar{U}([b, c], \eta) = b$,
- (iii) $\bar{\kappa}(x) \cap \bar{\kappa}(g_2(x)) \neq \emptyset$ for all $x \in \bar{U}([b, c], \eta)$.

We then modify g_2 to a map f' with fixed point set $\{c, a_2, a_3, \dots, a_m\}$

by putting $f'(x) = g_2(x)$ for $x \in |K| \setminus \bar{U}([b, c], \eta)$, and changing g_2 on $\bar{U}([b, c], \eta)$ by a method which is completely analogous to the one employed in the proof of Lemma 2.4 of [2]. The construction in [2] shows that f' is homotopic to g_2 and hence to f . As $f' = f$ on a neighbourhood of $\{a_2, a_3, \dots, a_m\}$, we have

$$\begin{aligned} \sum_{k=1}^m i(|K|, f, a_k) &= i(|K|, f', c) + \sum_{k=2}^m i(|K|, f', a_k) \\ &= i(|K|, f', c) + \sum_{k=2}^m i(|K|, f, a_k), \end{aligned}$$

and therefore $i(|K|, f', c) = i(|K|, f, a_1)$.

3. Mapping with prescribed fixed points and fixed point indices. We now proceed to construct deformations, and maps within an arbitrarily given homotopy class, with prescribed fixed points and indices. The method will be the same in both cases: we use Shi's [4] results to find a map with a minimum number of fixed points, and then use the splitting and the moving lemma of § 2 in order to obtain a map with the prescribed fixed points.—By $\chi(K)$ we understand the Euler characteristic of $|K|$.

THEOREM 1. *Let points c_k of a polyhedron $|K|$ of type W and integers i_k be given, where $k = 1, 2, \dots, m$. Then there exists a deformation which has the c_k as its fixed points and the i_k as its fixed point indices if and only if*

$$\sum \{i_k \mid 1 \leq k \leq m\} = \chi(K).$$

Proof. The necessity of the condition follows from the fact that the Lefschets number of a deformation equals the Euler characteristic of the polyhedron. (See [1], pp. 32, 52.) To prove its sufficiency, we construct a deformation $f_1: |K| \rightarrow |K|$ with exactly one fixed point b which is contained in a maximal simplex in the same way as in the proof of Theorem 1 on p. 143 of [1] (or use Theorem 3.1 of [2]). By subdividing K and using Lemma 2, if needed, we can change f_1 to a deformation f_2 with $\Phi(f_2) = a \in \sigma$, where σ is a maximal simplex with $\bar{\sigma} \cap \{c_1, c_2, \dots, c_m\} = \emptyset$. As f_2 is a deformation, we have $i(|K|, f_2, a) = \chi(K)$. Now we use Lemma 1 to construct a deformation f_3 with $\Phi(f_3) = \{a_1, a_2, \dots, a_m\} \subset \sigma$ and $i(|K|, f_3, a_k) = i_k$, and then make m -fold use of Lemma 2 to obtain the desired deformation f_4 with fixed points $\{c_1, c_2, \dots, c_m\}$ and indices $i(|K|, f_4, c_k) = i_k$.

In dealing with arbitrary maps rather than deformations we have to restrict the polyhedron $|K|$ further and assume that it is

of type S ([1], p. 139). This means that the dimension of $|K|$ is at least three, and that the boundary of the star of each vertex of $|K|$ is connected. A polyhedron of type S is always of type W , but the converse need not be true. (See [1], p. 143.)

The conditions on the indices in the generalization of Theorem 1 to arbitrary selfmaps will be more complicated. They require the concept of a *fixed point class* F_i of a selfmap f of $|K|$, and the *index* $i(F_i)$ of the fixed point class F_i , as defined in [1], pp. 86-87. The number of fixed point classes of a polyhedron is finite ([1], p. 86). A fixed point class F_i of f is called *essential* if $i(F_i) \neq 0$, and the *Nielsen number* $N = N(f)$ is the number of essential fixed point classes of f .

Both the number $N(f)$ and the indices $i(F_i)$ of the essential fixed point classes are homotopy invariant ([1], Chapter VI), and the conditions in Theorem 2 express precisely this fact. Theorem 2 coincides with Theorem 1 if $|K|$ is of type S and f is a deformation, as the Nielsen number of a deformation is 0 or 1.

THEOREM 2. *Let the selfmap f of the polyhedron $|K|$ of type S have the essential fixed point classes $F_l (l = 1, 2, \dots, N)$, with indices $i(F_l)$, and let points c_k of $|K|$ and integers $i_k (k = 1, 2, \dots, m)$ be given. Then there exists a selfmap which is homotopic to f , has the c_k as its fixed points and the i_k as its fixed point indices if and only if the c_k can be relabelled c_j , where $1 \leq j = j(k) \leq m$, such that either $N(f) = 0$ and $\sum \{i_j | 1 \leq j \leq m\} = 0$, or there exists a sequence of integers m_0, m_1, \dots, m_N with*

$$0 = m_0 < m_1 < \dots < m_N = m$$

for which

$$\sum \{i_j | m_{l-1} < j \leq m_l\} = i(F_l) \text{ for } l = 1, 2, \dots, N.$$

Proof. If $N(f) = 0$, then Theorem 1 on p. 140 of [1] shows that there exists a map $f_1: |K| \rightarrow |K|$ which is homotopic to f and fixed point free. In order to modify f_1 to a map with one fixed point, choose a point a in a maximal simplex σ of $|K|$, and arrange it so (by subdividing K , if necessary) that $\sigma \cap \{c_1, c_2, \dots, c_m\} = \emptyset$. Take $\eta > 0$ so that (i) $\bar{U}(a, \eta) \subset \sigma$, (ii) $f_1(\bar{U}(a, \eta)) \subset \text{st } \kappa(f_1(a))$, and (iii) $\bar{U}(a, \eta) \cap \text{st } \kappa(f_1(a)) = \emptyset$. (Again it may be necessary to subdivide K to make (iii) possible.) Let $\{\gamma(t) | 0 \leq t \leq 1\}$ be a path in $|K|$ from a to $f_1(a)$ for which $d(a, \gamma(t)) \neq t\eta/2$ for all $0 < t \leq 1$. Denote, for any $x \in \bar{U}(a, \eta) \setminus a$, by $x(\eta/2)$ and $x(\eta)$ the two unique points in which the ray from a to x intersects the boundaries of $\bar{U}(a, \eta)_2$, and $\bar{U}(a, \eta)$,

and define a map $f_2: |K| \rightarrow |K|$ by

$$f_2(x) = \begin{cases} f_1(x) & \text{if } x \in K \setminus \bar{U}(a, \eta) , \\ tf_1(x(\eta)) + (1 - t)f_1(a) & \text{if } x = tx(\eta) + (1 - t)x(\eta/2) , \\ \gamma(t) & \text{if } x = tx(\eta/2) + (1 - t)a , \end{cases}$$

where $0 \leq t \leq 1$.

It follows from the choice of η and $\gamma(t)$ that $\Phi(f_2) = a$. As f_2 is homotopic to f_1 , we have $N(f_2) = N(f_1) = 0$ and hence $i(|K|, f_2, a) = 0$. We now use Lemma 1 to find a map f_3 with $\Phi(f_3) = \{a_1, a_2, \dots, a_m\} \subset \sigma$ and $i(|K|, f_3, a_k) = i_k$, and then Lemma 2 to find f_4 with $\Phi(f_4) = \{c_1, c_2, \dots, c_m\}$ and $i(|K|, f_4, c_k) = i_k$ for $k = 1, 2, \dots, m$. Both f_3 and f_4 are homotopic to f , so that Theorem 2 holds if $N(f) = 0$.

If $N(f) > 0$, then Theorem 1 on p. 140 of [1], and its proof, show that there exists a map $f_1: |K| \rightarrow |K|$ in the homotopy class of f which has $N = N(f)$ fixed points b_1, b_2, \dots, b_N , where each b_i is contained in a maximal simplex. Again we can, if needed, subdivide K and use Lemma 2 to change f_1 to a map f_2 which is homotopic to f and has fixed points a_1, a_2, \dots, a_N , where each a_i is contained in a maximal simplex σ_i with $\bar{\sigma}_i \cap \{a_1, a_2, \dots, a_m, c_1, c_2, \dots, c_m\} = a_i$. We have $i(|K|, f_2, a_i) = i(|K|, f_1, b_i) = i(F_i)$. We now use Lemma 1 to split each a_i on σ_i into $m_i - m_{i-1}$ fixed points with indices i_j , for $m_{i-1} < j \leq m_i$, and then Lemma 2 to move these fixed points to the prescribed locations c_k .

4. Outlook. Theorems 1 and 2 deal only with mappings within a given homotopy class. The following, much more general question arises naturally:

If points c_k , where $k = 1, 2, \dots, m$, of a polyhedron $|K|$ and integers i_k are given, when does there exist a selfmap of $|K|$ which has the c_k as its fixed points and the i_k as its fixed point indices?

An answer to this question with present means will be difficult to obtain, as it seems necessary to know the number of essential fixed point classes, and their indices, which can occur for the different homotopy classes of selfmaps. We restrict our attention to one very special case of the problem, namely the one where $|K|$ is a sphere. In this case no restriction on the fixed points and indices is necessary.

THEOREM 3. *Let points c_k of a sphere S^2 and integers i_k be given, where $k = 1, 2, \dots, m$. Then there exists a selfmap of S^2 which has the c_k as its fixed points and the i_k as its fixed point indices.*

Proof. As S^2 is not of type S , we cannot make use of Theorem 2. Therefore we start by giving a direct construction of a selfmap which has precisely one fixed point with index $\sum \{i_k \mid 1 \leq k \leq m\}$.

We identify the unit sphere S^2 of R^3 with the suspension of the unit circle $S^1 = \{z \in C \mid |z| = 1\}$ in the complex plane, and therefore write the points of S^2 as $\langle z, s \rangle$, with $z \in S^1$ and $s \in [-1, 1]$. Let d be the integer determined by

$$1 + d = \sum_{k=1}^m i_k ,$$

and let

$$\lambda = \lambda(s) = \begin{cases} 1/2(1 + s) & \text{if } -1 \leq s \leq 0 , \\ 1/2(1 - s) & \text{if } 0 \leq s \leq 1 . \end{cases}$$

Then the map $f_0: S^2 \rightarrow S^2$ defined by

$$f_0(\langle z, s \rangle) = \langle z^d, s + \lambda \rangle$$

is of degree d and has the fixed points $u = \langle z, 1 \rangle$ and $v = \langle z, -1 \rangle$.

We now change f_0 to a map f_1 which is of the same degree, but has only one fixed point. For this purpose we select a point $a = \langle z_0, 0 \rangle$ so that $a \in \{c_1, c_2, \dots, c_m\}$ and $z_0^d \neq -z_0$. Denote by $\gamma = \gamma(s)$, where $-1 \leq s \leq 1$, the unique great arc through u, v , and a , and put for $\varepsilon > 0$

$$\bar{U}(\gamma, \varepsilon) = \{x \in S^2 \mid \widehat{\gamma(s)x} \leq \varepsilon \text{ for some } s \in [-1, 1]\} ,$$

where $\widehat{\gamma(s)x}$ is the distance from $\gamma(s)$ to x measured along a shortest arc on S^2 .

Select $\delta > 0$ so that $z^d \neq -z_0$ if $|z - z_0| \leq \delta$, and choose $\varepsilon = \varepsilon(\delta) > 0$ so that $\langle z, -1/3 \rangle \in \bar{U}(\gamma, \varepsilon)$ implies $|z - z_0| < \delta$, and also so that $\bar{U}(\gamma, \varepsilon)$ does not contain the antipode $b = \langle -z_0, 0 \rangle$ of a . If $f_0(x) = b$ for some $x = \langle z, s \rangle \in S^2$, then $z^d = -z_0$ and $s = -1/3$, therefore $f_0(x) \neq b$ for all $x \in \bar{U}(\gamma, \varepsilon)$.

Any $x \in \bar{U}(\gamma, \varepsilon) \setminus a$ determines a point $y = y(x)$ as the unique point on the boundary of $\bar{U}(\gamma, \varepsilon)$ for which the shortest arc from a to y contains x . Hence we can write any $x \in \bar{U}(\gamma, \varepsilon) \setminus a$ in the form $x = (1 - t)a + ty$, where $0 < t \leq 1$. Let $f_1: S^2 \rightarrow S^2$ be the map defined by

$$f_1(x) = \begin{cases} f_0(x) & \text{if } x \in S^2 \setminus \bar{U}(\gamma, \varepsilon) , \\ (1 - t)a + tf_0(y(x)) & \text{if } x \in \bar{U}(\gamma, \varepsilon) \setminus a , \\ a & \text{if } x = a . \end{cases}$$

As both f_0 and f_1 map $\bar{U}(\gamma, \varepsilon)$ into $S^2 \setminus b$, they are homotopic, and f_1 has degree d . The index $i(S^2, f_1, a) = i(S^2, f_1, S^2)$ equals the Lefschetz number of f_1 (see [1], p. 52), and hence is $1 + d = \sum_{k=1}^m i_k$.

It is always possible to choose a simplicial structure of S^2 for which a is contained in a maximal simplex σ with $\bar{\sigma} \cap \{c_1, c_2, \dots, c_m\} = \emptyset$. As S^2 is a polyhedron of type W , we can now use Lemma 1 to split a into fixed points a_1, a_2, \dots, a_m in σ with indices i_k , and then Lemma 2 to move each a_k to c_k in order to obtain the desired map.

REMARK. An n -sphere S^n with $n \geq 3$ is of type S , and hence an extension of Theorem 3 to such spheres is an almost immediate consequence of Theorem 2. It is easy to see that S^2 in Theorem 3 cannot, however, be replaced by S^1 .

REFERENCES

1. R. F. Brown, *The Lefschetz Fixed Point Theorem*, Glenview, Ill., 1971.
2. H. Schirmer, *Fixed point sets of polyhedra*, Pacific J. Math., **52** (1974), 221-226.
3. ———, *Attractive and repulsive fixed points on dendrites*, to appear.
4. Shi Gen-Hua, *On the least number of fixed points and Nielsen numbers*, Chinese Math., **8** (1966), 234-243.

Received September 20, 1974 and in revised form January 22, 1975. This research was partially supported by the National Research Council of Canada (Grant A-7579).

CARLETON UNIVERSITY, OTTAWA, CANADA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Pacific Journal of Mathematics

Vol. 63, No. 2

April, 1976

Joseph Anthony Ball and Arthur R. Lubin, <i>On a class of contractive perturbations of restricted shifts</i>	309
Joseph Becker and William C. Brown, <i>On extending higher derivations generated by cup products to the integral closure</i>	325
Andreas Blass, <i>Exact functors and measurable cardinals</i>	335
Joseph Eugene Collison, <i>A variance property for arithmetic functions</i>	347
Craig McCormack Cordes, <i>Quadratic forms over nonformally real fields with a finite number of quaternion algebras</i>	357
Freddy Delbaen, <i>Weakly compact sets in H^1</i>	367
G. D. Dikshit, <i>Absolute Nörlund summability factors for Fourier series</i>	371
Edward Richard Fadell, <i>Nielsen numbers as a homotopy type invariant</i>	381
Josip Globevnik, <i>Analytic extensions of vector-valued functions</i>	389
Robert Gold, <i>Genera in normal extensions</i>	397
Solomon Wolf Golomb, <i>Formulas for the next prime</i>	401
Robert L. Griess, Jr., <i>The splitting of extensions of $SL(3, 3)$ by the vector space F_3^3</i>	405
Thomas Alan Keagy, <i>Matrix transformations and absolute summability</i>	411
Kazuo Kishi, <i>Analytic maps of the open unit disk onto a Gleason part</i>	417
Kwangil Koh, Jiang Luh and Mohan S. Putcha, <i>On the associativity and commutativity of algebras over commutative rings</i>	423
James C. Lillo, <i>Asymptotic behavior of solutions of retarded differential difference equations</i>	431
John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues</i>	445
Anna Maria Mantero, <i>Sets of uniqueness and multiplicity for L^p</i>	467
J. F. McClendon, <i>Embedding metric families</i>	481
L. Robbiano and Giuseppe Valla, <i>Primary powers of a prime ideal</i>	491
Wolfgang Ruess, <i>Generalized inductive limit topologies and barrelledness properties</i>	499
Judith D. Sally, <i>Bounds for numbers of generators of Cohen-Macaulay ideals</i>	517
Helga Schirmer, <i>Mappings of polyhedra with prescribed fixed points and fixed point indices</i>	521
Cho Wei Sit, <i>Quotients of complete multipartite graphs</i>	531
S. Sznajder and Zbigniew Zielezny, <i>Solvability of convolution equations in \mathfrak{K}'_p, $p > 1$</i>	539
Mitchell Herbert Taibleson, <i>The existence of natural field structures for finite dimensional vector spaces over local fields</i>	545
William Yslas Vélez, <i>A characterization of completely regular fields</i>	553
P. S. Venkatesan, <i>On right unipotent semigroups</i>	555
Kenneth S. Williams, <i>A rational octic reciprocity law</i>	563
Robert Ross Wilson, <i>Lattice orderings on the real field</i>	571
Harvey Eli Wolff, <i>V-localizations and V-monads. II</i>	579