

Pacific Journal of Mathematics

ON RIGHT UNIPOTENT SEMIGROUPS

P. S. VENKATESAN

ON RIGHT UNIPOTENT SEMIGROUPS

P. S. VENKATESAN

We investigate the implications of certain conditions on right unipotent semigroups. We describe the greatest idempotent-separating congruence β on a right unipotent semigroup S . Necessary and sufficient conditions for (i) S to be a union of groups, (ii) S to be an inverse semigroup, (iii) the idempotents of S to be in the centre of S and (iv) the quotient semigroup S/β to be isomorphic with the subsemigroup of idempotents of S are also obtained.

It is known that any regular semigroup has the greatest idempotent-separating congruence [5], [6]. Such a congruence on an inverse semigroup was obtained by Howie [4]. For the general terminology and notation the reader is referred to [1], [2].

The author wishes to express his thanks to the referee for his valuable comments.

1. **Preliminary matters.** An orthodox semigroup S is a regular semigroup in which the idempotents form a subsemigroup. An inverse of an idempotent of S is an idempotent, and if a', b' are inverses of the elements a, b in S then $b'a'$ is an inverse of ab [7].

A semigroup S is called a *right (left) unipotent semigroup* if every principal right (left) ideal of S has a unique idempotent generator. Such semigroups are called left (right) inverse by the author [9], [10]. Lemma 1 below is a part of the left-right dual of Theorem 1 in [10].

LEMMA 1. *Let S be a regular semigroup. Then the following statements are equivalent.*

- (A) $fef = fe$ for any two idempotents e, f in S .
- (B) If a' and a'' are inverses of the element a in S then $aa' = aa''$.
- (C) S is a right unipotent semigroup.

LEMMA 2. *Let S be a right unipotent semigroup and e be an idempotent of S . Let $x \in S$ and x', x'' be inverses of x . Then xex' is an idempotent and $xex' = xex''$.*

Proof. By Lemma 1 we have $xe = xx'xe = x(x'exex') = xex'x$. So xex' is an idempotent. Also $xex' = (xex')xx' = (xex')xx'' = (xex'x)x'' = xex''$, using Lemma 1.

2. The statements (Px) , (Qx) and (Rx) . Let S be a right unipotent semigroup and $x \in S$. Throught $E = E(S)$ denotes the subsemigroup of idempotents of S and $V(x)$ denotes the set of inverses of the element x . The symbols (Px) , (Qx) and (Rx) stand for the statements indicated below.

(Px) $exe = ex$ and $ex'e = ex'$ for all $e \in E$ and for at least one $x' \in V(x)$.

(Qx) $xex' = xx'e$ for all $e \in E$ and $x' \in V(x)$.

(Rx) $xex' = exx'$ for all $e \in E$ and $x' \in V(x)$.

REMARK. Let S be a left unipotent semigroup. Then the left-right dual of (Px) , (Qx) and (Rx) are obtained by replacing respectively the equations in them by $xe = xe$ and $ex'e = x'e$, $x'ex = ex'x$ and $x'ex = x'xe$.

THEOREM 1. *Let S be a right unipotent semigroup and $E = E(S)$. Then*

(1) $(Rx) \Rightarrow (Qx) \Rightarrow (Px)$ for any $x \in S$.

(2) E is contained in the center of S if and only if (Rx) is satisfied for all $x \in S$.

Proof. (1) Let $x \in S$ and $x' \in V(x)$.

Assume (Rx) . Then for any $e \in E$ we have $xex' = exx'$ and hence $ex = (exx')x = xex'x = x(x'exx') = x(x'xe) = xe$ by Lemma 1. So $xx'e = xx'(ex)x' = xx'(xe)x' = xex'$, giving (Qx) .

Assume (Qx) . Then $xex' = xx'e$ for any $e \in E$. Therefore, by Lemma 1, we get $exe = ex(x'xe) = ex(x'exx') = e(xex')x = e(xx'e)x = exx'x = ex$ and $ex'e = ex'(xx'e) = ex'(xex') = ex'xx' = ex'$, giving (Px) .

(2) The only if part is trivial. The if part follows since, for any $x \in S$ and $e \in E$, as shown above, (Rx) implies $ex = xe$.

Let S be a right unipotent semigroup. Then the statements (1) S is union of groups, (2) each \mathcal{L} -class of S is a left group and (3) each \mathcal{R} -class of S is a group are equivalent [8]. An alternate characterization for S to be a union of groups is obtained in the following

THEOREM 2. *Let S be a right unipotent semigroup and $E = E(S)$. Then S is a union of groups if and only if (Px) is satisfied for all x in S .*

Proof. Let S be a union of groups. Let $x \in S$ and $e \in E$. Let x^{-1} be the inverse of x in the group H_x . Then $x^{-1}x = xx^{-1}$. Let a and b respectively be the group inverses of ex and ex^{-1} . As $x^{-1}e$ is

an inverse of ex , and xe is an inverse of ex^{-1} , by Lemma 1 we have $exa = exx^{-1}e$ and $ex^{-1}b = ex^{-1}xe$. But $exx^{-1}e = exx^{-1}$ and $ex^{-1}xe = exx^{-1}e = exx^{-1}$ by Lemma 1. So both ex and ex^{-1} and hence their product $exex^{-1}$ belong to the group with identity element exx^{-1} . As $exex^{-1}$ is an idempotent we conclude that $exex^{-1} = exx^{-1}$. Therefore $axe = ex(x^{-1}xe) = ex(x^{-1}xex^{-1}x) = (exex^{-1})x = exx^{-1}x = ex$ by Lemma 1. Further since ex^{-1} belongs to the group with identity element exx^{-1} , we have $ex^{-1} = ex^{-1}(exx^{-1}) = ex^{-1}(xx^{-1}exx^{-1}) = ex^{-1}(xx^{-1}e) = ex^{-1}e$, by Lemma 1. So we get (Px) .

Conversely let (Px) be satisfied for all $x \in S$. (This part of the proof holds for any regular semigroup S). Let $x \in S$ and $x' \in V(x)$. Taking $e = xx'$ in $ex = exe$ we have $x = x^2x' \in x^2S$. So S is a right regular semigroup and hence a union of group [1], [3].

Let S be a right unipotent semigroup. Then S is an inverse semigroup if and only if S satisfies the left-right dual of any of the statements of Lemma 1. We now obtain a necessary and sufficient condition in terms of (Px) and (Rx) for S to be an inverse semigroup.

THEOREM 3. *Let S be a right unipotent semigroup and $E = E(S)$. Then S is an inverse semigroup if and only if (Px) implies (Rx) for all x in S .*

Proof. Let S be an inverse semigroup. Let $x \in S$. Assume (Px) . Then for any $e \in E$ we have $axe = ex$ and $ex^{-1}e = ex^{-1}$. As the idempotents in S commute we get $exx^{-1} = (axe)x^{-1} = e(xex^{-1}) = (xx^{-1})e = x(ex^{-1}e) = xex^{-1}$, giving (Rx) .

Conversely let (Px) imply (Rx) for all $x \in S$. Let $g, h \in E$. Then for any $e \in E$, by Lemma 1, we have $e(gh)e = egh$. As $gh \in V(gh)$, by hypothesis we conclude $gh egh = eghgh$. So, by Lemma 1, we get $ghe = egh$. Taking $e = h$, by Lemma 1, we have $gh = hg$. Thus S is an inverse semigroup.

COROLLARY. *Let S be a right unipotent semigroup and $E = E(S)$. Then S is an inverse semigroup if and only if (Px) , (Qx) and (Rx) are equivalent for all x in S .*

REMARK. The left-right dual of Theorems 1, 2 and 3 hold for a left unipotent semigroup.

3. The congruences α and β . In this section we construct the greatest idempotent-separating congruence on a right (left) unipotent semigroup.

Theorems 4 and 7 below generalize known results for inverse semigroups [4]. In [6] Munn relates the greatest idempotent-

separating congruence on an inverse semigroup to a certain full inverse semigroup. We need the following

LEMMA 3. *Let S be an orthodox semigroup and σ be an idempotent-separating congruence on S . If $(x, y) \in \sigma$ then there exist $u \in V(x)$ and $v \in V(y)$ such that $(u, v) \in \sigma$.*

Proof. Let $(x, y) \in \sigma$, $x' \in V(x)$ and $y' \in V(y)$. Since σ is a congruence we get $(x'x, x'y) \in \sigma$ and hence $(x'xy'y, x'y) \in \sigma$. By transitivity of σ we conclude $(x'xy'y, x'x) \in \sigma$. This, since σ is idempotent-separating, implies $x'xy'y = x'x$. So $xy'y = x$. Similarly we get $xx'yy' = yy'$ and $xx'y = y$.

Set $u = y'yx'$ and $v = y'xx'$. Then $u \in V(x)$ and $v \in V(y)$. Now from $(x, y) \in \sigma$ we have $(y'xx', y'yx') \in \sigma$, that is $(v, u) \in \sigma$ and thus $(u, v) \in \sigma$. Hence the lemma.

Let S be a regular semigroup and E be the set of idempotents of S . Define the binary relations α and β on S thus:

$$\alpha = \{(x, y) \in S \times S : x'ex = y'ey \text{ for all } e \in E, x' \in V(x) \text{ and } y' \in V(y)\}.$$

$$\beta = \{(x, y) \in S \times S : xex' = yey' \text{ for all } e \in E, x' \in V(x) \text{ and } y' \in V(y)\}.$$

THEOREM 4. *Let S be a right (left) unipotent semigroup and $E = E(S)$. Then $\beta(\alpha)$ is an idempotent-separating congruence on S . Further, if σ is any idempotent-separating congruence on S then $\sigma \subseteq \beta(\sigma \subseteq \alpha)$.*

Proof. We prove the theorem for the right unipotent semigroup S . Clearly β is an equivalence relation on S . Let $(x, y) \in \beta$. Let $c \in S$ and $c' \in V(c)$ and $x' \in V(x)$. Then $x'c' \in V(cx)$ and $y'c' \in V(cy)$. As $c(xex')c' = c(yey')c'$, by Lemma 2, we get $(cx, cy) \in \beta$ and β is a left congruence. Further, since cec' is an idempotent for any $e \in E$, $c'x' \in V(xc)$ and $c'y' \in V(yc)$ we have $x(cec')x' = y(cec')y'$. So by Lemma 2, $(xc, yc) \in \beta$. Therefore β is a right congruence and hence a congruence relation on S .

Now let $g, h \in E$ and suppose that $(g, h) \in \beta$. Then by Lemma 2, for any $e \in E$ we have $geg = heh$. Taking $e = g$ and $e = h$ in turn we obtain $g = hgh = hg$ and $h = ghg = gh$ using Lemma 1. Therefore $g = h(gh) = hh = h$ proving that β is idempotent-separating.

Now let σ be any idempotent-separating congruence on S . Let $(x, y) \in \sigma$. Then by Lemma 3 there exist $x' \in V(x)$ and $y' \in V(y)$ such that $(x', y') \in \sigma$. As σ is a congruence, for any $e \in E$ we have $(xe, ye) \in \sigma$ and hence $(xex', yey') \in \sigma$. But xex' and yey' are idempotents and σ is idempotent-separating. Therefore $xex' = yey'$. This, by

Lemma 2, implies $(x, y) \in \beta$ and thus $\sigma \subseteq \beta$. Hence the theorem.

COROLLARY [4]. *Let S be an inverse semigroup. Then $\alpha(=\beta)$ is the greatest idempotent-separating congruence on S .*

THEOREM 5. *Let S be a right unipotent semigroup and $E = E(S)$. For each $x \in S$ let $\theta_x: E \rightarrow E$ be the mapping defined by $\theta_x(e) = xex'$ where $x' \in V(x)$. Then*

- (1) θ_x is an endomorphism, and
- (2) the following statements are equivalent.
 - (A) θ_x is an idempotent.
 - (B) The \mathcal{H} -class H_x is a subgroup of S and $xx^{-1}e = xex^{-1} = x^{-1}ex$ for all $e \in E$ where x^{-1} is the group inverse of x in H_x .
 - (C) $\theta_x = \theta_g$ where $g = xx'$.

Proof. Let $x \in S$ and $x' \in V(x)$.

(1) For any $e, f \in E$, by Lemma 1, we have $(xex')(xfx') = (xex'x)fx' = xefx'$, proving (1).

(2) Assume (A). Then $xex' = xxex'x'$ for all $e \in E$. Taking $x'x$ for e we have $xx' = xxx'x'$ and therefore $x'x = x'(xx')x = x'xxx'x'x = x'xxx'$ using Lemma 1. So $x = xx'x = x^2x'$ and $x\mathcal{R}x^2$.

Now taking $x'x'xx$ for e in $xex' = xxex'x'$, and using $x = x^2x'$ and Lemma 1, we get $xx'x'x = xx'$ and hence $x'x'x = x'$. Therefore $x = xx'x = xx'x'x^2$ and $x\mathcal{L}x^2$. Thus $x\mathcal{H}x^2$ and H_x is a subgroup of S .

By hypothesis and Lemma 2, for all $e \in E$ we have $xex^{-1} = x^2ex^{-2}$ and therefore $x^{-1}xe = x^{-1}xex^{-1}x = x^{-1}(x^2ex^{-2})x = xex^{-1}$ since $x^{-1}x = xx^{-1}$. Again taking $x^{-2}ex^2$ for e in $xex^{-1} = x^2ex^{-2}$ we get $x^{-1}ex = xx^{-1}e$ using Lemma 1. So we get (B).

Assume (B). Then by Lemma 1, for all $e \in E$ we have $xex^{-1} = xx^{-1}e = xx^{-1}e$, giving (C). Clearly (C) implies (A).

THEOREM 6. *Let S be a right unipotent semigroup and $E = E(S)$. Then*

- (1) $T = \{\theta_x: x \in S\}$ is a right unipotent semigroup.
- (2) The mapping $\theta: S \rightarrow T$ defined by $\theta(x) = \theta_x$ is an onto homomorphism and $\theta \cdot \theta^{-1} = \beta$.
- (3) Set $\gamma = \theta|_E$ (θ restricted to E). Then γ is an isomorphism of E upon $\theta(E)$.

Proof. As $\theta_x\theta_y = \theta_{xy}$ it follows that T is a regular semigroup. We now show directly that T is right unipotent. Let θ_x and θ_y be idempotents of T . Then, for all $e \in E$, using (B) of Theorem 5 repeatedly we have $x(yxex^{-1}y^{-1})x^{-1} = xx^{-1}y(xex^{-1})y^{-1} = xx^{-1}yy^{-1}(xex^{-1}) = xx^{-1}(yy^{-1})xex^{-1} = xx^{-1}x(yy^{-1})ex^{-1} = x(yy^{-1}e)x^{-1} = xyey^{-1}x^{-1}$, and hence

$\theta_{xyx} = \theta_{xy}$ by Lemma 2. So, by Lemma 1, T is a right unipotent semigroup, proving (1).

(2) follows directly. As for (3) we need only to show that γ is one-to-one. Let $\gamma(g) = \gamma(h)$ for $g, h \in E$. Then by Lemma 2, we have $(g, h) \in \beta$. This, by Theorem 4, implies $g = h$ and so γ is an isomorphism.

We now consider the quotient semigroup S/β . The following theorem gives a necessary and sufficient condition for S/β to be an idempotent semigroup.

THEOREM 7. *Let S be a right unipotent semigroup and $E = E(S)$. Then the quotient semigroup S/β is isomorphic with E if and only if the statement (Qx) is satisfied for all x in S . (The left-right dual holds for a left unipotent semigroup).*

Proof. Let S/β be isomorphic with E . As S/β is a homomorphic image of S , each idempotent of S/β is the image of an idempotent of S [10]. So each β -class of S contains at least one and hence exactly one idempotent of S . Let $x \in S$. Then there exists $h \in E$ such that $(x, h) \in \beta$. So for any $e \in E$ and $x' \in V(x)$ we have $xx' = heh$. In particular taking $e = x'x$ and $e = h$ in turn we get $xx' = hx'xh$ and $xx' = h$. The first equation gives $xx'h = xx'$ and the second $xx'h = h$. So $xx' = h$. Hence for any $e \in E$, by Lemma 1, we have $xxe' = heh = he = xx'e$ giving (Qx) .

Conversely let (Qx) be satisfied for all $x \in S$. Let $x \in S$. Then for any $e \in E$ and $x' \in V(x)$, by Lemma 1, we have $xxe' = xx'e = xx'exe'$. Therefore $(x, xx') \in \beta$ and hence each β -class contains a unique idempotent. Let β^* be the natural homomorphism of S upon S/β . Then the mapping β^* restricted to E is an isomorphism of E upon S/β . This completes the proof of the theorem.

REMARK. One may appeal to Theorems 5 and 6 to prove Theorem 7. Clearly, for any $x \in S$, the statements (Qx) , and (C) of Theorem 5 are equivalent. Therefore (Qx) is satisfied for all $x \in S$ if and only if $T = \theta(E)$ and hence if and only if S/β is isomorphic with E .

From Theorems 1, 2, 7 and the corollary of Theorem 3 we have the following.

COROLLARY [4]. *Let S be an inverse semigroup. Then the following statements are equivalent.*

- (A) *The quotient semigroup S/β is isomorphic with E .*
- (B) *S is a union groups.*
- (C) *The idempotents of S are contained in the centre of S .*

REFERENCES

1. A. H. Clifford and G. B. Preston, *The Algebraic Theory of Semigroups*, Vol. 1, Math. Surveys, Amer. Math. Soc., **7** (1961).
2. ———, Vol. 2, Math. Surveys, Amer. Math. Soc., **7** (1967).
3. R. Croisot, *Demi-groupes inversifs et demigroupes réunions de demi-groupes simples*, Ann. Sci. école normale supérieure, **70** (1953), 361-379.
4. J. M. Howie, *The maximum idempotent-separating congruence on an inverse semigroup*, Proc. Edinburg Math. Soc., (2), **14** (1964/65), 71-79.
5. G. Lallement, *Demi-groupes réguliers*, Ann. Mat. Pura e Appl., **77** (1967), 47-130.
6. W. D. Munn, *Fundamental inverse semigroups*, Quart. J. Math. Oxford, (2) **21** (1970), 157-170.
7. N. R. Reilly and H. E. Scheiblich, *Congruences on regular semigroups*, Pacific J. Math., **23** (1967), 349-360.
8. P. S. Venkatesan, *On decomposition of semigroups with zero*, Math. Zeitschr., **92** (1966), 164-174.
9. ———, *Bisimple left inverse semigroups*, Semigroup Forum, **4** (1972), 34-45.
10. ———, *Right (left) inverse semigroups*, J. Algebra, (2), **31** (1974), 209-217.

Received May 3, 1973 and in revised form November 20, 1975.

UNIVERSITY OF IBADAN, IBADAN, NIGERIA.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Joseph Anthony Ball and Arthur R. Lubin, <i>On a class of contractive perturbations of restricted shifts</i>	309
Joseph Becker and William C. Brown, <i>On extending higher derivations generated by cup products to the integral closure</i>	325
Andreas Blass, <i>Exact functors and measurable cardinals</i>	335
Joseph Eugene Collison, <i>A variance property for arithmetic functions</i>	347
Craig McCormack Cordes, <i>Quadratic forms over nonformally real fields with a finite number of quaternion algebras</i>	357
Freddy Delbaen, <i>Weakly compact sets in H^1</i>	367
G. D. Dikshit, <i>Absolute Nörlund summability factors for Fourier series</i>	371
Edward Richard Fadell, <i>Nielsen numbers as a homotopy type invariant</i>	381
Josip Globevnik, <i>Analytic extensions of vector-valued functions</i>	389
Robert Gold, <i>Genera in normal extensions</i>	397
Solomon Wolf Golomb, <i>Formulas for the next prime</i>	401
Robert L. Griess, Jr., <i>The splitting of extensions of $SL(3, 3)$ by the vector space F_3^3</i>	405
Thomas Alan Keagy, <i>Matrix transformations and absolute summability</i>	411
Kazuo Kishi, <i>Analytic maps of the open unit disk onto a Gleason part</i>	417
Kwangil Koh, Jiang Luh and Mohan S. Putcha, <i>On the associativity and commutativity of algebras over commutative rings</i>	423
James C. Lillo, <i>Asymptotic behavior of solutions of retarded differential difference equations</i>	431
John Alan MacBain, <i>Local and global bifurcation from normal eigenvalues</i>	445
Anna Maria Mantero, <i>Sets of uniqueness and multiplicity for L^p</i>	467
J. F. McClendon, <i>Embedding metric families</i>	481
L. Robbiano and Giuseppe Valla, <i>Primary powers of a prime ideal</i>	491
Wolfgang Ruess, <i>Generalized inductive limit topologies and barrelledness properties</i>	499
Judith D. Sally, <i>Bounds for numbers of generators of Cohen-Macaulay ideals</i>	517
Helga Schirmer, <i>Mappings of polyhedra with prescribed fixed points and fixed point indices</i>	521
Cho Wei Sit, <i>Quotients of complete multipartite graphs</i>	531
S. Sznajder and Zbigniew Zielezny, <i>Solvability of convolution equations in \mathcal{H}'_p, $p > 1$</i>	539
Mitchell Herbert Taibleson, <i>The existence of natural field structures for finite dimensional vector spaces over local fields</i>	545
William Yslas Vélez, <i>A characterization of completely regular fields</i>	553
P. S. Venkatesan, <i>On right unipotent semigroups</i>	555
Kenneth S. Williams, <i>A rational octic reciprocity law</i>	563
Robert Ross Wilson, <i>Lattice orderings on the real field</i>	571
Harvey Eli Wolff, <i>V-localizations and V-monads. II</i>	579