

Pacific Journal of Mathematics

NON-HAUSDORFF MULTIFUNCTION GENERALIZATION OF THE KELLEY-MORSE ASCOLI THEOREM

GEOFFREY FOX AND PEDRO MORALES

NON-HAUSDORFF MULTIFUNCTION GENERALIZATION OF THE KELLEY-MORSE ASCOLI THEOREM

GEOFFREY FOX AND PEDRO MORALES

The paper generalizes the Kelley-Morse theorem to continuous point-compact multifunction context. The generalization, which is non-Hausdorff, contains the Ascoli theorem for continuous functions on a k_3 -space by the authors and the known multifunction Ascoli theorems of Mancuso and of Smithson.

1. Introduction. The Kelley-Morse theorem [3, p. 236] is central among the topological Ascoli theorems for continuous functions on a k -space. It generalizes to the k_3 -space theorem of [1], which contains all known Ascoli theorems for k -spaces or k_3 -spaces.

Obviously a multifunction generalization depends on a multifunction extension of "even continuity". One such extension is that of Lin and Rose [5], but this was not applied in Kelley-Morse context. Another which was so applied [7, p. 24] is two-fold and leads to a two-fold multifunction Kelley-Morse theorem which, however, does not contain the Mancuso theorem [6, p. 470], nor the Smithson theorem [9, p. 259]. This paper gives a natural multifunction extension of the definition and leads to a multifunction theorem containing all the above-mentioned theorems.

2. Tychonoff sets. Let X and Y be nonempty sets. A *multifunction* is a point to set correspondence $f: X \rightarrow Y$ such that, for all $x \in X$, fx is a nonempty subset of Y . For $A \subseteq X$, $B \subseteq Y$ it is customary to write $f(A) = \bigcup_{x \in A} fx$, $f^-(B) = \{x: x \in X \text{ and } fx \cap B \neq \emptyset\}$ and $f^+(B) = \{x: x \in X \text{ and } fx \subseteq B\}$. If Y is a topological space, a multifunction $f: X \rightarrow Y$ is *point-compact* if fx is compact for all $x \in X$.

Let $\{Y_x\}_{x \in X}$ be a family of nonempty sets. The *m-product* $P\{Y_x: x \in X\}$ of the Y_x is the set of all multifunctions $f: X \rightarrow \bigcup_{x \in X} Y_x$ such that $fx \subseteq Y_x$ for all $x \in X$. In the case $Y_x = Y$ for all $x \in X$, the *m-product* of the Y_x , denoted Y^{mX} , is the set of all multifunctions on X to Y . In particular, if Y is a topological space, the symbol $(Y^{mX})_0$ will denote the set of all point-compact members of Y^{mX} . For $x \in X$, the *x-projection* $\text{pr}_x: P\{Y_x: x \in X\} \rightarrow Y_x$ is the multifunction defined by $\text{pr}_x f = fx$. If the Y_x are topological spaces, the *pointwise topology* τ_p on $P\{Y_x: x \in X\}$ is defined to be the topology having as open subbase the sets of the forms $\text{pr}_x^-(U_x)$, $\text{pr}_x^+(U_x)$, where U_x is open in Y_x , $x \in X$.

For $F \subseteq Y^{mX}$, $x \in X$, we write $F[x] = \bigcup_{f \in F} fx$. Let Y be a topological space. A subset F of Y^{mX} is *pointwise bounded* if $F[x]$ has compact

closure in Y for all $x \in X$. A subset T of Y^{m^x} is *Tychonoff* if, for every pointwise bounded subset F of T , $T \cap P\{\overline{F[x]}: x \in X\}$ is τ_p -compact.

The following sets are Tychonoff:

- (1) Y^X , by the classical Tychonoff theorem.
- (2) Y^{m^x} , by the theorem of Lin [4, p. 400].
- (3) The set of all point-closed members of Y^{m^x} , by Corollary 7.5 of [7, p. 17].
- (4) $(Y^{m^x})_0$, by Corollary 7.6 of [7, p. 17].

LEMMA 2.1. *If F is a pointwise bounded subset of a Tychonoff set T , then the τ_p -closure of F in T is compact.*

Proof. Let \overline{F} denote the τ_p -closure of F in T . Since $T \cap P\{\overline{F[x]}: x \in X\}$ is a τ_p -compact subset of T , it suffices to show that $\overline{F} \subseteq P\{\overline{F[x]}: x \in X\}$. But this follows from Lemma 7.7 of [7, p. 17].

3. Even continuity. Let X and Y be topological spaces. A multifunction $f: X \rightarrow Y$ is *lower semi-continuous* (*upper semi-continuous*) if $f^-(U)$ ($f^+(U)$) is open in X whenever U is open in Y . If f is both lower semi-continuous and upper semi-continuous it is called *continuous*. Henceforth, the set of all continuous multifunctions on X to Y will be denoted $\mathcal{C}(X, Y)$. The multifunction $(f, x) \rightarrow fx$ on $Y^{m^x} \times X$ to Y , or any restriction, will be denoted by the symbol ω . Let $F \subseteq Y^{m^x}$. A topology τ on F is said to be *jointly continuous* if $\omega: (F, \tau) \times X \rightarrow Y$ is continuous.

A subset F of Y^{m^x} is *evenly continuous* if, whenever $x \in X$, K is a compact subset of Y and V is a neighborhood of K , there exist neighborhoods U, W of x, K , respectively, such that

- (a) $f \in F$ and $fx \cap W \neq \emptyset$ imply $U \subseteq f^-(V)$, and
- (b) $f \in F$ and $fx \subseteq W$ imply $U \subseteq f^+(V)$.

This extends the original Kelley-Morse definition [3, p. 235] by the substitution of compact subsets of Y for points of Y . It is easily verified that every member of an evenly continuous subset of Y^{m^x} is lower semi-continuous. Moreover, every member of an evenly continuous subset of $(Y^{m^x})_0$ is also upper semi-continuous, hence continuous.

LEMMA 3.1. *Let Y be a regular space. If F is an evenly continuous subset of Y^{m^x} , then the τ_p -closure of F in Y^{m^x} is evenly continuous.*

Proof. Let \overline{F} denote the τ_p -closure of F in Y^{m^x} . Let $x \in X$, let K be a compact subset of Y and let V be a closed neighborhood of K . There exist open neighborhoods U, W of x, K , respectively, such that, for all $f \in F$, $fx \cap W \neq \emptyset$ implies $U \subseteq f^-(V)$ and $fx \subseteq W$ implies $U \subseteq f^+(V)$. Let $g \in \overline{F}$ be such that $gx \cap W \neq \emptyset$. Let $\{g_\alpha\}$ be a net in F

which is τ_p -convergent to g . Since $\{h: h \in Y^{m \times}$ and $hx \cap W \neq \emptyset\}$ is a τ_p -neighborhood of g , $g_\alpha x \cap W \neq \emptyset$ eventually, so $U \subseteq g_\alpha^-(V)$ eventually. Suppose that $U \not\subseteq g^-(V)$. Then, for some $u \in U$, $gu \subseteq Y - V$, so $g_\alpha u \subseteq Y - V$ eventually, which is a contradiction.

Now let $g \in \bar{F}$ be such that $gx \subseteq W$. Let $\{g_\alpha\}$ be a net in F which is τ_p -convergent to g . Since $\{h: h \in Y^{m \times}$ and $hx \subseteq W\}$ is a τ_p -neighborhood of g , $g_\alpha x \subseteq W$ eventually, so $U \subseteq g_\alpha^+(V)$ eventually. Suppose that $U \not\subseteq g^+(V)$. Then, for some $u \in U$, $gu \cap (Y - V) \neq \emptyset$, so $g_\alpha u \cap (Y - V) \neq \emptyset$ eventually, which is a contradiction.

LEMMA 3.2. *If F is an evenly continuous subset of $(Y^{m \times})_0$, then τ_p on F is jointly continuous.*

Proof. Let $\omega: (F, \tau_p) \times X \rightarrow Y$. Suppose that $(f, x) \in \omega^-(G)$, where G is open in Y . Choose $y \in fx \cap G$. There are neighborhoods U, W of x, y , respectively, such that $g \in F$ and $gx \cap W \neq \emptyset$ imply $U \subseteq g^-(G)$. Then $\{h: h \in F$ and $hx \cap W \neq \emptyset\} \times U$ is a neighborhood of (f, x) which is contained in $\omega^-(G)$.

Now suppose that $(f, x) \in \omega^+(G)$, where G is open in Y . There are neighborhoods U, W of x, fx , respectively, such that $g \in F$ and $gx \subseteq W$ imply $U \subseteq g^+(G)$. Then $\{h: h \in F$ and $hx \subseteq W\} \times U$ is a neighborhood of (f, x) which is contained in $\omega^+(G)$.

The following lemma generalizes an implicit lemma of Noble [8], stated explicitly as Lemma 1.4 in [7, p. 7]:

LEMMA 3.3. *Let $f \in \mathcal{C}(X \times Y, Z)$. If X is compact and Z is regular, then the set $F = \{f(x, \cdot): x \in X\}$ is evenly continuous.*

Proof. Let $y \in Y$, let K be a compact subset of Z and let V be an open neighborhood of K . Let W be a closed neighborhood of K which is contained in V . We construct a neighborhood U of y as follows: Since $f(\cdot, y)$ is continuous, $K_1 = f(\cdot, y)^-(W)$ and $K_2 = f(\cdot, y)^+(W)$ are closed in X , therefore compact. Thus the second projections $\text{pr}_2: K_1 \times Y \rightarrow Y, \text{pr}_2: K_2 \times Y \rightarrow Y$ are closed, so that

$$U_1 = Y - \text{pr}_2[(K_1 \times Y) - f^-(V)], \quad U_2 = Y - \text{pr}_2[(K_2 \times Y) - f^+(V)]$$

are open in Y . Because $K_1 \subseteq f(\cdot, y)^-(V), K_2 \subseteq f(\cdot, y)^+(V)$, we have $K_1 \times \{y\} \subseteq f^-(V), K_2 \times \{y\} \subseteq f^+(V)$. Hence $y \notin \text{pr}_2[(K_1 \times Y) - f^-(V)], y \notin \text{pr}_2[(K_2 \times Y) - f^+(V)]$, that is, $y \in U_1 \cap U_2 = U$.

We show that the neighborhoods U, W of y, K , respectively, satisfy the required implications: Let $g \in F$ be such that $gy \cap W \neq \emptyset$, so that $g = f(x, \cdot)$ for some $x \in K_1$. Let $u \in U$, so that $u \notin \text{pr}_2[(K_1 \times Y) - f^-(V)]$.

Then $(x, y) \in f^-(V)$, that is, $gu \cap V \neq \emptyset$. Now let $g \in F$ be such that $gy \subseteq W$, so that $g = f(x, \cdot)$ for some $x \in X_2$. Let $u \in U$, so that $u \notin \text{pr}_2[(K_2 \times Y) - f^+(V)]$. Then $(x, u) \in f^+(V)$, that is, $gu \subseteq V$.

Let X and Y be topological spaces. The *compact open topology* τ_c on Y^{mX} is defined to be the topology having as open subbase the sets of the forms $\{f: f(K) \subseteq U\}$, $\{f: fx \cap U \neq \emptyset \text{ for all } x \in K\}$, where K is a compact subset of X and U is open in Y . Obviously, τ_c is larger than τ_p .

A subset F of Y^{mX} satisfies the condition (G) if, for every τ_c -closed subset F_0 of F , $\bigcap_{f \in F_0} f^-(U)$ and $\bigcap_{f \in F_0} f^+(U)$ are open in X whenever U is open in Y . The following two lemmas relate this condition to even continuity:

LEMMA 3.4. *If Y is regular, then every subset of Y^{mX} satisfying the condition (G) is evenly continuous.*

Proof. Let F be a subset of Y^{mX} which satisfies the condition (G). Let $x \in X$, let K be a compact subset of Y and let V be an open neighborhood of K . Let W be an open neighborhood of K such that $K \subseteq W \subseteq \bar{W} \subseteq V$. Since $F_1 = \{h: h \in F \text{ and } hx \cap \bar{W} \neq \emptyset\}$, $F_2 = \{h: h \in F \text{ and } hx \subseteq \bar{W}\}$ are τ_c -closed in F , $U_1 = \bigcap_{h \in F_1} h^-(V)$ and $U_2 = \bigcap_{h \in F_2} h^+(V)$ are open in X . Then $U = U_1 \cap U_2$ is an open neighborhood of x .

Let $f \in F$ be such that $fx \cap W \neq \emptyset$. Then $f \in F_1$, so that $U \subseteq U_1 \subseteq f^-(V)$. Now let $f \in F$ be such that $fx \subseteq W$. Then $f \in F_2$, so that $U \subseteq U_2 \subseteq f^+(V)$.

LEMMA 3.5. *Every τ_c -compact evenly continuous subset of $(Y^{mX})_0$ satisfies the condition (G).*

Proof. Let F be a τ_c -compact evenly continuous subset of $(Y^{mX})_0$. Since F is τ_p -compact, it suffices, by Corollary 10.6 of [7, p. 23], to show that τ_p on F is jointly continuous. For this we apply Lemma 3.2.

Let X be a topological space and let $Y = (Y, \mathcal{U})$ be a uniform space. A subset F of Y^{mX} is *equicontinuous* if, for $(x, U) \in X \times \mathcal{U}$, there exists a neighborhood V of x such that, for all $f \in F$, $f(V) \subseteq U[fx]$ and $fz \cap U[y] \neq \emptyset$ whenever $(z, y) \in V \times fx$. The following two lemmas relate equicontinuity to even continuity:

LEMMA 3.6. *If $Y = (Y, \mathcal{U})$ is a uniform space, then every equicontinuous subset of Y^{mX} is evenly continuous.*

Proof. Let F be an equicontinuous subset of Y^{mX} . Let $x \in X$, let K be a compact subset of Y and let U be a symmetric member of \mathcal{U} . There is a neighborhood V of x such that, for all $f \in F$, $f(V) \subseteq$

$U[fx]$ and $fx \subseteq U[fz]$ for all $z \in V$. Write $W = U[K]$. Let $f \in F$ be such that $fx \cap W \neq \emptyset$. If $z \in V$ then $fx \subseteq U[fz]$, so that $U[fz] \cap W \neq \emptyset$, therefore $V \subseteq f^-(U^2[K])$. Now let $f \in F$ be such that $fx \subseteq W$. Then $f(V) \subseteq U[fx] \subseteq U^2[K]$, that is, $V \subseteq f^+(U^2[K])$.

LEMMA 3.7. *If Y is a uniform space, then every evenly continuous pointwise bounded subset of $(Y^{m^X})_0$ is equicontinuous.*

Proof. Let F be an evenly continuous pointwise bounded subset of $(Y^{m^X})_0$. Let \bar{F} denote the τ_p -closure of F in $(Y^{m^X})_0$. By the Lemmas 3.1, 3.2, τ_p on \bar{F} is jointly continuous. Since $(Y^{m^X})_0$ is a Tychonoff set, by Lemma 2.1, \bar{F} is τ_p -compact. Then, by the Lemma 8 of Smithson [9, p. 258], \bar{F} is equicontinuous.

4. Ascoli theorem. Let $X = (X, \tau)$ be a topological space. The k -extension of τ is the family $k(\tau)$ of all subsets U of X such that $U \cap K$ is open in K for every compact subset K of X . It is clear that $k(\tau)$ is a topology on X which is larger than τ . The topological space $kX = (X, k(\tau))$ is called the k -extension of X . A topological space X is called a k -space if $kX = X$. For an arbitrary topological space X , $kkX = kX$, so kX is a k -space. Familiar examples of k -spaces are the locally compact spaces and the spaces satisfying the first countability axiom.

Let X and Y be topological spaces. A function $f: X \rightarrow Y$ is called k -continuous if its restriction to each compact subset of X is continuous. Henceforth, the set of all continuous (k -continuous) functions on X to Y will be denoted $C(X, Y)$ ($C_k(X, Y)$). It can be shown that a topological space X is a k -space if and only if $C_k(X, Y) = C(X, Y)$ for every topological space Y [7, p. 9]. A topological space X is a k_3 -space if $C_k(X, Y) = C(X, Y)$ for every regular space Y . Thus a k -space is a k_3 -space but not conversely. In fact, the product of uncountably many copies of the real line, which is not a k -space, is a k_3 -space. We write $\mathcal{C}_0(X, Y) = (Y^{m^X})_0 \cap \mathcal{C}(X, Y)$.

We note that if Y is regular, then $(\mathcal{C}_0(X, Y), \tau_c)$ is a regular space for every topological space X .

In a regular space there was introduced in [7, p. 11] the following equivalence relation $R: xRy$ if every open neighborhood of x contains y . For a subset F of such a space, F^* denotes its R -saturation, that is, the smallest R -saturated set containing F .

THEOREM 4.1. *Let X, Y be topological spaces, let T be a Tychonoff subset of $(Y^{m^X})_0$ and let $F \subseteq (T \cap \mathcal{C}(X, Y), \tau_c)$. If Y is regular, the following conditions are sufficient for the compactness of F :*

- (a) F^* is closed in $T \cap \mathcal{C}(X, Y)$.
- (b) F is pointwise bounded, and
- (c) F is evenly continuous.

If X is a k -space and Y is regular, then the conditions (a), (b) and (c) are necessary for the compactness of F .

Proof. Sufficiency. Let \bar{F} denote the τ_p -closure of F in T . Since $T \subseteq (Y^{mX})_0$, (c) implies, by Lemmas 3.1 and 3.2, that $\omega: (\bar{F}, \tau_p) \times X \rightarrow Y$ is continuous, and, in particular, that $\bar{F} \subseteq \mathcal{C}(X, Y)$. By Lemma 8.1 of [7, p. 18], $\tilde{\omega}: (\bar{F}, \tau_p) \rightarrow (\mathcal{C}(X, Y), \tau_c)$ is continuous. Since T is a Tychonoff set, (b) implies, by Lemma 2.1, that \bar{F} is τ_p -compact, so $\tilde{\omega}(\bar{F}) = \bar{F}$ is a τ_c -compact subset of $T \cap \mathcal{C}(X, Y)$. But (a) implies $F \subseteq \bar{F} \subseteq F^*$, so, by Theorem 4.1 (b) of [7, p. 11], F is τ_c -compact.

Necessity. By Theorem 4.1 (c) of [7, p. 11], F^* is closed in $(T \cap \mathcal{C}(X, Y), \tau_c)$. It is clear that F is pointwise bounded. Since X is a k -space, by Theorem 9.4 of [7, p. 21], $\omega: (F, \tau_c) \times X \rightarrow Y$ is continuous. So by Lemma 3.3, $F = \{\omega(f, \cdot): f \in F\}$ is evenly continuous.

COROLLARY 1. Let $F \subseteq (\mathcal{C}_0(X, Y), \tau_c)$. If Y is regular, the following conditions are sufficient for the compactness of F :

- (a) F^* is closed in $\mathcal{C}_0(X, Y)$,
- (b) F is pointwise bounded, and
- (c) F is evenly continuous.

If X is a k -space and Y is regular, then the conditions (a), (b) and (c) are necessary for the compactness of F .

COROLLARY 2. Let $F \subseteq (C(X, Y), \tau_c)$. If Y is regular, the following conditions are sufficient for the compactness of F :

- (a) F^* is closed in $C(X, Y)$,
- (b) F is pointwise bounded, and
- (c) F is evenly continuous.

If X is a k -space and Y is regular, then the conditions (a), (b) and (c) are necessary for the compactness of F .

COROLLARY 3. If Y is regular, a subset F of $(C_k(X, Y), \tau_c)$ is compact if and only if

- (a) F^* is closed in $C_k(X, Y)$,
- (b) F is pointwise bounded, and
- (c) F is evenly continuous on compacta.

Proof. For the sufficiency, we note that $C_k(X, Y) = C(kX, Y)$ and apply the Lemma 3.4 of [7, p. 11]. For the necessity, we consider F as a subset of $(C(kX, Y), \tau_c)$ and deduce from Corollary 2 the conditions (a),

(b) and the even continuity of F . Then it is clear that F , considered as a subset of $(C_k(X, Y), \tau_c)$, is evenly continuous on compacta.

COROLLARY 4. ([1, p. 635]). *Let $F \subseteq (C(X, Y), \tau_c)$. If Y is regular, the following conditions are sufficient for the compactness of F :*

- (a) F is closed in $C(X, Y)$,
- (b) F is pointwise bounded, and
- (c) F is evenly continuous.

If X is a k_3 -space and Y is regular, then the conditions (a), (b) and (c) are necessary for the compactness of F .

Proof. For the necessity, we note that, since Y is regular, $C_k(X, Y) = C(X, Y)$; then we apply Corollary 3 and Lemma 3.4 of [7, p. 11].

REMARKS. (1) By Lemmas 3.4, 3.5, the Corollary 1 is equivalent to the Theorem 10.10 of [7, pp. 23–24], which contains the Ascoli theorem of Gale [2, p. 304] and the multifunction Ascoli theorem of Mancuso [6, p. 470].

(2) Let Y be a uniform space. By Lemmas 3.6, 3.7 and Theorem 12.2 of [7, p. 28], the Corollary 1 in this context is equivalent to the Theorem 12.8 of [7, p. 31], which contains the multifunction Ascoli theorem of Smithson [9, p. 259].

REFERENCES

1. G. Fox and P. Morales, *A non-Hausdorff Ascoli theorem for k_3 -spaces*, Proc. Amer. Math. Soc., **39** (1973), 633–636.
2. D. Gale, *Compact sets of functions and function rings*, Proc. Amer. Math. Soc., **1** (1950), 303–308.
3. J. Kelley, *General Topology*, New York 1965.
4. Y. F. Lin, *Tychonoff's theorem for the space of multifunctions*, Amer. Math. Monthly, **74** (1967), 399–400.
5. Y. F. Lin and D. A. Rose, *Ascoli's theorem for spaces of multifunctions*, Pacific J. Math., **34** (1970), 741–747.
6. V. J. Mancuso, *An Ascoli theorem for multi-valued functions*, J. Austral. Math. Soc., **12** (1971), 466–472.
7. P. Morales, *Non-Hausdorff Ascoli theory*, Dissertationes Math. 119 (1974), 1–37.
8. N. Noble, *Ascoli theorems and the exponential map*, Trans. Amer. Math. Soc., **143** (1969), 393–411.
9. R. Smithson, *Uniform convergence for multifunctions*, Pacific J. Math., **39** (1971), 253–259.

Received June 11, 1975.

UNIVERSITÉ DE MONTRÉAL

| | |
|---|-----|
| Walter Allegretto, <i>Nonoscillation theory of elliptic equations of order $2n$</i> | 1 |
| Bruce Allen Anderson, <i>Sequencings and starters</i> | 17 |
| Friedrich-Wilhelm Bauer, <i>A shape theory with singular homology</i> | 25 |
| John Kelly Beem, <i>Characterizing Finsler spaces which are pseudo-Riemannian of constant curvature</i> | 67 |
| Dennis K. Burke and Ernest A. Michael, <i>On certain point-countable covers</i> | 79 |
| Robert Chen, <i>A generalization of a theorem of Chacon</i> | 93 |
| Francis H. Clarke, <i>On the inverse function theorem</i> | 97 |
| James Bryan Collier, <i>The dual of a space with the Radon-Nikodým property</i> | 103 |
| John E. Cruthirds, <i>Infinite Galois theory for commutative rings</i> | 107 |
| Artatrana Dash, <i>Joint essential spectra</i> | 119 |
| Robert M. DeVos, <i>Subsequences and rearrangements of sequences in FK spaces</i> | 129 |
| Geoffrey Fox and Pedro Morales, <i>Non-Hausdorff multifunction generalization of the Kelley-Morse Ascoli theorem</i> | 137 |
| Richard Joseph Fleming, Jerome A. Goldstein and James E. Jamison, <i>One parameter groups of isometries on certain Banach spaces</i> | 145 |
| Robert David Gulliver, II, <i>Finiteness of the ramified set for branched immersions of surfaces</i> | 153 |
| Kenneth Hardy and István Juhász, <i>Normality and the weak cb property</i> | 167 |
| C. A. Hayes, <i>Derivation of the integrals of $L^{(q)}$-functions</i> | 173 |
| Frederic Timothy Howard, <i>Roots of the Euler polynomials</i> | 181 |
| Robert Edward Jamison, II, Richard O'Brien and Peter Drummond Taylor, <i>On embedding a compact convex set into a locally convex topological vector space</i> | 193 |
| Andrew Lelek, <i>An example of a simple triod with surjective span smaller than span</i> | 207 |
| Janet E. Mills, <i>Certain congruences on orthodox semigroups</i> | 217 |
| Donald J. Newman and A. R. Reddy, <i>Rational approximation of e^{-x} on the positive real axis</i> | 227 |
| John Robert Quine, Jr., <i>Homotopies and intersection sequences</i> | 233 |
| Nambury Sitarama Raju, <i>Periodic Jacobi-Perron algorithms and fundamental units</i> | 241 |
| Herbert Silverman, <i>Convexity theorems for subclasses of univalent functions</i> | 253 |
| Charles Frederick Wells, <i>Centralizers of transitive semigroup actions and endomorphisms of trees</i> | 265 |
| Volker Wrobel, <i>Spectral approximation theorems in locally convex spaces</i> | 273 |
| Hidenobu Yoshida, <i>On value distribution of functions meromorphic in the whole plane</i> | 283 |