

Pacific Journal of Mathematics

HOMOTOPIES AND INTERSECTION SEQUENCES

JOHN ROBERT QUINE, JR.

HOMOTOPIES AND INTERSECTION SEQUENCES

J. R. QUINE

For $\gamma_t: S^1 \rightarrow \mathbb{C}$, a smooth homotopy of closed curves, the changing configuration of vertices and cusps is studied by considering the set in $I \times S^1 \times S^1$ given by $(\gamma_t(z) - \gamma_t(\zeta))/(z - \zeta) = 0$. The main tool is oriented intersection theory from differential topology. The results relate to previous work by Whitney and Titus on normal curves and intersection sequences.

Consider a closed curve as a smooth map $\gamma: S^1 \rightarrow \mathbb{C}$. Let γ_t for $t \in I$ be a smooth homotopy of closed curves. A vertex of γ_t is a point w such that $w = \gamma_t(z) = \gamma_t(\zeta)$ for $z \neq \zeta$. A cusp is a point where the tangent vanishes and changes direction. Let $X = I \times S^1 \times S^1$. We study the changing configuration of vertices and cusps of γ_t by studying the set $Z = \{x \in X \mid G(x) = 0\}$ where $G(t, z, \zeta) = (\gamma_t(z) - \gamma_t(\zeta))/(z - \zeta)$, and the limiting value is taken when $z = \zeta$. If 0 is a regular value for G , then Z has the structure of an oriented 1-submanifold of X . If for fixed t , Z intersects $t \times S^1 \times S^1$ transversely, then the oriented intersection gives a set of pairs in $S^1 \times S^1$ with corresponding orientation numbers $+1$ or -1 . If γ_t is a normal immersion, these pairs and their orientation numbers give the Titus intersection sequence of γ_t . The changes in the intersection sequence are reflected in the behavior of Z . If Z crosses $I \times \Delta$, where Δ is the diagonal of $S^1 \times S^1$, then we have a cusp and a change in the tangent winding number. The difference between the tangent winding numbers of γ_0 and γ_1 is just $N(Z, I \times \Delta)$, the total number of oriented intersections of Z with $I \times \Delta$.

1. Intersection sequences. In the complex plane, let S^1 be the set $|z| = 1$. Consider S^1 as a 1-manifold with functions $\theta \rightarrow e^{i\theta}$ giving local coordinate systems. The tangent vector $d/d\theta$ is defined independently of the choice of coordinate system. On $T(S^1)$, the tangent space, let $d/d\theta$ give the positive orientation at each point. This gives S^1 the structure of an oriented 1-manifold.

Suppose $\gamma: S^1 \rightarrow \mathbb{C}$ is a smooth (C^∞) map. Let $\beta(z) = (d\gamma/d\theta)(z)$ be the tangent at $\gamma(z)$. Let $S^1 \times S^1 = Y$ and let the maps $(\theta, \phi) \rightarrow (e^{i\theta}, e^{i\phi})$ give local coordinate systems for Y . Let $S^1 \times S^1$ have the product orientation, i.e., $T(S^1 \times S^1)$ has positive orientation given by the ordered basis $\{\partial/\partial\theta, \partial/\partial\phi\}$ at each point. Let $\Delta \subseteq Y = \{(z, \zeta) \mid z = \zeta\}$.

Let $\theta \rightarrow (e^{i\theta}, e^{i\theta})$ be local coordinate systems on Δ and let positive

orientation be given on Δ by $d/d\theta$. Thus Δ is an oriented 1-submanifold of Y . Now we define $g: Y \rightarrow \mathbf{C}$ as follows

$$g(z, \zeta) = \begin{cases} \frac{\gamma(z) - \gamma(\zeta)}{z - \zeta}, & z \neq \zeta \\ \frac{-i\beta(z)}{z}, & z = \zeta. \end{cases}$$

We can check that g is a smooth function on Y .

Letting $y = (z, \zeta)$, we compute that for $y \in g^{-1}(0)$ we have

$$(1) \quad dg_y = \begin{cases} \frac{\beta(z)d\theta - \beta(\zeta)d\phi}{z - \zeta}, & z \neq \zeta \\ \frac{1}{iz} \frac{d\beta}{d\theta}(z)(d\theta + d\phi), & z = \zeta. \end{cases}$$

Now let $y = (z, \zeta) \in g^{-1}(0)$, and consider dg_y as a linear map from $T_y(Y)$ to $T_0(\mathbf{C})$. Then from (1):

(a) If $z \neq \zeta$, then dg_y has rank 2 iff the tangents $\beta(z)$ and $\beta(\zeta)$ are linearly independent. In this case, dg_y preserves orientation iff $\{\beta(z), -\beta(\zeta)\}$ is a positively oriented basis of \mathbf{C} (where \mathbf{C} has the usual orientation).

(b) If $z = \zeta$, then $\beta(z) = 0$ and dg_y has rank 1 iff $(d\beta/d\theta)(z) \neq 0$. Otherwise dg_y has rank 0. We may check that if $(d\beta/d\theta)(z) \neq 0$, then there is a cusp at $\gamma(z)$ and the limiting tangential directions at $\gamma(z)$ are the directions of $\pm(d\beta/d\theta)(z)$.

The point $0 \in \mathbf{C}$ is said to be a regular value for g if dg_y has rank 2 at every point of $g^{-1}(0)$. By remarks (a) and (b) above we see that 0 is a regular value for g iff γ is an immersion ($\beta(z) \neq 0$ for $z \in S^1$), and the tangents $\beta(z)$ and $\beta(\zeta)$ are linearly independent for each point $(z, \zeta) \in g^{-1}(0)$. Also if 0 is a regular value of g , $g^{-1}(0)$ is a finite subset of the compact set Y (a torus). In this case if $y \in g^{-1}(0)$ we set $\lambda(y) = +1$ if dg_y preserves orientation and $\lambda(y) = -1$ if dg_y reverses orientation. We say that $g^{-1}(0)$ with the sign λ gives the set of signed intersection pairs for γ .

We say that γ is a normal immersion if γ is an immersion, each point of \mathbf{C} has at most two preimages under γ , and the tangents are linearly independent at each double point. Another way to say this is that 0 is a regular value for g , and projection on the first coordinate is one-to-one on $g^{-1}(0)$. ($g^{-1}(0)$ as a set of ordered pairs is a function.) If γ is a normal immersion, let $\{z_1, \dots, z_{2n}\}$ be the preimages under γ of the double points, numbered sequentially along S^1 in a counterclockwise direction from a point z_0 on S^1 , not a preimage of a double point. Then $g^{-1}(0)$ defines an involution $*$ on the integers $1, \dots, 2n$, such that $(z_i, z_{i'}) \in g^{-1}(0)$

for $j = 1, \dots, 2n$. Now define the sign ν by $\nu(j) = -\lambda((z_j, z_{j^*}))$. We say that the involution $*$ together with the sign ν defines the intersection sequence of γ with respect to z_0 . Usually z_0 is chosen so that $\gamma(z_0)$ is on the outer boundary, i.e., the boundary of the component of $\mathbb{C} - \gamma(S^1)$ containing ∞ . In this case ν and $*$ give the Titus intersection sequence (see Titus [5] or Francis [1]). We remark that signed intersection pairs are defined if 0 is a regular value for g . To define the intersection sequence also, we need in addition that $g^{-1}(0)$ is a function.

2. The fundamental theorem. In this context, we would like to prove what we call the fundamental theorem on intersection sequences. The use of intersection pairs allows a slightly more general statement than that of Whitney [6] and Titus [5]. Let γ be a normal immersion and let $[\gamma]$ denote the image of γ . For $a \in \mathbb{C} - \gamma(R)$ we define j_a on $S_a = S^1 - \gamma^{-1}(a)$ by $j_a = (\gamma - a)/|\gamma - a|$. We define

$$\omega(\gamma, a) = \frac{1}{2\pi i} \int_{S_a} \frac{dj_a}{j_a}.$$

If $a \notin \gamma$, this is just the winding number of γ about a . If $a \in [\gamma]$, we may check that $\omega(\gamma, a)$ is the average of the winding numbers of γ on the components near $\gamma(a)$.

Now, for fixed $z_0 \in S^1$, consider $z_0 \times S^1$ and $S^1 \times z_0$ as subsets of Y . Let $\theta \rightarrow (z_0, e^{i\theta})$ and $\phi \rightarrow (e^{i\phi}, z_0)$ be coordinate systems on $z_0 \times S^1$ and let these define the orientations. Thus, $z_0 \times S^1$ and $S^1 \times z_0$ have the structures of oriented 1-submanifolds of Y . Now $W = z_0 \times S^1 + S^1 \times z_0 - \Delta$ divides the torus Y into 2 simply connected 2-manifolds with boundary, Y^+ and Y^- . Here Y^+ denotes the one for which W is a positively oriented boundary and Y^- the one for which W is a negatively oriented boundary (see Fig. 1).




FIG. 1

If γ is an immersion, and $\beta = d\gamma/d\theta$ is the tangent, then the tangent winding number, $tw\ \gamma$, is defined to be

$$\frac{1}{2\pi i} \int_{S^1} \frac{d\beta}{\beta}.$$

We now have

THEOREM 1 (Titus–Whitney). *If 0 is a regular value for g , $z_0 \in S^1$, and Y^+ is the oriented 2-submanifold of $S^1 \times S^1$ with positively oriented boundary $z_0 \times S^1 + S^1 \times z_0 - \Delta$, then*

$$tw\ \gamma = - \sum_{y \in Y^+ \cap g^{-1}(0)} \lambda(y) + 2\omega(\gamma, \gamma(z_0)).$$

Proof. Let $g^{-1}(0) \cap Y^+ = \{y_1, \dots, y_n\}$. Let D_1, \dots, D_n be closed disjoint coordinate discs in Y^+ such that $D_j \cap g^{-1}(0) = Y_j$ for $j = 1, \dots, n$. Let these have orientation inherited from Y and let ∂D_j be the oriented boundary of D_j for $j = 1, \dots, n$. Recall that for $j = 1, \dots, n$, $\lambda(y_j) = +1$ iff dg preserves orientation at y_j . Therefore we may choose each D_j so that

$$\frac{1}{2\pi i} \int_{D_j} \frac{dg}{g} = \lambda(y_j).$$

Now dg/g is closed on $Y^+ - \bigcup_{j=1}^n D_j$ so the integral of dg/g over its boundary is 0. The boundary is the cycle $z_0 \times S^1 + S^1 \times z_0 - \Delta - \sum_{j=1}^n \partial D_j$. From the definition of g ,

$$\frac{1}{2\pi i} \int_{z_0 \times S^1} \frac{dg}{g} = \frac{1}{2\pi i} \int_{S^1 \times z_0} \frac{dg}{g} = \omega(\gamma, \gamma(z_0)) - 1/2$$

and $(1/2\pi i) \int_{\Delta} dg/g = tw\ \gamma - 1$. The theorem now follows. We remark that if $\gamma(z_0)$ is on the outer boundary of γ and its image is not a multiple point of γ , then $\omega(\gamma, \gamma(z_0)) = \pm \frac{1}{2}$. In this case, if γ is a normal immersion, then Theorem 1 is Lemma 3 of Titus [5].

3. Homotopies. Let $I = [0, 1]$ considered as an oriented 1-manifold with boundary having the usual orientation. Let $I \times S^1$ be an oriented 2-manifold with boundary with the product orientation. A smooth map $F: I \times S^1 \rightarrow \mathbf{C}$ is called a homotopy. Let $\gamma_t(z) = F(t, z)$ and $\beta_t(z) = (d\gamma_t/d\theta)(z)$. Let $X = I \times S^1 \times S^1$ and $Y_t = t \times S^1 \times S^1 \subseteq X$ where both are given the product orientations. Define $G: X \rightarrow \mathbf{C}$ by

$$G(t, z, \zeta) = \begin{cases} \frac{F(t, z) - F(t, \zeta)}{z - \zeta}, & z \neq \zeta \\ \frac{-i\beta_i(z)}{z}, & z = \zeta. \end{cases}$$

Define $g_i: S^1 \times S^1 \rightarrow \mathbf{C}$ by $g_i(z, \zeta) = G(t, z, \zeta)$. Let $Z = \{x \in X \mid G(x) = 0\}$. We say 0 is a regular value for G if dG has rank 2 everywhere on Z . In this case, by the implicit function theorem, Z has the structure of a 1-submanifold of X , with boundary. We intend to study the change in the intersection sequence under the homotopy F by looking at the smooth manifold $Z \subseteq X$, therefore we will make the assumption that 0 is a regular value for G .

To justify this assumption, we prove the following lemma.

LEMMA 1. *If $F(t, z) = \gamma_i(z)$ is a smooth homotopy of closed curves and $G(F): I \times S^1 \times S^1 \rightarrow \mathbf{C}$ is defined by $G(F)(t, z, \zeta) = (F(t, z) - F(t, \zeta))/(z - \zeta)$, then F may be deformed by an arbitrarily small amount into a homotopy F for which 0 is a regular value for $G(F)$.*

Proof. Let D be the open disc $|w| < 1$. For $w \in D$, define $F_w(t, z) = F(t, z) + wz$. Note that $F_0(t, z) = F(t, z)$. Then $G(F_w)(t, z, \zeta) = G(F)(t, z, \zeta) + w$. Clearly the map $(t, z, \zeta, w) \rightarrow G(F_w)(t, z, \zeta) + w$ from $(I \times S^1 \times S^1) \times D$ to \mathbf{C} is a submersion, and therefore 0 is a regular value for this function. By the transversality theorem (Guillemin and Pollack [3] p. 68), 0 is a regular value of $G(F_w)$ for almost all $w \in D$. This proves the lemma.

4. The orientation on Z . Assume that 0 is a regular value of G so that Z is a 1-manifold with boundary. We will define an orientation on Z such that we get a set of signed intersection pairs for γ_i by intersecting Z with Y_r . At each intersection point, the sign will be defined by the orientation of Z and Y_r .

First we indicate how to define a direct sum orientation on vector spaces. If V and W are oriented subspaces of a vector space and if the ordered bases $\{v_1, \dots, v_n\}$ and $\{w_1, \dots, w_m\}$ define positive orientation of V and W respectively, then the sum orientation on $V \oplus W$ (in that order) is defined by the ordered basis $\{v_1, \dots, v_n, w_1, \dots, w_m\}$.

We now orient Z as follows: If $x \in Z$, write $T_x(X) = T_x(Z) \oplus H$. Then $dG_x: H \rightarrow T_0(\mathbf{C})$ and the mapping is a vector space isomorphism. In a natural way, this isomorphism induces an orientation on H from the usual orientation on $T_0(\mathbf{C})$. We now choose an orientation on $T_x(Z)$ so that the sum orientation agrees with the prescribed orientation on $T_x(X)$. In this way Z is given the structure of an oriented 1-manifold.

Now as before let $Y_i = I \times S^1 \times S^1$ with the product orientation. Suppose $x = (t, z, \zeta) \in Z \cap Y_i$ and $d(g_i)_{(z, \zeta)}$ preserves orientation. Then dG_x preserves orientation on $T_x(Y_i)$. Now we can write $T_x(X) = T_x(Z) \oplus T_x(Y_i)$ where by definition, the orientations sum to the prescribed orientation on $T_x(X)$. In this case the intersection number at $x \in Z \cap Y_i$ is said to be $+1$ (here the order in which we list Z and Y_i is important (see Guillemin and Pollack [3])). Likewise if $d(g_i)_{(z, \zeta)}$ reverses orientation, the intersection number of $x \in Z \cap Y_i$ is -1 . Thus if $d(g_i)_{(z, \zeta)}$ has rank 2 at each point $x \in Z$ then the set $Z \cap Y_i$ along with the intersection number at each point gives us the set of signed intersection pairs for γ_i .

5. The change in the intersection sequences. The configuration of the oriented 1-manifold Z as a submanifold of X indicates how the intersection pairs and the intersection sequence changes under the homotopy F . (We may take the intersection sequence with respect to a continuously moving point whose image stays on the outer boundary.) We mention here only some general considerations:

(a) Z is symmetric with respect to $I \times \Delta$, i.e., $(t, z, \zeta) \in Z$ iff $(t, \zeta, z) \in Z$.

(b) The components of Z are oriented 1-manifolds homeomorphic to either S^1 or I (see Guillemin and Pollack [3] Appendix 2 or Milnor [4] Appendix).

(c) Each component either crosses $I \times \Delta$ and is symmetric with respect to $I \times \Delta$ or has another component symmetric to it with respect to $I \times \Delta$ (see Fig. 2).

(d) When a component of Z crosses $I \times \Delta$ we have a change in $\text{tw} \gamma_i$. We will describe this fully in the next section.

(e) Each component of Z represents a continuously moving vertex on γ_i . Components homeomorphic to I and joining points on Y_0 represent vertices lost in homotopy. Components homeomorphic to I and joining points in Y_1 represent vertices gained.




FIG. 2

Finally, suppose that $\Pi: X = I \times S^1 \times S^1 \rightarrow I \times S^1$ is the projection on the first two coordinates. Then $\Pi(Z) \subseteq I \times S^1$ consists of smooth curves. If the intersection sequence of γ_t changes at t_0 , then either some vertices coincide, in which case $\Pi(Z)$ crosses itself at a point (t_0, z) or else a vertex appears or disappears, in which case the real valued function t on Z has a relative maximum or minimum at a point (t_0, z, ζ) on Z .

6. Change in *tw* γ_t . Let $I \times \Delta \subseteq X$ have the usual product orientation. Say Z intersects $I \times \Delta$ transversely if $T_x(Z) \oplus T_x(I \times \Delta) = T_x(X)$ at each point $x \in Z \cap (I \times \Delta)$. Let $N(Z, I \times \Delta)$ be the intersection multiplicity of Z with $I \times \Delta$, i.e., the sum of the intersection numbers at points of $Z \cap (I \times \Delta)$. We prove the following theorem concerning the change in *tw* γ_t for the homotopy.

THEOREM 2. *If Z intersects $I \times \Delta$ transversely, then $\text{tw } \gamma_1 - \text{tw } \gamma_0 = N(Z, I \times \Delta)$.*

Proof. Let $Z \cap (I \times \Delta) = \{y_1, \dots, y_n\}$. At $y = y_j$ write $T_y(X) = T_y(Z) \oplus T_y(I \times \Delta)$. By definition of the intersection number at y_j and by definition of the orientation of Z we see that the intersection number at $y = y_j$ is $+1$ iff dG_y preserves orientation on $T_y(I \times \Delta)$. Now we can choose closed disjoint coordinate discs D_1, \dots, D_n in $I \times \Delta$ such that $D_j \cap Z = y_j$ for $j = 1, \dots, n$ and $(1/2\pi i) \int_{\partial D_j} dG/G =$ the orientation number at $y_j \in Z \cap (I \times \Delta)$. Now dG/G is closed on $I \times \Delta - \bigcap_{j=1}^n D_j$ and the boundary is $1 \times \Delta - 0 \times \Delta - \sum_{j=1}^n \partial D_j$. Now $(1/2\pi i) \int_{0 \times \Delta} dG/G = \text{tw } \gamma_0$ and $(1/2\pi i) \int_{1 \times \Delta} dG/G = \text{tw } \gamma_1$, and integration of dG/G over the boundary gives 0. This proves the theorem.

We have the following well-known:

COROLLARY 1. *Regular homotopies preserve the tangent winding number.*

Proof. In this case $Z \cap (I \times \Delta) = \emptyset$.

Finally, we remark that the fundamental theorem of Titus and Whitney becomes in this context:

THEOREM 3. *Suppose for fixed $t \in I$ and $z_0 \in S^1$, Y_t^+ is the oriented submanifold of $I \times S^1 \times S^1$ with positively oriented boundary $t \times z_0 \times S^1 + t \times S^1 \times z_0 - t \times \Delta$. If Z intersects Y_t^+ transversely,*

$$N(Z, Y_t^+) = \text{twn } \gamma_t - 2\omega(\gamma_t, \gamma_t(z_0)).$$

Proof. We observe that if $\dot{x} = (t, z, \zeta) \in Z \cap Y_t$ then the intersection number is +1 iff $d(g_t)_{(z, \zeta)}$ preserves orientation. Now the theorem follows from Theorem 1.

REFERENCES

1. G. Francis, *Null genus realizability criterion for abstract intersection sequences*, J. Combinatorial Theory, **7** (1969), 331–341.
2. ———, *Titus' homotopies of normal curves*, Proc. Amer. Math. Soc., **30** (1971), 511–518.
3. V. Guillemin and A. Pollack, *Differential Topology*, Prentice-Hall, Inc., Englewood Cliffs, New Jersey, 1974.
4. J. Milnor, *Topology from the Differentiable Viewpoint*, The University Press of Virginia, Charlottesville, 1965.
5. C. J. Titus, *A theory of normal curves and some applications*, Pacific J. Math., **10** (1960), 1083–1096.
6. H. Whitney, *On regular closed curves in the plane*, Comp. Math., **4** (1937), 276–284.

Received November 5, 1975 and in revised form February 24, 1976.

THE FLORIDA STATE UNIVERSITY

Pacific Journal of Mathematics

Vol. 64, No. 1

May, 1976

Walter Allegretto, <i>Nonoscillation theory of elliptic equations of order $2n$</i>	1
Bruce Allem Anderson, <i>Sequencings and starters</i>	17
Friedrich-Wilhelm Bauer, <i>A shape theory with singular homology</i>	25
John Kelly Beem, <i>Characterizing Finsler spaces which are pseudo-Riemannian of constant curvature</i>	67
Dennis K. Burke and Ernest A. Michael, <i>On certain point-countable covers</i>	79
Robert Chen, <i>A generalization of a theorem of Chacon</i>	93
Francis H. Clarke, <i>On the inverse function theorem</i>	97
James Bryan Collier, <i>The dual of a space with the Radon-Nikodým property</i>	103
John E. Cruthirds, <i>Infinite Galois theory for commutative rings</i>	107
Artatrana Dash, <i>Joint essential spectra</i>	119
Robert M. DeVos, <i>Subsequences and rearrangements of sequences in FK spaces</i>	129
Geoffrey Fox and Pedro Morales, <i>Non-Hausdorff multifunction generalization of the Kelley-Morse Ascoli theorem</i>	137
Richard Joseph Fleming, Jerome A. Goldstein and James E. Jamison, <i>One parameter groups of isometries on certain Banach spaces</i>	145
Robert David Gulliver, II, <i>Finiteness of the ramified set for branched immersions of surfaces</i>	153
Kenneth Hardy and István Juhász, <i>Normality and the weak cb property</i>	167
C. A. Hayes, <i>Derivation of the integrals of $L^{(q)}$-functions</i>	173
Frederic Timothy Howard, <i>Roots of the Euler polynomials</i>	181
Robert Edward Jamison, II, Richard O'Brien and Peter Drummond Taylor, <i>On embedding a compact convex set into a locally convex topological vector space</i>	193
Andrew Lelek, <i>An example of a simple triod with surjective span smaller than span</i>	207
Janet E. Mills, <i>Certain congruences on orthodox semigroups</i>	217
Donald J. Newman and A. R. Reddy, <i>Rational approximation of e^{-x} on the positive real axis</i>	227
John Robert Quine, Jr., <i>Homotopies and intersection sequences</i>	233
Nambury Sitarama Raju, <i>Periodic Jacobi-Perron algorithms and fundamental units</i>	241
Herbert Silverman, <i>Convexity theorems for subclasses of univalent functions</i>	253
Charles Frederick Wells, <i>Centralizers of transitive semigroup actions and endomorphisms of trees</i>	265
Volker Wrobel, <i>Spectral approximation theorems in locally convex spaces</i>	273
Hidenobu Yoshida, <i>On value distribution of functions meromorphic in the whole plane</i>	283