

Pacific Journal of Mathematics

**CENTRALIZERS OF TRANSITIVE SEMIGROUP ACTIONS AND
ENDOMORPHISMS OF TREES**

CHARLES FREDERICK WELLS

CENTRALIZERS OF TRANSITIVE SEMIGROUP ACTIONS AND ENDOMORPHISMS OF TREES

CHARLES WELLS

A tree is *locally finite* if the interval between any two points is finite. A *local isomorphism* of a tree with itself is a homomorphism which is an isomorphism when restricted to any interval. Two theorems are proved. One characterizes those locally finite trees which have transitive automorphism groups, and those which have transitive local-isomorphism monoids. The other theorem gives necessary and sufficient conditions for a non-injective transformation to be centralized by a transitive permutation group, and necessary and sufficient conditions for it to be centralized by a transitive transformation semigroup. Also, an example is given of a nonlocally-finite tree with transitive automorphism group.

1. Preliminaries and statements of the theorems. All functions will be written to the right of the argument, and functional composition will read from left to right. If $\alpha: X \rightarrow X$ is a function, and $x \in X$, then $x\alpha^{-1}$ denotes the inverse image of x under α . If X is any set, $|X|$ denotes the cardinality of X .

A semigroup S acts on a set X on the right if, for every $x \in X$ and $s \in S$, xs denotes an element of X , and

$$(1) \quad (xs)t = x(st) \quad (x \in X, s \in S).$$

Then X is an S -set or an S -operand.

An action by a semigroup S on X is *transitive* if for every $x, y \in X$, there is an element $s \in S$ such that $xs = y$. A subset G of X generates the S -set X if for every $x \in X$ there is $g \in G$ and $s \in S$ such that $gs = x$. Thus an S -set is transitive if and only if every one-element subset of X generates X .

Let X and Y be S -sets. A function $\alpha: X \rightarrow Y$ is an S -homomorphism (equivariant map) if

$$(2) \quad (x\alpha)s = (xs)\alpha \quad (x \in X, s \in S).$$

S -endomorphisms and S -automorphisms are defined in the obvious way. It is easy to see that the S -endomorphisms of an S -set X form a semigroup $\text{End}_S X$ and the S -automorphisms form a group $\text{Aut}_S X$.

Let T be a partially ordered set; its order relation, like all those in

this paper, will be denoted \leq . T is a *tree* if the set $tA \stackrel{\text{def}}{=} \{u \in T: t < u\}$ is well-ordered (whenever it is nonempty) for every $t \in T$. T is *nontrivial* if tA is nonempty for some $t \in T$. An element of tA is an *ancestor* of t .

If T is a tree, define for each $t \in T$:

tP = the *parent* of t = the minimum of the set $tA - \{t\}$, if it exists.

$tB = tPP^{-1}$ = the set of *siblings* of t .

$tD = \{u \in T: u < t\}$ = the set of *descendants* of t .

$tC = tP^{-1}$ = the set of *children* of t .

Observe that tC may be empty even if tD is nonempty.

If $t, u \in T$, the *interval from t to u* is the set $[t, u] = \{x \in T: t \leq x \leq u\}$; the notation will be used only when $t \leq u$. T is *locally finite* if $[t, u]$ is a finite set for all $t, u \in T$ with $t \leq u$. (This could be called *vertical* local finiteness, contrasted with *horizontal* local finiteness which would require that every sibling class be finite).

A *filter* of T is a subset X such that if $x \in X$ and $x < y$, then $y \in X$. X is an *ideal* of T if $x \in X$ and $y < x$ implies $y \in X$.

Let T and T' be trees. A bijection $f: T \rightarrow T'$ is an *isomorphism* if $t \leq u \Leftrightarrow tf \leq uf$ for all $t, u \in T$. A function $g: T \rightarrow T'$ is a *local isomorphism* if g restricted to $[t, u]$ is an isomorphism from $[t, u]$ to $[tg, ug]$ for all $t, u \in T$ with $t \leq u$. The set of isomorphisms of a tree T with itself forms a group $\text{Aut } T$ under functional composition, and similarly the set of local isomorphisms of T with itself forms a semigroup $LI(T)$.

Finally, let ω denote the set of nonnegative integers regarded as a well-ordered set.

In this paper, the following two theorems will be proved.

THEOREM 1. *Let T be a nontrivial locally finite tree. Then*

(a) *$LI(T)$ is transitive if and only if T has no maximal or minimal elements.*

(b) *$\text{Aut } T$ is transitive if and only if T has no maximal or minimal elements and $|tB| = |uB|$ for all $t, u \in T$.*

THEOREM 2. *Let S be a semigroup and X a transitive S -set with S -endomorphism α . Then α is surjective, and if α is not injective, then α^k has no fixed points for any positive integer k . Conversely, if $\alpha: X \rightarrow X$ is a surjective function with the property that α^k has no fixed points for any positive integer k , then there is a transitive semigroup of functions from X to X with respect to which α is an endomorphism (in other words, α commutes with every function of the semigroup). This semigroup of functions can be taken to be a group of permutations if and only if $|x\alpha^{-1}| = |y\alpha^{-1}|$ for all $x, y \in X$.*

Observe that as a corollary of Theorem 2, a transitive semigroup action on a finite set can only be centralized by permutations.

2. Proof of Theorem 1. We need three lemmas.

LEMMA 1. *Let T be a tree. Then T is locally finite and has no maximal elements if and only if tA is order-isomorphic to ω for every $t \in T$.*

Proof. Let T be locally finite and have no maximal elements. Let $t \in T$. Define $\beta: \omega \rightarrow tA$ by

$$\begin{aligned} 0\beta &= t \\ (n + 1)\beta &= (n\beta)P \qquad (n = 1, 2, \dots). \end{aligned}$$

Then for $k \in \omega$, $k\beta$ is clearly tP^k . It is easy to see that β is an injective, order-preserving map. It is also surjective: If $u \in tA$, then $[t, u]$ is finite, so that $u = tP^k$ for some integer k ; hence $u = k\beta$.

The converse is obvious.

LEMMA 2. *Let T and T' be trees, and $f: T \rightarrow T'$ a function. Then the following are equivalent:*

- (a) *f is a local isomorphism.*
- (b) *For all $t \in T$, f restricted to tA is an injective order-preserving function onto an initial segment of tfA .*
- (c) *For all $t, u \in T$ and $w \in T'$,*
 - (i) *$t < u \Rightarrow tf < uf$*
 - and (ii) *$tf < w < uf \Rightarrow$ there is $v \in T$ such that $t < v < u$ and $vf = w$.*

Proof. The proof, particularly in the order (a) \Rightarrow (c) \Rightarrow (b) \Rightarrow (a), is an easy consequence of the definitions of the terms involved and is omitted.

LEMMA 3. *Let T be a tree, $t \in T$ and $f \in LI(T)$. Then $tfP = tPf$.*

Proof. Immediate from Lemma 2(c).

We now prove the forward half of Theorem 1(a). Let $LI(T)$ be transitive. Since T is nontrivial, there are $t, u \in T$ with $t < u$. If $x \in T$, there are $f, g \in LI(T)$ with $tf = x$ and $ug = x$. Then $tg < x < uf$, so that x is neither maximal nor minimal.

The forward half of Theorem 1(b) now follows, since if $\text{Aut } T$ is transitive, $LI(T)$ certainly is, and an automorphism restricted to a sibling set must be a bijection onto another sibling set.

Now for the converse half of Theorem 1(b). Let T be a locally finite tree with no maximal or minimal elements. Let $t, u \in T$. We must construct an automorphism f of T with $tf = u$. To do this, we first construct a sequence f_0, f_1, f_2, \dots of functions with increasing domains D_0, D_1, \dots .

Take f_0 to be an order isomorphism from tA to uA ; it exists (in fact it is unique) by Lemma 1, and clearly $tf_0 = u$. The domain D_0 of f_0 is thus tA .

Let $D_1 = D_0 \cup (\bigcup_{x \in D_0} xB)$. For each $x \in D_0$, let $g_x: xB \rightarrow xf_0B$ be a bijection such that $xg_x = xf_0$. Define $f_1 = f_0 \cup (\bigcup_{x \in D_0} g_x)$. Then f_1 is well-defined because for $x, y \in D_0$, xB and yB are disjoint if $x \neq y$, and xB intersects D_0 precisely in $\{x\}$.

It is not difficult to show that (3), (4) and (5) below hold for $i = 1$.

- (3) $D_{i-1} \subset D_i$ and $f_i|_{D_{i-1}} = f_{i-1}$.
- (4) D_i and $D_i f_i$ are both filters of T .
- (5) f_i is an isomorphism from D_i to $D_i f_i$.

Now suppose that f_i has been defined with domain D_i so that (3), (4) and (5) are true. Suppose also $i > 0$.

We define f_{i+1} and D_{i+1} as follows. Let M_i denote the set of minimal elements of D_i (M_i is nonempty, but we do not need that fact). Let $E_i = \bigcup_{x \in M_i} xC$; E_i is the set of immediate descendants of elements of M_i . It is important to note that E_i is the *disjoint* union of the xC 's.

For each $x \in M_i$, let $g_x: xC \rightarrow xf_i C$ be a bijection. (Note: xC is nonempty because $[y, x]$ is finite for any $y < x$.) Let $D_{i+1} = D_i \cup E_i$, and $f_{i+1} = f_i \cup (\bigcup_{x \in M_i} g_x)$. Then f_{i+1} is well-defined because of the fact that E_i is a disjoint union of the xC 's. It is straightforward to check (it involves numerous cases) that f_{i+1} and D_{i+1} satisfy (3) through (5) with $i + 1$ instead of i .

Now let f_ω be the union of the functions f_i . Condition (3) implies that f_ω is well-defined, and conditions (4) and (5) that f_ω is an isomorphism between filters of T . Actually, more is true: The domain and the image of f_ω are each both filters and ideals of T . To see the latter, observe that if $x \in D_i$, then any element which is k levels below x will be included in D_{i+k} ; thus every element below x is included because of local finiteness. An analogous argument works for the image of f_ω .

Furthermore, either $\text{dom} f_\omega = \text{im} f_\omega$ or they are disjoint. (In fact, they are connected components of T). For let $w, x, y \in \text{dom} f_\omega$ and $xf_\omega = y$. By construction of $\text{dom} f_\omega$, there is $z \in tA$ with $w \leq z$ and $y \leq z$. Since $\text{im} f_\omega$ is both a filter and an ideal, it follows that $w \in \text{im} f_\omega$. A similar argument shows that if x and y are as before and $w \in \text{im} f_\omega$, then $w \in \text{dom} f_\omega$.

It follows that either f_ω or $f_\omega \cup f_\omega^{-1}$ is an automorphism of a subtree X

such that if $t \in T - X$, then t is unrelated to any element of X . Thus that automorphism can be safely extended to all of T by stipulating that it be the identity on $T - X$. By construction it takes t to u as required.

Finally, to prove the converse half of Theorem 1(a), the preceding construction can be modified easily as follows: At the first stage, take $g_x: xB \rightarrow xf_0B$ to be any function with $xg_x = xf_0$. At each later stage, g_x can be any function from xC to xf_iC . Conditions (3) and (4) will still be true at each stage, and condition (5) will be true if the word "isomorphism" is replaced by the phrase "local isomorphism". The function f thus constructed will be a local isomorphism from its domain to its image. This function (rather than $f_\omega \cup f_\omega^{-1}$ as in the automorphism case) may then be extended to all of T by taking it to be the identity function outside the domain of f_ω . The resulting function will then be a local isomorphism taking t to u .

3. Proof of Theorem 2. The following two facts are needed for the proof of the forward part of Theorem 2. Both follow easily from (2). In both, S is a semigroup.

LEMMA 4. *Let X be an S -set with endomorphism α . Then α is surjective if and only if $\text{Im } \alpha$ generates X .*

LEMMA 5. *Let X be a transitive S -set with endomorphism α . Then α has a fixed point if and only if $\alpha = \text{id}_X$.*

Now assume that X is a transitive S -set with endomorphism α . Then α is surjective by Lemma 4, because every element of X generates X . Furthermore, Lemma 5 applies equally well to α^k for any k . But no power of a noninjective function can be the identity function. This proves the forward part of Theorem 2.

The converse part of Theorem 2 follows from Theorem 1. Suppose that $\alpha: X \rightarrow X$ is a surjective function with the property that no power of α has a fixed point. Define a relation \cong on X this way:

(6) $x \cong y$ if $y = x\alpha^k$ for some nonnegative integer k .

It is easy to see that \cong is reflexive and transitive. It is also antisymmetric: If $y = x\alpha^k$ and $x = y\alpha^m$ for some nonnegative integers k and m , then y is a fixed point of α^{k+m} , which means that $k + m = 0$, hence $k = m = 0$, hence $x = y$. Thus (X, \cong) is a poset. Furthermore, if $x \in X$, then xA is obviously isomorphic to the set ω , so that (X, \cong) is a locally finite tree with no maximal elements by Lemma 1.

It is immediate that α is the parent mapping of the tree (X, \cong) . Since α is surjective, (X, \cong) has no minimal elements, so by Theorem 1, $LI(T)$ is transitive. But by Lemma 3, every function in $LI(T)$ commutes with α , as required.

The last sentence of Theorem 2 clearly follows from Theorem 1(b).

4. Remarks. A congruence on an S -set X is a partition π of X with the property that if $u \in \pi$ and $u, v \in U$, then us and vs are in the same block of π for any $s \in S$. If X has an S -endomorphism α , the set $\{x\alpha^{-1} \mid x \in X\}$ is a congruence on X (this follows easily from (2)). It follows that if X is a transitive S -set with no nontrivial congruences, then α must be a permutation of X . In particular, a primitive group action, finite or infinite, is centralized only by permutations.

It is easy to see, using (2), that a doubly transitive semigroup action is centralized only by the identity function.

The way to construct permutations that centralize finite transitive group actions has been known since the turn of the century. A recent exposition is in Kerber [2]. Expositions of the elementary theory of semigroup actions may be found in Banaschewski [1], Knauer [3], Petrich [5] and Wells [8] with varying emphases. This subject is treated from an axiomatic point of view by Šaĭn [6].

Infinite trees have been used as tools in various areas, for example in mathematical linguistics. References may be found in Maxson [4]. Serre [7] describes the structure of groups acting on trees. (Warning: My terminology is quite different from his!).

It would be interesting to characterize all the trees with transitive automorphism semigroup. Such a tree need not be locally finite; the only examples I know, however, are all quite complicated. The one which can be described most succinctly is the following:

Let T consist of all ordered triples (a, b, γ) with $a, b \in Z$ (Z the set of all integers) and $\gamma: Z \rightarrow Z$ a function for which

$$(7) \quad n\gamma = 0 \quad \text{for} \quad n < b.$$

Define

$$(8) \quad (a, b, \gamma) \cong (a', b', \gamma') \Leftrightarrow \begin{cases} a \cong a', b = b' \text{ and } \gamma = \gamma' \\ \text{or } b < b', (b' - 1)\gamma \cong a' \text{ and} \\ n\gamma = n\gamma' \text{ for all } n \cong b'. \end{cases}$$

A tedious but straightforward proof shows that T is a nonlocally finite tree. The element (a, b, γ) may be visualized as the integer a in a copy of Z which is hanging down from the element $(b\gamma, b + 1, \gamma')$, where γ' agrees with γ except that $b\gamma' = 0$. Each element of T has a copy of Z hanging below it.

If (a, b, γ) and (a', b', γ') are elements of T , the following defines an automorphism φ of T which takes (a, b, γ) to (a', b', γ') .

Let $\eta: T \rightarrow T$ be defined by

$$(9) \quad y\eta = \begin{cases} (y + b' - b - 1)\gamma' - (y - 1)\gamma & \text{for } y > b. \\ a' - a & \text{for } y \leq b. \end{cases}$$

If $(x, y, \xi) \in T$, define $\bar{\xi}: Z \rightarrow Z$ by

$$(10) \quad n\bar{\xi} = \begin{cases} (n + b - b')\xi + n\gamma' - (n + b - b')\gamma & \text{for } n \geq y + b' - b \\ \text{and } n \geq b'. \\ (n + b - b')\xi + a' - a & \text{for } n \geq y + b' - b \text{ and } n < b'. \\ 0 & \text{for } n < y + b' - b. \end{cases}$$

Now set, for $(x, y, \xi) \in T$,

$$(11) \quad (x, y, \xi)\varphi = (x + y\eta, y + b' - b, \bar{\xi}).$$

Then φ will be an automorphism of T taking (a, b, γ) to (a', b', γ') .

The author wishes to thank Steve McCleary for pointing out an error in the original version of Theorem 1, and Paul Kainen, who came up with the idea behind the preceding example.

REFERENCES

1. B. Banaschewski, *Functions into categories of M-sets*, Abh. Math. Sem. Univ. Hamburg, **38** (1972), 49–64.
2. A. Kerber, *Representations of permutation groups I*, Lecture Notes in Mathematics 240. Springer Verlag, 1971.
3. U. Knauer, *Projectivity of acts and Morita equivalence of monoids*, Semigroup Forum, **3** (1971/72), 359–370.
4. C. Maxson, *Semigroups of order-preserving partial endomorphisms on trees, I*, To be published.
5. M. Petrich, *Dense extensions of semigroups I*, J. reine ang. Math., **258** (1973), 103–125, section 2.
6. B. M. Šain, *Transformative semigroups of transformations*, Mat. Sbornik (N.S.) **71** (113), (1968), 65–82.
7. J. P. Serre, *Arbes, Amalgams et SL_2* , Mimeographed notes.
8. C. Wells, *Some applications of the wreath product construction*, Amer. Math. Monthly **83**, May, 1976.

Received March 18, 1975.

EIDGEN TECHNISCHE HOCHSCHULE
AND
CASE WESTERN RESERVE UNIVERSITY

Pacific Journal of Mathematics

Vol. 64, No. 1

May, 1976

Walter Allegretto, <i>Nonoscillation theory of elliptic equations of order $2n$</i>	1
Bruce Allem Anderson, <i>Sequencings and starters</i>	17
Friedrich-Wilhelm Bauer, <i>A shape theory with singular homology</i>	25
John Kelly Beem, <i>Characterizing Finsler spaces which are pseudo-Riemannian of constant curvature</i>	67
Dennis K. Burke and Ernest A. Michael, <i>On certain point-countable covers</i>	79
Robert Chen, <i>A generalization of a theorem of Chacon</i>	93
Francis H. Clarke, <i>On the inverse function theorem</i>	97
James Bryan Collier, <i>The dual of a space with the Radon-Nikodým property</i>	103
John E. Cruthirds, <i>Infinite Galois theory for commutative rings</i>	107
Artatrana Dash, <i>Joint essential spectra</i>	119
Robert M. DeVos, <i>Subsequences and rearrangements of sequences in FK spaces</i>	129
Geoffrey Fox and Pedro Morales, <i>Non-Hausdorff multifunction generalization of the Kelley-Morse Ascoli theorem</i>	137
Richard Joseph Fleming, Jerome A. Goldstein and James E. Jamison, <i>One parameter groups of isometries on certain Banach spaces</i>	145
Robert David Gulliver, II, <i>Finiteness of the ramified set for branched immersions of surfaces</i>	153
Kenneth Hardy and István Juhász, <i>Normality and the weak cb property</i>	167
C. A. Hayes, <i>Derivation of the integrals of $L^{(q)}$-functions</i>	173
Frederic Timothy Howard, <i>Roots of the Euler polynomials</i>	181
Robert Edward Jamison, II, Richard O'Brien and Peter Drummond Taylor, <i>On embedding a compact convex set into a locally convex topological vector space</i>	193
Andrew Lelek, <i>An example of a simple triod with surjective span smaller than span</i>	207
Janet E. Mills, <i>Certain congruences on orthodox semigroups</i>	217
Donald J. Newman and A. R. Reddy, <i>Rational approximation of e^{-x} on the positive real axis</i>	227
John Robert Quine, Jr., <i>Homotopies and intersection sequences</i>	233
Nambury Sitarama Raju, <i>Periodic Jacobi-Perron algorithms and fundamental units</i>	241
Herbert Silverman, <i>Convexity theorems for subclasses of univalent functions</i>	253
Charles Frederick Wells, <i>Centralizers of transitive semigroup actions and endomorphisms of trees</i>	265
Volker Wrobel, <i>Spectral approximation theorems in locally convex spaces</i>	273
Hidenobu Yoshida, <i>On value distribution of functions meromorphic in the whole plane</i>	283