

Pacific Journal of Mathematics

QUASITRIANGULAR OPERATOR ALGEBRAS

JOAN KATHRYN PLASTIRAS

QUASITRIANGULAR OPERATOR ALGEBRAS

JOAN K. PLASTIRAS

Fix a sequence $\mathcal{P} = \{P_n\}_{n=1}^\infty$ of finite dimensional projections increasing to the identity on a separable Hilbert space \mathcal{H} and let $\mathcal{L}(\mathcal{H})$ denote the algebra of all bounded operators on \mathcal{H} . The quasitriangular algebra associated with \mathcal{P} and denoted as $\mathcal{QT}(\mathcal{P})$ is defined to be the set of those operators T in $\mathcal{L}(\mathcal{H})$ for which $\|P_n^\perp TP_n\| \rightarrow 0$.

In this paper we will examine the structure of the $\mathcal{QT}(\mathcal{P})$ algebras. Specifically, if $\mathcal{R} = \{R_n\}_{n=1}^\infty$ is another sequence of finite dimensional projections increasing to the identity on the same Hilbert space, when is $\mathcal{QT}(\mathcal{R})$ equal to $\mathcal{QT}(\mathcal{P})$? By an algebraic isomorphism between two algebras we shall mean a bijection which preserves algebraic structure: that is to say — addition, scalar multiplication, multiplication, but we do not impose any topological condition. When are two quasitriangular algebras isomorphic?

In [5] we asked the same questions of $\mathcal{D}(\mathcal{E}) + \mathcal{C}(\mathcal{H}) = \{T + K : T \text{ belongs to the commutant of } E \text{ and } K \text{ is compact}\}$ and answered them completely by arguments very different from those presented here; the conclusions were different too. The concept of quasitriangularity for operators was first isolated for systematic study in [3]. The quasitriangular algebra was introduced later in [1] and a formula expressing the distance from such an algebra to an arbitrary operator was obtained. We begin our discussion with an algebraic property:

DEFINITION 1. A subset \mathcal{S} of $\mathcal{L}(\mathcal{H})$ is said to be inverse-closed if whenever T in \mathcal{S} is invertible in $\mathcal{L}(\mathcal{H})$ then T^{-1} belongs to \mathcal{S} .

LEMMA 2. $\mathcal{QT}(\mathcal{P})$ is inverse-closed for every sequence $\mathcal{P} = \{P_n\}_{n=1}^\infty$ of finite dimensional projections increasing to the identity on a Hilbert space.

Before verifying Lemma 2 we remark that the assumption that the P_k be finite dimensional is essential.

Proof. From [1, Corollary following 2.2] we know that $\mathcal{QT}(\mathcal{P}) = \mathcal{T}(\mathcal{P}) + \mathcal{C}(\mathcal{H})$, where $\mathcal{T}(\mathcal{P})$ is the set of operators T such that $P_n^\perp TP_n = 0$ for all n . Hence, it suffices to assume that S belongs to $\mathcal{T}(\mathcal{P}) + \mathcal{C}(\mathcal{H})$ and is invertible in $\mathcal{L}(\mathcal{H})$ and show that S^{-1} belongs to $\mathcal{QT}(\mathcal{P})$. So, $S = T + C$, where $T \in \mathcal{T}(\mathcal{P})$ and $C \in \mathcal{C}(\mathcal{H})$. Since $S_m = T + P_m CP_m$ tends in norm to S , S_m is invertible for all m greater than a positive

integer l . Fix $m > l$ and note that $S_m P_n = P_n S_m P_n$ for all $n \geq m$, and since $\dim P_n < \infty$, S_m maps $P_n \mathcal{H}$ onto itself, so that $P_n S_m^{-1} P_n = S_m^{-1} P_n$ (or equivalently, $P_n^{\perp} S_m^{-1} P_n \equiv 0$). Hence, S_m^{-1} belongs to $\mathcal{QT}(\mathcal{P})$ by definition. As S_m^{-1} tends in norm to $(T + C)^{-1}$ and $\mathcal{QT}(\mathcal{P})$ is norm-closed [1, Proposition 2.1], we conclude that $(T + C)^{-1}$ belongs to $\mathcal{QT}(\mathcal{P})$.

THEOREM 3. *Suppose that T is an invertible operator in $\mathcal{L}(\mathcal{H})$. Then T implements an automorphism of $\mathcal{QT}(\mathcal{P})$ (i.e. $T\mathcal{QT}(\mathcal{P})T^{-1} = \mathcal{QT}(\mathcal{P})$) if and only if T belongs to $\mathcal{QT}(\mathcal{P})$.*

Proof. \Leftarrow : Assume that T belongs to $\mathcal{QT}(\mathcal{P})$. To show that T implements an inner automorphism of $\mathcal{QT}(\mathcal{P})$ it will suffice to show that T^{-1} also belongs to $\mathcal{QT}(\mathcal{P})$. But that is immediate from Lemma 2.

\Rightarrow : Assume that T implements an automorphism of $\mathcal{QT}(\mathcal{P})$. First we conclude from [1, Theorem 3.3] that T admits a factorization $T = UA$, where A belongs to $\mathcal{T}(\mathcal{P})$ and U is a partial isometry. Note that $A = U^*T$ has closed range; since $\ker A = \{0\}$, A is semi-Fredholm by definition. Since A belongs to $\mathcal{QT}(\mathcal{P})$ the index of A is nonnegative [2] so that $\ker A^* = \{0\}$ and A is consequently invertible. This forces U to be unitary. Since $A \in \mathcal{QT}(\mathcal{P})$ is invertible, then by the previous argument, A implements an automorphism of $\mathcal{QT}(\mathcal{P})$ so that we are reduced to showing that if U is a unitary operator which implements an automorphism of $\mathcal{QT}(\mathcal{P})$, then U belongs to $\mathcal{QT}(\mathcal{P})$.

So, we assume that U does not belong to $\mathcal{QT}(\mathcal{P})$ and arrive at a contradiction. Since U does not belong to $\mathcal{QT}(\mathcal{P})$ then by the definition of $\mathcal{QT}(\mathcal{P})$ there is an $\alpha > 0$ and a subsequence $\{P_{n(k)}\}_{k=1}^{\infty}$ of \mathcal{P} for which $\liminf_k \|P_{n(k)}^{\perp} U P_{n(k)}\| \geq \alpha$. From Lemma 2 we know that U^* does not belong to $\mathcal{QT}(\{P_{n(k)}\}_{n=1}^{\infty})$, so that by definition, there is $\beta > 0$ and a subsequence $\{m(k)\}_{k=1}^{\infty}$ of $\{n(k)\}_{k=1}^{\infty}$ for which $\liminf_k \|P_{m(k)}^{\perp} U^* P_{m(k)}\| \geq \beta$. If we let $\epsilon = \min(\alpha, \beta)/2$, then we can conclude that $\|P_n^{\perp} U P_n\|$ and $\|P_n U P_n^{\perp}\|$ ($= \|P_n^{\perp} U^* P_n\|$) are both greater than ϵ for all n in an infinite subset M of \mathbf{N} .

We will obtain a sequence $\{m_i, n_i\}_{i=1}^{\infty}$ of positive integers such that $0 < m_1 < n_1 < m_2 < n_2 < \dots$ and projections $\{F_k, E_k\}_{k=1}^{\infty}$ such that $F_k = P_{m_k} P_{n_{k-1}}^{\perp}$ and $E_k = P_{n_k} P_{m_k}^{\perp}$ for which $\|F_k U E_k\|$ and $\|E_k U F_k\|$ are both greater than $\epsilon/2$. We do so inductively.

For $k = 1$, define $F_1 = P_{m_1}$, where m_1 is the first integer in M . Let n_1 be the first integer such that $\|P_{n_1} P_{m_1}^{\perp} U P_{m_1}\|$ and $\|P_{m_1} U P_{n_1}^{\perp} P_{n_1}\|$ are both greater than $\epsilon/2$ (such an n_1 exists because $\|P_{m_1}^{\perp} U P_{m_1}\|$ and $\|P_{m_1} U P_{m_1}^{\perp}\|$ are greater than ϵ and the P_n tend strongly to the identity).

Assume that we have obtained $\{E_k, F_k\}_{k=1}^l$. To obtain m_{l+1} and n_{l+1} , note that $U P_{n_l}$ and $P_{n_l} U$ are compact; hence, there is a positive integer j such that $\|P_n^{\perp} U P_n\|$ and $\|P_n U P_n^{\perp}\|$ are both less than $\epsilon/4$ for all $n \geq j$. Let m_{l+1} be the first integer in M greater than j .

Then

$$\begin{aligned} \|P_{m_{l+1}}^\perp UP_{m_{l+1}} P_{n_l}^\perp\| &\geq \|P_{m_{l+1}}^\perp UP_{m_{l+1}}\| - \|P_{m_{l+1}}^\perp UP_{m_{l+1}} P_{n_l}\| \\ &\geq \epsilon - \|P_{m_{l+1}}^\perp UP_{n_l}\| \\ &\geq \epsilon - \epsilon/4 = \frac{3}{4}\epsilon. \end{aligned}$$

Similarly, $\|P_{m_{l+1}} P_{n_l}^\perp UP_{m_{l+1}}^\perp\| \geq 3\epsilon/4$ by the same argument. Let n_{l+1} be the first positive integer greater than m_{l+1} for which $\|P_{n_{l+1}} P_{m_{l+1}}^\perp UP_{m_{l+1}}^\perp P_{n_l}^\perp\|$ and $\|P_{m_{l+1}}^\perp P_{n_l}^\perp UP_{m_{l+1}}^\perp P_{n_{l+1}}\|$ are both greater than $\epsilon/2$. Let $F_{l+1} = P_{m_{l+1}} P_{n_l}^\perp$ and let $E_{l+1} = P_{n_{l+1}} P_{m_{l+1}}^\perp$. Continue inductively.

We select a subsequence $\{E_{i_j}, F_{i_j}\}_{j=1}^\infty$ of $\{E_i, F_i\}_{i=1}^\infty$ as follows: first, we let $\{\alpha_{ij}\}_{i,j=1}^\infty$ be any sequence of positive real numbers such that $\sum_{i,j} \alpha_{ij}^2 \leq \epsilon^2/16$. Let $i_1 = 1$. Assuming that we have obtained i_k , let i_{k+1} be the next positive integer such that for all $l \not\leq k + 1$, $\|E_{i_{k+1}} UF_{i_l}\|$ and $\|F_{i_{k+1}} UE_{i_l}\|$ are less than $\alpha_{k+1,l}$ while $\|E_{i_l} UF_{i_{k+1}}\|$ and $\|F_{i_l} UE_{i_{k+1}}\|$ are less than $\alpha_{l,k+1}$. This is possible because $UF_{i_l}, F_{i_l}U$ (respectively $UE_{i_l}, E_{i_l}U$) are compact and the E_i (respectively F_i) tend weakly to zero. Continue inductively. Now for each i_k there is a rank one partial isometry $T_{i_k} \in \mathcal{L}(E_{i_k}\mathcal{H}, F_{i_k}\mathcal{H})$ such that $\|E_{i_k} UT_{i_k} U^* F_{i_k}\| \geq \epsilon^2/4$. Clearly, $T = \sum_{k=1}^\infty T_{i_k}$ is a partial isometry in $\mathcal{T}(\mathcal{P})$. So, for arbitrary l in \mathbf{N} ,

$$E_{i_l}(UTU^*)F_{i_l} = \sum_{k=1}^\infty E_{i_l} UT_{i_k} U^* F_{i_l} = E_{i_l} UT_{i_l} U^* F_{i_l} + \sum_{\substack{k=1 \\ k \neq l}}^\infty E_{i_l} UT_{i_k} U^* F_{i_l}.$$

Hence,

$$\begin{aligned} \|E_{i_l}(UTU^*)F_{i_l}\| + \left\| \sum_{\substack{k=1 \\ k \neq l}}^\infty E_{i_l} UT_{i_k} U^* F_{i_l} \right\| &\geq \|E_{i_l} UT_{i_l} U^* F_{i_l}\| \\ \|E_{i_l}(UTU^*)F_{i_l}\| + \sum_{\substack{k=1 \\ k \neq l}}^\infty \|E_{i_l} UT_{i_k} U^* F_{i_l}\| &\geq (\epsilon/2)^2 = \epsilon^2/4. \end{aligned}$$

Therefore,

$$\begin{aligned} \|E_{i_l}(UTU^*)F_{i_l}\| &\geq \frac{\epsilon^2}{4} - \sum_{k \neq l} \|E_{i_l} UF_{i_k}\| \cdot \|E_{i_k} U^* F_{i_l}\| \\ &\geq \frac{3\epsilon^2}{16}. \end{aligned}$$

Since i_l was arbitrary, it follows from the construction that

$$\frac{3\epsilon^2}{16} \leq \|E_{i_l}(UTU^*)F_{i_l}\| \leq \|P_{m_{i_l}}^\perp (UTU^*) P_{m_{i_l}}\|.$$

Hence,

$$\overline{\lim}_k \| P_k^\perp (UTU^*) P_k \| > 0$$

and it follows by definition of $\mathcal{QT}(\mathcal{P})$ that UTU^* does not belong to $\mathcal{QT}(\mathcal{P})$. This contradicts our assumption that U implements an automorphism of $\mathcal{QT}(\mathcal{P})$ and thus concludes the argument of the proof of Theorem 3.

DEFINITION 4. Let $\mathcal{P} = \{P_n\}_{n=1}^\infty$ be a sequence of finite dimensional projections increasing to the identity on a Hilbert space \mathcal{H} . An operator T is said to be *strictly upper triangular for \mathcal{P}* if $P_n^\perp T P_{n+1} = 0$ for all n in \mathbf{N} .

REMARK 5. Note that in the proof of Theorem 3 we showed that if U does not belong to $\mathcal{QT}(\mathcal{P})$ then there is an operator T , which is strictly upper triangular for \mathcal{P} , and such that UTU^* does not belong to $\mathcal{QT}(\mathcal{P})$.

REMARK 6. Let $\mathcal{S} = \{S_n\}_{n=1}^\infty$ be any sequence of finite dimensional projections increasing to the identity on \mathcal{H} . Let $\mathcal{P} = \{P_n\}_{n=1}^\infty$ be a subsequence of \mathcal{S} . Then $\mathcal{QT}(\mathcal{S}) \subseteq \mathcal{QT}(\mathcal{P})$. Equality may fail; however, if T is strictly upper triangular for \mathcal{P} then T belongs to $\mathcal{QT}(\mathcal{S})$.

DEFINITION 7. A sequence of projections $\mathcal{S} = \{S_n\}_{n=1}^\infty$ increasing to the identity on a Hilbert space \mathcal{H} is said to be a *defining sequence* for a quasitriangular algebra \mathcal{A} if and only if $\mathcal{A} = \{T \in \mathcal{L}(\mathcal{H}) : \|S_n^\perp T S_n\| \rightarrow 0\}$.

REMARK 8. Suppose that U is a unitary operator which implements an isomorphism $T \rightarrow UTU^*$ from $\mathcal{QT}(\mathcal{P})$ onto $\mathcal{QT}(\mathcal{S})$. Then U maps defining sequences of $\mathcal{QT}(\mathcal{P})$ to defining sequences of $\mathcal{QT}(\mathcal{S})$.

LEMMA 9. *Suppose that $\mathcal{P} = \{P_n\}_{n=1}^\infty$ and $\mathcal{S} = \{S_n\}_{n=1}^\infty$ are sequences of finite dimensional projections increasing to the identity such that $\mathcal{P} \cup \mathcal{S}$ is totally ordered by inclusion. Then $\mathcal{QT}(\mathcal{P}) = \mathcal{QT}(\mathcal{S})$ if and only if there exist positive integers m_0 and n_0 such that $P_{m_0+k} = S_{n_0+k}$ for all k in \mathbf{N} .*

Proof. \Leftarrow : This conclusion is clear.

\Rightarrow : Assume that $\mathcal{QT}(\mathcal{P}) = \mathcal{QT}(\mathcal{S})$. Then $\mathcal{QT}(\mathcal{P}) = \mathcal{QT}(\mathcal{P} \cup \mathcal{S})$. We assert that \mathcal{P} contains all but perhaps finitely many of the projections in $\mathcal{P} \cup \mathcal{S}$. Contrapositively, assume not. Let $\mathcal{R} = \{R_n\}_{n=1}^\infty$ be a total ordering of $\mathcal{P} \cup \mathcal{S}$ and choose an infinite subsequence $\{n_k\}_{k=1}^\infty$ for which $R_{n_k} \notin \mathcal{P}$ but $R_{n_k+1} \in \mathcal{P}$. Let T_k be any rank one partial isometry with initial space $(R_{n_k} \ominus R_{n_k-1})\mathcal{H}$ and final space $(R_{n_k+1} \ominus R_{n_k})\mathcal{H}$. Then $T = \sum_{k=1}^\infty T_k$ is a partial isometry which belongs to $\mathcal{QT}(\mathcal{P})$ but not to $\mathcal{QT}(\mathcal{P} \cup \mathcal{S})$.

Hence, $\mathcal{QT}(\mathcal{P} \cup \mathcal{S}) \not\subseteq \mathcal{QT}(\mathcal{P})$. We conclude that \mathcal{P} contains all but perhaps finitely many of the projections in $\mathcal{P} \cup \mathcal{S}$.

By symmetry, \mathcal{S} contains all but perhaps finitely many of the projections in $\mathcal{P} \cup \mathcal{S}$. So there exists a positive integer k such that $\{P_n : \dim P_n \geq k\} \subseteq \mathcal{S}$ and $\{S_n : \dim S_n \geq k\} \subseteq \mathcal{P}$. Let m_0 be the first positive integer such that $\dim(P_{m_0}) \geq k$ and let n_0 be the first integer such that $\dim(S_{n_0}) \geq k$. Then $P_{m_0+k} = S_{n_0+k}$ for all $k \in \mathbb{N}$.

THEOREM 10. $\mathcal{S} = \{S_n\}_{n=1}^\infty$ is a defining sequence for $\mathcal{QT}(\mathcal{P})$ if and only if there exist positive integers m_0 and n_0 such that $\lim_k \|P_{m_0+k} - S_{n_0+k}\| = 0$.

Proof. \Leftarrow : We note that $\mathcal{QT}(\mathcal{S}) \subseteq \mathcal{QT}(\mathcal{P})$ since for T in $\mathcal{QT}(\mathcal{S})$,

$$\begin{aligned} \|P_{m_0+k}^\perp T P_{m_0+k}\| &\leq \|S_{n_0+k}^\perp T S_{n_0+k}\| + \|(P_{m_0+k}^\perp - S_{n_0+k}^\perp) T S_{n_0+k}\| \\ &\quad + \|P_{m_0+k}^\perp T (P_{m_0+k} - S_{n_0+k})\| \\ &\leq \|S_{n_0+k}^\perp T S_{n_0+k}\| + \|P_{m_0+k}^\perp - S_{n_0+k}^\perp\| \cdot \|T\| \\ &\quad + \|T\| \cdot \|P_{m_0+k} - S_{n_0+k}\|, \end{aligned}$$

and the other inclusion follows by symmetry.

\Rightarrow : We assume that $\mathcal{S} = \{S_n\}_{n=1}^\infty$ is a defining sequence for $\mathcal{QT}(\mathcal{P})$. Let V be any unitary operator such that $\{VS_n V^*\}_{n=1}^\infty \cup \{P_n\}_{n=1}^\infty$ is a sequence of projections totally ordered by set inclusion.

Let $\mathcal{W} = \{W_n\}_{n=1}^\infty$ with $W_n = VS_n V^*$ for each n . We assert that V belongs to $\mathcal{QT}(\mathcal{W})$. So assume that T is strictly upper triangular for \mathcal{W} ; it suffices to show that VTV^* belongs to $\mathcal{QT}(\mathcal{W})$ by Remark 5. By Remark 6, T belongs to $\mathcal{QT}(\mathcal{P} \cup \mathcal{W}) \subseteq \mathcal{QT}(\mathcal{P})$ so that it remains to observe that $V\mathcal{QT}(\mathcal{P})V^* \subseteq \mathcal{QT}(\mathcal{W})$: $W_n^\perp (VTV^*) W_n = (VS_n^\perp V^*) (VTV^*) (VS_n V^*) = VS_n^\perp T S_n V^*$, so that $\|W_n^\perp (VTV^*) W_n\| = \|VS_n^\perp T S_n V^*\| = \|S_n^\perp T S_n\| \rightarrow 0$.

Hence, we conclude that V belongs to $\mathcal{QT}(\mathcal{W})$. Since $\mathcal{QT}(\mathcal{W})$ is inverse-closed by Lemma 2, it follows that $\|W_n^\perp V W_n\| \rightarrow 0$ and $\|W_n V W_n^\perp\| = \|W_n^\perp V^* W_n\| \rightarrow 0$.

(1) Since $W_n V = VS_n$, we have that $W_n V W_n^\perp = VS_n W_n^\perp$ so that $\|W_n V W_n^\perp\| = \|VS_n W_n^\perp\| = \|S_n W_n^\perp\| \rightarrow 0$ and

(2) Since $W_n^\perp V = VS_n^\perp$, we have that $W_n^\perp V W_n = VS_n^\perp W_n$ so that $\|W_n^\perp V W_n\| = \|VS_n^\perp W_n\| = \|S_n^\perp W_n\| \rightarrow 0$.

Since $\|S_n - W_n\| = \max\{\|S_n^\perp W_n\|, \|S_n W_n^\perp\|\}$ [5, Lemma 6] it follows that $\lim_n \|S_n - W_n\| = 0$ and by a previous argument that \mathcal{W} is a defining sequence for $\mathcal{QT}(\mathcal{P})$. It follows from Lemma 9 that there are integers m_0 and n_0 such that $W_{m_0+k} = P_{m_0+k}$ for all k in \mathbb{N} . Hence

$$\lim_k \|S_{m_0+k} - P_{m_0+k}\| = 0,$$

which concludes the proof.

EXAMPLE 11. As an easy consequence of Theorem 10, it follows that there exist defining sequences $\mathcal{P} = \{P_n\}_{n=1}^\infty$ and $\mathcal{R} = \{R_n\}_{n=1}^\infty$ for a quasitriangular algebra \mathcal{A} such that $\{P_n \vee R_n\}_{n=1}^\infty$ is not a defining sequence for \mathcal{A} (“ \vee ” denotes the supremum of two projections). This phenomenon is suggested by an example in [3, p. 285].

We shall say that two subsets of $\mathcal{L}(\mathcal{H})$, \mathcal{S} and \mathcal{T} , are *locally isomorphic* if each operator in \mathcal{S} is unitarily equivalent to an operator in \mathcal{T} and conversely. Because every quasitriangular operator is a compact perturbation of a triangular operator, it follows that any two quasitriangular algebras are locally isomorphic; from Theorem 12 we conclude that they are not necessarily isomorphic.

THEOREM 12. *Let $\mathcal{QT}(\mathcal{P})$ and $\mathcal{QT}(\mathcal{S})$ be quasitriangular algebras. Then $\mathcal{QT}(\mathcal{P})$ and $\mathcal{QT}(\mathcal{S})$ are algebraically isomorphic if and only if there exist positive integers j_0 and l_0 such that $\dim(P_{j_0+k}) = \dim(S_{l_0+k})$ for all k in \mathbf{N} .*

Proof. \Leftarrow : If we assume that there exist positive integers j_0 and l_0 such that $\dim(P_{j_0+k}) = \dim(S_{l_0+k})$ for all k in \mathbf{N} , then we can define a unitary operator U such that $UP_{j_0+k}U^* = S_{l_0+k}$ for all k in \mathbf{N} . We assert that U implements an isomorphism from $\mathcal{QT}(\mathcal{P})$ to $\mathcal{QT}(\mathcal{S})$.

\Rightarrow : Assume that there is a map α from $\mathcal{QT}(\mathcal{P})$ to $\mathcal{QT}(\mathcal{S})$ which preserves algebraic structure. Since $\mathcal{QT}(\mathcal{P})$ and $\mathcal{QT}(\mathcal{S})$ are Banach algebras, each of which contains the set of finite rank operators, it follows from [6, Theorem 2.5.19] that there exists an invertible operator S such that $\alpha(T) = STS^{-1}$ for all T in $\mathcal{QT}(\mathcal{P})$.

We conclude from [1, Theorem 3.3] that S has a factorization $S = UA$ where A belongs to $\mathcal{T}(\mathcal{P})$ and U is unitary. Then we note that $R_n = UP_nU^*$ is a defining sequence for $\mathcal{QT}(\mathcal{S})$; by Theorem 10, we note that there exist positive integers m_0 and n_0 such that $\|R_{m_0+k} - S_{n_0+k}\| \rightarrow 0$. So, there exists a positive integer d such that $\|R_{m_0+d+k} - S_{n_0+d+k}\| < 1$ for all k in \mathbf{N} . Hence, $\dim(R_{m_0+d+k}) = \dim(S_{n_0+d+k})$ for all k in \mathbf{N} . Since $\dim(P_n) = \dim(R_n)$ for all n in \mathbf{N} , let $j_0 = m_0 + d$ and let $l_0 = n_0 + d$ to obtain the theorem.

REFERENCES

1. W. B. Arveson, *Interpolation problems in nest algebras*, J. Functional Analysis, **20** (1975), 208–233.

2. R. G. Douglas and C. Pearcy, *A note on quasitriangular operators*, Duke Math. J., **37** (1970), 177–188.
3. P. R. Halmos, *Quasitriangular operators*, Acta Sci. Mat. (Szeged), **29** (1968), 283–293.
4. ———, *Ten problems in Hilbert space*, Bulletin Amer. Math. Soc., **76** (1970), 887–933.
5. J. K. Plastiras, *Compact perturbations of certain von Neumann algebras*, Trans. Amer. Math. Soc., (to appear).
6. C. E. Rickart, *General Theory of Banach Algebras*, D. van Nostrand Company, Inc. (1960).
7. B. Sz.-Nagy, *Spektraldarstellung linearer Transformation des Hilbertschen Raumes*, Berlin, (1942).

Received February 23, 1976.

UNIVERSITY OF PENNSYLVANIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed). double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in duplicate, may be sent to any one of the four editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Printed at Jerusalem Academic Press, POB 2390, Jerusalem, Israel.

Copyright © 1976 Pacific Journal of Mathematics
All Rights Reserved

Pacific Journal of Mathematics

Vol. 64, No. 2

June, 1976

Richard Fairbanks Arnold and A. P. Morse, <i>Plus and times</i>	297
Edwin Ogilvie Buchman and F. A. Valentine, <i>External visibility</i>	333
R. A. Czerwinski, <i>Bonded quadratic division algebras</i>	341
William Richard Emerson, <i>Averaging strongly subadditive set functions in unimodular amenable groups. II</i>	353
Lynn Harry Erbe, <i>Existence of oscillatory solutions and asymptotic behavior for a class of third order linear differential equations</i>	369
Kenneth R. Goodearl, <i>Power-cancellation of groups and modules</i>	387
J. C. Hankins and Roy Martin Rakestraw, <i>The extremal structure of locally compact convex sets</i>	413
Burrell Washington Helton, <i>The solution of a Stieltjes-Volterra integral equation for rings</i>	419
Frank Kwang-Ming Hwang and Shen Lin, <i>Construction of 2-balanced (n, k, λ) arrays</i>	437
Wei-Eihn Kuan, <i>Some results on normality of a graded ring</i>	455
Dieter Landers and Lothar Rogge, <i>Relations between convergence of series and convergence of sequences</i>	465
Lawrence Louis Larmore and Robert David Rigdon, <i>Enumerating immersions and embeddings of projective spaces</i>	471
Douglas C. McMahon, <i>On the role of an abelian phase group in relativized problems in topological dynamics</i>	493
Robert Wilmer Miller, <i>Finitely generated projective modules and TTF classes</i>	505
Yashaswini Deval Mittal, <i>A class of isotropic covariance functions</i>	517
Anthony G. Mucci, <i>Another martingale convergence theorem</i>	539
Joan Kathryn Plastiras, <i>Quasitriangular operator algebras</i>	543
John Robert Quine, Jr., <i>The geometry of $p(S^1)$</i>	551
Tsuan Wu Ting, <i>The unloading problem for severely twisted bars</i>	559