

Pacific Journal of Mathematics

CHAINED RINGS

PAUL FROESCHL

CHAINED RINGS

PAUL A. FROESCHL III

All rings considered are commutative with identity. A chained ring is any ring whose set of ideals is totally ordered by inclusion. The main object of this paper is stating conditions in which every valuation overring of a given ring is a chained ring. It is shown that every valuation overring of a ring R is a chained ring if and only if the ideal of zero divisors of $T(R)$, the total quotient ring of R , is the conductor of \bar{R} , the integral closure of R , in $T(R)$. An example is provided of a valuation ring which is not a chained ring even though its total quotient ring is a chained ring.

1. Preliminaries. In this short section we set notation and establish some elementary results.

A *total quotient ring* is a ring consisting entirely of zero divisors and units. If R is a ring, then let $T(R)$ denote the smallest total quotient ring containing R . We will call $T(R)$ *the total quotient ring of R* .

Let G be a totally ordered abelian group. A *valuation* on a ring T is a function v from T onto $G \cup \{\infty\}$, such that for all x and y in T :

- (1) $v(xy) = v(x) + v(y)$;
- (2) $v(x + y) \geq \min\{v(x), v(y)\}$;
- (3) $v(1) = 0$ and $v(0) = \infty$.

Let V be a ring, let P be a prime ideal of V , and let T be a ring containing V . Manis [11] defined the pair (V, P) to be a *valuation pair* of T if the following equivalent conditions are satisfied:

- (a) If S is a subring of T containing V and if M is a prime ideal of S such that $M \cap V = P$, then $V = S$;
- (b) For each x in $T \setminus V$, there exists y in P such that xy is in $V \setminus P$;
- (c) There is a valuation (v, G) on T such that

$$V = \{x \in T \mid v(x) \geq 0\} \quad \text{and} \quad P = \{x \in T \mid v(x) > 0\}.$$

If (V, P) is a valuation pair of T , then V is called a *valuation ring*. We will sometimes represent V by T_v . In this paper we always assume that T is $T(V)$. Chained rings are valuation rings, and in the domain case the concepts are equivalent to the concept of a valuation domain.

When we deal with rings containing zero divisors, we say that

an ideal is *regular* provided it contains a regular element, a nonzero divisor. We therefore define a *Prüfer ring* as a ring in which every finitely generated regular ideal is invertible. Fourteen equivalent definitions for a Prüfer ring were given by Griffin [7]. Moreover, examples by Gilmer [4] and Boisen and Larsen [1] show that a valuation ring need not be a Prüfer ring. Consequently, we define a *Prüfer valuation ring* as a valuation ring that is also a Prüfer ring.

A ring R has *few zero divisors* if the set of zero divisors of R is a finite union of prime ideals. A ring R is *additively regular* if for each x in $T(R)$ there exists a u in R such that $x + u$ is a regular element of $T(R)$. Davis [3] showed that a ring with few zero divisors is additively regular. Gilmer and Huckaba [6] have shown that the converse is not true. In his paper Davis defined what he called a quasi-valuation ring as a ring R with few zero divisors such that if x and y are in R , both regular, then x divides y or y divides x in R . However, Griffin [7] showed that if R is an additively regular ring, then R is a valuation ring if and only if R is a quasi-valuation ring. But then Huckaba [8] established that an additively regular valuation ring is a Prüfer ring. Consequently, we define R to be a *quasi-valuation ring* if x and y in R , both regular, implies that x divides y or y divides x in R .

2. **Structure theorems for chained rings.** If R is a chained ring, then obviously any homomorphic image of R is also a chained ring. Consequently, the homomorphic image of a valuation domain is a chained ring. By making use of recent results of Hungerford [9] and Boisen and Larsen [2], we are able to characterize a chained ring with Noetherian total quotient ring:

THEOREM 1. *Let R be a chained ring whose total quotient ring is a Noetherian ring. Then R is the homomorphic image of a valuation domain.*

Proof. Since R is a chained ring with $T(R)$ a Noetherian ring, then it is easy to see that $T(R)$ is a principal ideal ring and satisfies the descending chain condition. Hungerford [9] has shown that $T(R)$ is the homomorphic image of a valuation domain. Indeed, $T(R)$ is the homomorphic image of a Prüfer domain. However, since R is a Prüfer ring, Boisen and Larsen [2] have shown that R is the homomorphic image of a Prüfer domain, say D . Let π be the desired homomorphism and let M be the unique maximal ideal of R . There is an ideal M' of D such that $\pi(M') = M$. If $b \notin M'$, then $\pi(b) \notin M$, and hence $\pi(b)$ is a unit of R . The function defined by

$\pi^*: D_{M'} \rightarrow R$ such that $\pi^*(a/b) = \pi(a)\pi(b)^{-1}$ is an epimorphism. Therefore, R is the homomorphic image of the valuation domain $D_{M'}$.

The question remains open as to whether every chained ring can be realized as the homomorphic image of a valuation domain.

Let V be an arbitrary valuation ring with corresponding valuation v . If z is a zero divisor of V , then nothing can be determined about the image of z under v . This is not the case in chained rings, as our next result shows. It is basic to what follows.

THEOREM 2. (1) *Every chained ring is a valuation ring of its total quotient ring.*

(2) *If v is a valuation on T and T_v is chained, then $Z(T)$ is a maximal ideal of T and there is a valuation v' on the field $T/Z(T)$ such that v is given by $v(x) = v'(x + Z(T))$ for all x in T . Moreover, in this case $Z(T_v) = \{x \in T \mid v(x) = \infty\}$.*

Proof. For (1) let v' be the valuation of R_P/P_P whose valuation ring is R/P , where P is the prime ideal of zero divisors of the chained ring R . One sees easily that $R = T(R)_u$, where u is given by the formula $u(a/s) = v'(a + P) - v'(s + P)$, where a and s are in R and s is regular.

For the proof of (2), suppose $R = T_v$ is chained and P the prime ideal of zero divisors of R , so that $T = R_P$. Then every element of R_P either belongs to R or has the form $1/s$ for some s in $R \setminus P$. If $R = T(R)$, the result is straightforward. Otherwise, we have $v(1/s) < 0$ for some regular s in R . But then if r is an element of $Z(R)$ and $v(r) = 0$, we have $rR \subseteq sR$ so that $v(s) = 0$ for all regular s , a contradiction. Hence, if r is an element of $Z(R)$, $v(r) > 0$. If $v(r) < \infty$, there exists s regular in R such that

$$v(1/s) = -v(r), \text{ i.e. } v(r/s) = 0.$$

But then $r = r's$ and r' is a zero divisor for which $v(r') = 0$, a contradiction. This proves the "moreover" of (2). Now define a valuation of $T/Z(T)$ by $v'(x + Z(T)) = v(x)$. This is easily seen to be well defined, etc., and $Z(T) = P_P$ is maximal in T ; clearly v is equivalent to u , where u is defined as in (1).

We now prove our first structure theorem:

THEOREM 3. *Let T be a chained total quotient ring and V a valuation ring of T . Then V is a chained ring iff $Z(T) \subseteq V$, in which case V is quasilocal. Otherwise, V has exactly two maximal ideals, namely $Z(V)$ and a maximal regular ideal.*

Proof. If V is chained, then x is an element of $Z(T)$ if and only if $x = a/s$ with a and s elements of V and s regular, a an element of $Z(V)$. But clearly $aV \subseteq sV$, so a/s is an element of V . Conversely, if $Z(T) \subseteq V$, then for all a in $Z(V)$ and regular s in V , a/s is in V . Now let x and y be in V , say $xT \subseteq yT$, i.e. $x = y(a/s)$ for a and s in V , s regular. If a is regular, then $v(a/s)$ or $v(s/a)$ is ≥ 0 , i.e. a/s or s/a is in V . Hence, $xV \subseteq yV$ or $yV \subseteq xV$. If a is a zero divisor, then x is in yV . In either case, xV and yV are comparable.

Finally, if $Z(T) \not\subseteq V$, then there is a regular nonunit s in V and zero divisor such that a/s is not in V , i.e. $aV \not\subseteq sV$. So let M be the *unique* maximal ideal containing sV . [Regular primes are comparable by an argument of Davis [3]: if M and N are incomparable such ideals, then $M \not\subseteq N \cup Z(V)$, since $Z(V) = Z(T) \cap V$ is a prime ideal, whence there is a regular element x in $M \setminus N$, and symmetrically, regular y in $N \setminus M$. Since x and y are comparable, this is impossible.] and N a maximal ideal containing aV . Since $N \neq M$, N consists of zero divisors, so $N = Z(V)$, as required.

This theorem tells us precisely when a valuation ring with chained total quotient ring is or is not a chained ring. It rests completely on whether the zero divisors of the total quotient ring are or are not contained in the ring. Any homomorphic image of a valuation domain satisfies the first part of Theorem 3. We now construct an example which show the second part of Theorem 3 is realizable; that is, we construct a chained total quotient ring T that admits a valuation ring V such that $T(V) = T$, but V is not a chained ring.

EXAMPLE 1. Let K be a Prufer 2^∞ -group [2-primary part of the group \mathbf{Q}/\mathbf{Z}]. Then K is a $\mathbf{Z}_{(2)}$ -module in a natural way. Let $T = K + \mathbf{Z}_{(2)}$ be the idealization of the $\mathbf{Z}_{(2)}$ -module K [addition is coordinate-wise and multiplication is $(k_1 + z_1)(k_2 + z_2) = (k_1z_2 + k_2z_1) + z_1z_2$]. The divisibility of K shows that the ideals of T are exactly the subgroups of K and ideals of the form $K + J$, J an ideal of $\mathbf{Z}_{(2)}$. Hence T is a chained ring and in fact $Z(T) = K + (2)$, so T is a total quotient ring. Define a valuation on T by $v(k + z) = v'(z)$, where v' is the 3-adic valuation of \mathbf{Q} . Then $T_v = K + (\mathbf{Z}_{(2)} \cap \mathbf{Z}_{(3)})$, clearly, and $\mathbf{Z}_{(2)} \cap \mathbf{Z}_{(3)} = \mathbf{Z}_{(2) \cup (3)}$ is not a quasilocal ring. Since it is a homomorphic image of T_v , T_v is not a chained ring, yet the total quotient ring of T_v is T itself. Incidentally, in this case $Z(T) = K + (2)$ and $K + (3)$ are the two maximal ideals of T_v .

We now list three immediate corollaries to Theorem 3.

COROLLARY 1. *Let T be a chained total quotient ring and V a proper valuation ring of T such that $T = T(V)$.*

(a) *If V has exactly two maximal ideals, then every overring of V , other than T , is a valuation ring with exactly two maximal ideals;*

(b) *If V is a chained ring, then every overring of V is chained ring.*

COROLLARY 2. *Let R be a ring such that $T(R)$ is a chained ring. Let $\{V_\alpha\}$ be a set of valuation rings, totally ordered by inclusion, between R and $T(R)$. If one member of $\{V_\alpha\}$ is a chained ring, then every member of $\{V_\alpha\}$ is a chained ring. If one member of $\{V_\alpha\}$ has exactly two maximal ideals, then every member of $\{V_\alpha\}$ has exactly two maximal ideals.*

COROLLARY 3. *Every overring of a chained ring is a chained ring.*

We have seen by Example 1 that there exist chained total quotient rings T that admit valuation rings which are not chained rings. The content of the next theorem is that this situation cannot arise when the total quotient ring is a 0-dimensional ring.

THEOREM 4. *Let R be a ring with 0-dimensional total quotient ring, then the following are equivalent:*

- (a) *R admits at least one chained overring;*
- (b) *$T(R)$ is a chained ring; and*
- (c) *Every valuation overring of R is a chained ring.*

Proof. (a) \Rightarrow (b): Let V be a chained overring of R , then $T(V) = T(R)$ and by Corollary 3 $T(R)$ is a chained ring.

(b) \Rightarrow (c): If V is a valuation overring of R , let $T = T(V) = T(R)$. Since T is 0-dimensional, we have that $Z(T) = N(T)$ [the nil radical of T]. Since z in $Z(T)$ implies that there is an integer $n > 1$ such that $z^n = 0$, then z is integral over V , and therefore z is in V since V is integrally closed. Thus $Z(T) \subseteq V$, and we have that V is a chained ring.

(c) \Rightarrow (a): Obvious.

A ring R is a *von Neumann regular ring* if for each r in R , there is a b in R such that $r^2b = r$. von Neumann regular rings are 0-dimensional rings and have no nonzero nilpotents.

COROLLARY 4. *If R admits a chained overring, then $T(R)$ is*

von Neumann regular if and only if R is an integral domain.

We characterize rings in which every valuation overring is a chained ring.

THEOREM 5. *Let R be ring with total quotient ring T . Let \bar{R} be the integral closure of R in T . Every valuation overring of R is a chained ring if and only if $Z(T)$ is the conductor of \bar{R} in T .*

Proof. (\Rightarrow): Let V be an arbitrary valuation overring of R , then V is a chained ring. Clearly, $Z(T) = Z(V)$ is the conductor of V in T . Therefore, $Z(R) = Z(T) \cap R$, and R is an additively regular ring. This implies that \bar{R} is the intersection of a family of valuation rings $\{V_a\}$. So, $Z(T) \subseteq \bigcap V_a = \bar{R}$.

(\Leftarrow): If $Z(T)$ is the conductor of \bar{R} in T , then $Z(T)$ is the unique maximal ideal of T and thus is a prime ideal. Hence, $Z(R) = Z(T) \cap R$ is a prime ideal of R , and thus R is additively regular. Whence $\bar{R} = \bigcap V_a$ where $\{V_a\}$ is the set of valuation rings between R and $T(R)$. Therefore, $Z(T) \subseteq V_a$ for all a and hence each V_a is a chained ring.

Our final result is of interest in the light that it extends the theory of chained rings although it is not directly concerned with overrings. It gives necessary and sufficient conditions for a Prüfer valuation ring to be a chained ring.

Let P be a prime ideal of a ring R . The *core* of P , denoted by $C(P)$, is the set of all elements b in R such that for each regular r in R , there exists an element e in $R \setminus P$ such that be/r is in R . For a valuation ring V with $M = \{x \in T(V) \mid v(x) > 0\}$, the core of M is the prime ideal of zero for which the valuation, v , is infinite.

THEOREM 6. *Let V be a valuation ring with a unique maximal regular ideal M . Then V is a chained ring if and only if the following three conditions hold:*

- (a) V is a Prüfer ring;
- (b) $C(M) = Z(V)$;
- (c) $a, b \in C(M)$ implies that a divides b or b divides a in V .

Proof. (\Rightarrow): The fact that chained rings are Prüfer is trivial, since finitely generated regular ideals are principal ideals generated by regular elements. Theorem 2 tells us that

$$C(M) = \{x \in V \mid v(x) = \infty\} = Z(V).$$

And certainly (c) is true.

(\Leftarrow): If a and b are in V , we must show that a divides b or b divides a in V . This follows from the hypotheses, if a and b are in $C(M)$; and by the quasi-valuation property, if a and b are regular. Assume that a is regular and that b is a zero divisor. Then b/a is in $T(V)$ and $v(b/a) = v(b) - v(a)$. But $v(b) = \infty$ and $v(a) < \infty$. Thus, $v(b/a) = \infty$ which implies that b/a is in V , or that a divides b in V .

REFERENCES

1. M. Boisen, and M. Larsen, *Prüfer and valuation rings with zero divisors*, Pacific J. Math., **40** (1972), 7-12.
2. ———, *On Prüfer rings as images of Prüfer domains*, Proc. Amer. Math. Soc., **40** (1973), 87-90.
3. E. Davis, *Overrings of commutative rings II. Integrally closed overrings*, Trans. Amer. Math. Soc., **110** (1964), 196-212.
4. R. Gilmer, *On Prüfer rings*, Bull. Amer. Math. Soc., **78** (1972), 223-224.
5. ———, *Multiplicative Ideal Theory*. Queens Papers on Pure and Applied Mathematics, Vol. 12. Kingston, Ontario, Canada: Queens University Press, 1968.
6. R. Gilmer, and J. Huckaba, *Δ -rings*, J. Algebra, **28** (1974), 414-432.
7. M. Griffin, *Prüfer rings with zero divisors*, J. Reine Angew. Math., **239/240** (1970), 55-67.
8. J. Huckaba, *On valuation rings that contain zero divisors*, Proc. Amer. Math. Soc., **40** (1973), 9-15.
9. T. Hungerford, *On the structure of principal ideal rings*, Pacific J. Math., **25** (1968), 543-547.
10. M. Larsen and P. McCarthy, *Multiplicative Theory of Ideals*, New York, N. Y.: Academic Press, 1971.
11. M. Manis, *Valuations on a commutative ring*, Proc. Amer. Math. Soc., **20** (1969), 193-198.

Received May 2, 1975 and in revised form January 27, 1976. This paper is part of the author's Ph. D. thesis written under the direction of James A. Huckaba at the University of Missouri-Columbia.

SAINT MARY'S COLLEGE

Current Address: Saint Mary's College
Winona, Minnesota 55987

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 65, No. 1

September, 1976

David Lee Armacost, <i>Compactly cogenerated LCA groups</i>	1
Sun Man Chang, <i>On continuous image averaging of probability measures</i>	13
J. Chidambaraswamy, <i>Generalized Dedekind ψ-functions with respect to a polynomial. II</i>	19
Freddy Delbaen, <i>The Dunford-Pettis property for certain uniform algebras</i>	29
Robert Benjamin Feinberg, <i>Faithful distributive modules over incidence algebras</i>	35
Paul Froeschl, <i>Chained rings</i>	47
John Brady Garnett and Anthony G. O'Farrell, <i>Sobolev approximation by a sum of subalgebras on the circle</i>	55
Hugh M. Hilden, José M. Montesinos and Thomas Lusk Thickstun, <i>Closed oriented 3-manifolds as 3-fold branched coverings of S^3 of special type</i>	65
Atsushi Inoue, <i>On a class of unbounded operator algebras</i>	77
Peter Kleinschmidt, <i>On facets with non-arbitrary shapes</i>	97
Narendrakumar Ramanlal Ladhawala, <i>Absolute summability of Walsh-Fourier series</i>	103
Howard Wilson Lambert, <i>Links which are unknottable by maps</i>	109
Kyung Bai Lee, <i>On certain g-first countable spaces</i>	113
Richard Ira Loebel, <i>A Hahn decomposition for linear maps</i>	119
Moshe Marcus and Victor Julius Mizel, <i>A characterization of non-linear functionals on W_1^p possessing autonomous kernels. I</i>	135
James Miller, <i>Subordinating factor sequences and convex functions of several variables</i>	159
Keith Pierce, <i>Amalgamated sums of abelian l-groups</i>	167
Jonathan Rosenberg, <i>The C^*-algebras of some real and p-adic solvable groups</i>	175
Hugo Rossi and Michele Vergne, <i>Group representations on Hilbert spaces defined in terms of ∂_b-cohomology on the Silov boundary of a Siegel domain</i>	193
Mary Elizabeth Schaps, <i>Nonsingular deformations of a determinantal scheme</i>	209
S. R. Singh, <i>Some convergence properties of the Bubnov-Galerkin method</i>	217
Peggy Strait, <i>Level crossing probabilities for a multi-parameter Brownian process</i>	223
Robert M. Tardiff, <i>Topologies for probabilistic metric spaces</i>	233
Benjamin Baxter Wells, Jr., <i>Rearrangements of functions on the ring of integers of a p-series field</i>	253
Robert Francis Wheeler, <i>Well-behaved and totally bounded approximate identities for $C_0(X)$</i>	261
Delores Arletta Williams, <i>Gauss sums and integral quadratic forms over local fields of characteristic 2</i>	271
John Yuan, <i>On the construction of one-parameter semigroups in topological semigroups</i>	285