

Pacific Journal of Mathematics

**GAUSS SUMS AND INTEGRAL QUADRATIC FORMS OVER
LOCAL FIELDS OF CHARACTERISTIC 2**

DELORES ARLETTA WILLIAMS

GAUSS SUMS AND INTEGRAL QUADRATIC FORMS OVER LOCAL FIELDS OF CHARACTERISTIC 2

DELORES A. WILLIAMS

The theory of Gauss sums is developed for integral quadratic forms over a local field of characteristic 2, and Gauss sums are used to characterize these forms. For a character χ and an integral lattice L , the Gauss sum $\chi(L)$ is either zero, a nonnegative power of two, or the negative of a positive power of two. Gauss sums alone characterize the integral equivalence classes for modular lattices. For arbitrary lattices, other invariants are required.

The classification given in this paper is an alternate to the one by C.-H. Sah [6]. The notation and terminology of [6] is used except when stated to the contrary. O. T. O'Meara [4] used Gauss sums to characterize local integral quadratic forms over a field of characteristic not 2, and R. Jacobowitz [3] classified hermitian forms over the integers of a local field of characteristic not 2 by Gauss sums. When needed, results from these papers are referred to when the proofs hold for the characteristic 2 case.

After a few preliminaries, we introduce Gauss sums and prove some results for Gauss sums of lines and planes that will in turn be used to study more complicated lattices. In Theorem 5.4 we show that Gauss sums alone are sufficient to characterize modular lattices, and in Theorem 4.2 we show that for nondefective lattices only a finite number of Gauss sums need be considered.

1. Preliminaries. Throughout this paper k denotes a local field of characteristic 2 with fixed prime element π , ring of integers \mathfrak{o} , and residue class field of order 2^f . We let Ω denote a complete set of representatives for the residue class field. We refer the reader to [6] for a discussion of the Arf invariant ΔV for a quadratic space V and the additive group $\mathcal{P}\Omega$. As in [6] we let $\{0, \lambda\}$ be a fixed set of representatives of $\Omega/\mathcal{P}\Omega$. The letter e always denotes a unit of k . For a nonnegative integer s , H_s and H'_s denote s -hyperbolic lattices.

Let L be a lattice. $K(L) = \{x \in L \mid \langle x, y \rangle = 0 \text{ for all } y \in L\}$. If $K(L) = 0$, then L is nondefective. Otherwise, it is defective. We assume that if $x \in K(L)$ and $Q(x) = 0$, then $x = 0$.

We now state some lemmas and definitions from [6] in the form in which they are used in this paper.

LEMMA 1.1. Let L be a lattice with $(qL)v = \pi^u v$.

(1) If $\text{rank}_{\mathfrak{o}} q(L) = 1$, then $L = K(L)$ and $q(L) = e\pi^u v^2$ with e a unit.

(2) If $\text{rank}_{\mathfrak{o}} q(L) \geq 2$, then $v = \min\{\text{ord } a \mid a \in q(L) \text{ and } \text{ord } a \equiv u + 1 \pmod{2}\}$ is a rational integer with $u < v$. If, in addition, $L \neq K(L)$, $v \leq s + 1$ with $sL = \pi^s v$. Moreover, for any unit e with $e\pi^u \in Q(L)$, $q(L) = e(\pi^u v^2 + \pi^v v^2)$; and there is at least one such e .

If more that one lattice is under consideration, the invariants u and v for L are denoted by u_L and v_L respectively.

As in [6], for a rational integer i , $u(i)$ and $v(i)$ denote the u and v of Lemma 1.1 for the lattice $L(i) \otimes H_i$.

LEMMA 1.2. Let L be an i -modular lattice such that $\text{rank } L \geq 4$ and $q(L) = e(\pi^u v^2 + \pi^v v^2)$ with $u < v \leq i + 1$, $u + v \equiv 1 \pmod{2}$, and $e\pi^u \in Q(L)$ (as in Lemma 1.1). Then $L \simeq \begin{pmatrix} \pi^i & \\ e\pi^v & e^{-1}\pi^{2i-u}\delta + e^{-1}\pi^v a^2 \end{pmatrix} \oplus \begin{pmatrix} \pi^i & \\ e\pi^v & e^{-1}\pi^{2i-v}\delta' \end{pmatrix} \oplus H$ with $\delta, \delta' \in \{0, \lambda\}$, H i -hyperbolic or 0, and $\delta' = 0$ when $v = i + 1$. Moreover, $q(L) = Q(L)$.

For a canonical decomposition $L = (\sum_{i=0}^t L_i) \oplus K(L)$, $s(i)$ always denotes $s(L_i)$.

DEFINITION 1.3. Let $L = (\sum_{i=0}^t L_i) \oplus K(L)$ be a canonical decomposition. Define $I = s(0)$ and $T = s(t) + \max\{u', v'\}$, where

$$u' = \begin{cases} u_{K(L)} - u(s(t)) & \text{if } L \text{ is defective} \\ 0 & \text{otherwise} \end{cases}$$

and

$$v' = \begin{cases} v_{K(L)} - v(s(t)) & \text{if } \text{rank}_{\mathfrak{o}} q(K(L)) = 2 \\ 0 & \text{otherwise} \end{cases},$$

and defined $R = (r(I), r(I + 1), \dots, r(T))$ where

$$r(j) = \begin{cases} \text{rank } L_i & \text{if } s(i) = j \text{ for some } i \in \{0, \dots, t\} \\ 0 & \text{otherwise} \end{cases}.$$

Then L is said to be of type (I, T, R) and length $(t + 1)$.

DEFINITION 1.4. For $L \neq K(L)$, let $L = (\sum_{i=0}^t L_i) \oplus K(L)$ be a canonical decomposition of L . If $s(0), \dots, s(t)$ are consecutive integers, $\text{rank } L_i \geq 8$, and $u' = v' = 0$, then L_i is a normal lattice of type $(s(0), s(t), R)$ and $R = (\text{rank } L_0, \dots, \text{rank } L_t)$. Observe that

the concept of normal is independent of the canonical decomposition.

LEMMA 1.5. Let $L = \sum_{i=0}^t L_i$ and $M = \sum_{i=0}^{t'} M_i$ be canonical decomposition of the non-defective lattices L and M respectively. If L and M are of the same type, then $t = t'$, $\text{rank } L_i = \text{rank } M_i$, and $s_L(i) = s_M(i)$.

2. Characters. A character of \mathfrak{o} is a map χ from \mathfrak{o} into the complex numbers such that $\chi(a+b) = \chi(a)\chi(b)$ for $a, b \in \mathfrak{o}$, and $\chi(\mathfrak{o}(r)) = 1$ for some nonnegative rational integer r . Let $m(\chi)$ denote the smallest $r \geq 0$ such that $\chi(\mathfrak{o}(r)) = 1$. Then $\mathfrak{o}(m(\chi))$ is called the *maximal support* of χ and $m(\chi)$ is called the *maximal support ordinal* of χ . When a single character χ is being considered, we use m for the maximal support ordinal of χ without stating that $m = m(\chi)$.

Note that if χ is a character and $a \in \mathfrak{o}$, then $\chi(a) = \pm 1$. In fact, the image of χ is $\{-1, 1\}$ except when $m(\chi) = 0$.

The set of all characters of \mathfrak{o} , denoted by X , together with the operation of function multiplication is a group. For $r \geq 0$ define $X(r) = \{\chi \in X \mid \chi(\mathfrak{o}(r)) = 1\} = \{\chi \in X \mid m(\chi) \leq r\}$. $X(r)$ is a subgroup of X and will play an important role in the study of Gauss sums.

For each $r \geq 0$ there is a one-to-one correspondence between $X(r)$ and the set of group characters on $\mathfrak{o}/\mathfrak{o}(r)$. This permits us to obtain information about $X(r)$ from the theory of group characters. (For details see [3].)

3. Gauss sums. A lattice L is said to be *integral* if $Q(L) \subseteq \mathfrak{o}$. If L is integral, then $q(L) \subseteq \mathfrak{o}$, since for $x, y \in L$, $\langle x, y \rangle = Q(x) + Q(y) + Q(x+y) \in \mathfrak{o}$. All lattices will be assumed to be integral. This causes no loss of generality in studying the classification problem.

For $r \geq 0$, $L(r)$ will denote $\pi^r L$ and a summation over $x \bmod L(r)$ will mean that x runs over a complete set of representatives for the additive group $L/L(r)$. The order of $L/L(r)$ is $2^{fr(\text{rank } L)}$.

For a lattice L and a character $\chi \in X(r)$, $\chi(L; \pi^r) = \sum_{x \bmod L(r)} \chi(Q(x))$ is called a *Gauss sum*. Write $\chi(L)$ for $\chi(L; \pi^m)$. Since L is integral and $L/L(r)$ is finite, all elements of $Q(L)$ are in the domain of χ and each Gauss sum is a finite sum of ones and negative-ones.

For a plane $P \cong \begin{pmatrix} b \\ a & c \end{pmatrix}$ (or a line $P \cong (a)$), we will sometimes write $\chi \begin{pmatrix} b \\ a & c \end{pmatrix}$ (or $\chi(a)$) for $\chi(P)$.

In view of [4, Propositions 1 and 2], it is clear that a great

deal can be learned about Gauss sums by studying $\chi(L)$ when L is a line or plane. Our investigation can be further simplified by noting:

$$(3.1) \quad \begin{pmatrix} b \\ a \ c \end{pmatrix} \oplus \begin{pmatrix} b \\ a \ c \end{pmatrix} \cong \begin{pmatrix} b \\ a \ 0 \end{pmatrix} \oplus \begin{pmatrix} b \\ c \ 0 \end{pmatrix}$$

and

$$(3.2) \quad \chi \begin{pmatrix} \pi^m \\ d \ 0 \end{pmatrix} = 2^{fm} \chi(d).$$

Let $L = \alpha x_1 + \cdots + \alpha x_n$. It follows from the uniqueness of the representation of elements with respect to a basis that $\alpha_1 x_1 + \cdots + \alpha_n x_n$ runs through a complete set of representatives for $L/L(m)$ as $\alpha_1, \dots, \alpha_n$ each runs through a complete set of representatives for $\mathfrak{o}/\mathfrak{o}(m)$.

LEMMA 3.1. For $\chi \in X$ and $L \cong \begin{pmatrix} \pi^s \\ a \ 0 \end{pmatrix}$,

$$\chi(L) = 2^{fm} \sum_{\alpha \bmod \mathfrak{o}(s)} \chi(\alpha^2 a \pi^{2m-2s})$$

if $s \leq m$ and $\chi(L) = 2^{fm} \sum_{\alpha \bmod \mathfrak{o}(m)} \chi(\alpha^2 a)$ if $m < s$.

Proof. Keeping in mind the preceding comment, $\chi(L) = \sum_{\alpha, \beta} \chi(\alpha^2 a + \alpha \beta \pi^s)$ with $\alpha \bmod \mathfrak{o}(m)$ and $\beta \bmod \mathfrak{o}(m)$. For a fixed α , by [3, (1.6)], $\sum_{\beta} \chi(\alpha \beta \pi^s) = \begin{cases} 2^{fm} & \text{if } \text{ord } \alpha + s \geq m \\ 0 & \text{otherwise} \end{cases}$ with $\beta \bmod \mathfrak{o}(m)$. Therefore $\chi(L) = 2^{fm} \sum \chi(\alpha^2 a)$ with $\alpha \bmod \mathfrak{o}(m)$ and $\text{ord } \alpha \geq m - s$.

Since the case $s > m$ is immediate, assume $s \leq m$. As γ runs through a complete set of representatives for $\mathfrak{o}/\mathfrak{o}(s)$, $\alpha = \gamma \pi^{m-s}$ runs through a complete set of those representatives for $\mathfrak{o}/\mathfrak{o}(m)$ such that $\text{ord } \alpha \geq m - s$. Thus $\chi(L) = 2^{fm} \sum_{\gamma} \chi(\gamma^2 a \pi^{2m-2s})$.

PROPOSITION 3.2. For $\chi \in X$ and H an i -hyperbolic plane, $\chi(H) = 2^{f \min(2m, m+i)}$.

Proof. This follows from Lemma 3.1 and the fact that $\chi(0) = 1$.

LEMMA 3.3. Let $\chi \in X$, $L \cong \begin{pmatrix} \pi^s \\ a \ 0 \end{pmatrix}$, and $M \cong \begin{pmatrix} \pi^{s+w} \\ a \pi^{2w} \ 0 \end{pmatrix}$ with $0 \leq w \leq s + w \leq m$. Then $\chi(M) = 2^{fw} \chi(L)$.

Proof. Using Lemma 3.1, $\chi(M) = 2^{fm} \sum \sum \chi((\alpha + \beta \pi^s)^2 a \pi^{2m-2s}) = 2^{f(m+w)} \sum \chi(\alpha^2 a \pi^{2m-2s}) = 2^{fw} \chi(L)$ with $\alpha \bmod \mathfrak{o}(s)$ and $\beta \bmod \mathfrak{o}(w)$.

PROPOSITION 3.4. *Let $L \cong \begin{pmatrix} \pi^s \\ e\pi^w & 0 \end{pmatrix}$ and $m \geq 0$. Then the following hold:*

- (1) *If $m \leq w$ or $2s - w \leq m$, then $\chi(L) = 2^{f \min(2m, m+s)}$.*
- (2) *If $w < m < 2s - w$ with w and m of opposite parity, then $\chi(L) = 0$.*
- (3) *If $w < m < 2s - w$ with w and m of the same parity, then there exists χ_1 and χ_2 in X with $m(\chi_1) = m(\chi_2) = m$ such that $\chi_1(L) = 0$ and $\chi_2(L) \neq 0$. Furthermore, if $m(\chi) = m$ and $\chi(L) \neq 0$, then $\chi(L) = 2^{f \min(2m, m+s)}$.*

Proof. (1) If $m \leq w$, $\chi(L) = \chi(H_s)$; otherwise use Lemma 3.1.

(2) *Claim.* If χ has maximal support $\mathfrak{o}(m)$, $r + m \equiv 1 \pmod{2}$, and $j \geq (m - r + 1)/2 > 0$, then $\sum_{\alpha} \chi(\alpha^2 e\pi^r) = 0$ with $\alpha \pmod{\mathfrak{o}(j)}$.

Justification. With $\beta \pmod{\mathfrak{o}((m - r - 1)/2)}$, $\gamma \pmod{\mathfrak{o}(1)}$, and $\delta \pmod{\mathfrak{o}(j - (m - r + 1)/2)}$, $\sum \chi(\alpha^2 e\pi^r) = \sum_{\beta} \sum_r \sum_{\delta} \chi(\beta + \pi^{(m-r-1)/2} \gamma + \pi^{(m-r+1)/2} \delta)^2 e\pi^r) = [\sum_{\beta} \chi(\beta^2 e\pi^r)] [\sum_r \chi(\gamma^2 e\pi^{m-1})] [\sum_{\delta} \chi(\delta^2 e\pi^{m+1})] = 0$ since $\sum_r \chi(\gamma^2 e\pi^{m-1}) = \sum_r \chi_{(m-1)}(\gamma) = 0$, and the claim holds.

If $s \leq m$, take $r = 2m - 2s + w$ and $j = s$; and if $m < s$, take $r = w$ and $j = m$.

(3) If $s < m$, then by Lemma 3.3 L may be replaced by $\begin{pmatrix} \pi^m \\ e\pi^{w+2m-2s} & 0 \end{pmatrix}$. Hence we assume $m \leq s$. Suppose $\chi(L) = 0$ for all χ with $m(\chi) = m$. Let $M \cong \begin{pmatrix} \pi^s \\ e\pi^w & \pi^{m-1} \end{pmatrix}$. For $\chi \in X(m)$, $\chi(M) = \chi(L)$. By [4, Proposition 3], $L(\pi^{m-1}; \pi^m) = M(\pi^{m-1}; \pi^m)$. Thus $\pi^{m-1} \in Q(L)$ which is impossible. Hence there exists χ_2 with $m(\chi_2) = m$ and $\chi_2(L) \neq 0$.

Now suppose $\chi(L)$ is nonzero for all χ with $m(\chi) = m$. Write $N \cong \begin{pmatrix} \pi^s \\ e\pi^{m-1} & 0 \end{pmatrix}$ and $N' \cong \begin{pmatrix} \pi^s \\ e\pi^w + e\pi^{m-1} & 0 \end{pmatrix}$. Using [6, LT₂ b)], $N \oplus L \cong N' \oplus L$. Now $e\pi^w + e\pi^{m-1}$ is a unit times π^w and therefore by the argument just gone through for L , $\chi(N') \neq 0$ for some χ with $m(\chi) = m$. Thus $\chi(N) \neq 0$; but this contradicts (2). Therefore there exists χ_1 with $m(\chi_1) = m$ and $\chi_1(L) = 0$.

If $\chi(L) \neq 0$, setting $H \cong \begin{pmatrix} \pi^s \\ 0 & 0 \end{pmatrix}$, $L \oplus L \cong H \oplus L$ and $\chi(L) = \chi(H) = 2^{f \min(2m, m+s)}$.

It now follows that if $\chi(L) \neq 0$ then $\chi(L) = \pm 2^n$ for some non-negative integer n .

4. A reduction.

LEMMA 4.1. *Let $L \cong \begin{pmatrix} \pi^s \\ a & c \end{pmatrix}$ and $M \cong \begin{pmatrix} \pi^r \\ a\pi^{2r-2s-2n} & c\pi^{2n} \end{pmatrix}$ with $r \geq s + n$ and $n \geq 0$. For any $\chi \in X$ such that $\chi \begin{pmatrix} \pi^{s+n} \\ a & 0 \end{pmatrix}, \chi \begin{pmatrix} \pi^s \\ 0 & c \end{pmatrix}$,*

and $\chi \begin{pmatrix} \pi^r & \\ 0 & c\pi^{2n} \end{pmatrix}$ are all nonzero, $\text{sgn}(\chi(L)) = \text{sgn}(\chi(M))$, and $\chi(L)$ and $\chi(M)$ are both nonzero.

Proof. Now $\begin{pmatrix} \pi^s & \\ a & c \end{pmatrix} \oplus \begin{pmatrix} \pi^{s+n} & \\ a & c\pi^{2n} \end{pmatrix} \cong \begin{pmatrix} \pi^s & \\ 0 & c \end{pmatrix} \oplus \begin{pmatrix} \pi^{s+n} & \\ a & 0 \end{pmatrix}$ and $\begin{pmatrix} \pi^{s+n} & \\ a & c\pi^{2n} \end{pmatrix} \oplus \begin{pmatrix} \pi^r & \\ a\pi^{2r-2s-2n} & c\pi^{2n} \end{pmatrix} \cong \begin{pmatrix} \pi^{s+n} & \\ a & 0 \end{pmatrix} \oplus \begin{pmatrix} \pi^r & \\ 0 & c\pi^{2n} \end{pmatrix}$. Since the Gauss sum of the lattice on the right-hand side in each isometry is positive, the result follows.

THEOREM 4.2. *Let $L = \sum_{i=0}^t L_i$ and $M = \sum_{i=0}^{t'} M_i$ be canonical decompositions for non-defective spaces L and M respectively, and let $u = \min(u_L(s(t)), u_M(s(t')))$. If L and M are of the same type (I, T, R) , and if $\chi(L) = \chi(M)$ for all $\chi \in X(2T - u + 1)$, then $\chi(L) = \chi(M)$ for all $\chi \in X$.*

Proof. Fix $\chi \in X$ such that $m > 2T - u + 1$. Write $m = 2T - u + 2n + j$ where $j = 0$ or $j = 1$. In view of Lemma 1.5 $\text{rank } L = \text{rank } M$ and $s_L(i) = s_M(i)$ for $0 \leq i \leq t$. Fix i . Adjoining hyperbolic planes if necessary, decompose L according to Lemma 1.2. Applying Proposition 3.2 to the hyperbolic part and Lemma 4.1 and Proposition 3.4 to the rest, we conclude that $\chi(L) = 2^{f_n \text{rank } L} \chi_{(2n)}(L)$. Similarly $\chi(M) = 2^{f_n \text{rank } M} \chi_{(2n)}(M)$.

5. Classification of modular lattices.

PROPOSITION 5.1. *If $sL = sM = v(s)$ and $\chi(L) = \chi(M)$ for all $\chi \in X(s)$, then $q(L) = q(M)$.*

Proof. *Case 1.* $s = 0$. Using the fact that L has an 0-modular orthogonal summand and Lemma 1.2, $q(L) = v^2 + \pi v^2 = q(M)$.

Case 2. $s \geq 1$. First assume there exists $\chi \in X$ such that $m(\chi) = s$ and $\chi(L) \neq 0$. Then $\text{sgn}(\chi(L))2^{f_m \text{rank } L} = \chi(L) = \chi(M) = \text{sgn}(\chi(M))2^{f_m \text{rank } M}$. Hence $\text{rank } L = \text{rank } M$, and by [4, Proposition 3] $L(a; \pi^s) \cong M(a; \pi^s)$ for all $a \in v$. It follows that $q(L) = q(M)$.

Now assume $\chi(L) = \chi(M) = 0$ for all $\chi \in X$ with $m(\chi) = s$. Assume without loss of generality that $\text{rank } L \leq \text{rank } M$. Let $L' \cong L \oplus H \oplus N$ such that $\text{rank } L' = \text{rank } M$, H is an s -hyperbolic lattice or 0, and $N \cong (\pi^s)$ or 0. Then $q(L') = q(L)$ and $\chi(L') = 0$ for all $\chi \in X(s)$. Hence $\chi(L') = \chi(M)$ for all $\chi \in X(s)$. Then [4, Proposition 3] is applicable and it follows that $q(L) = q(L') = q(M)$.

PROPOSITION 5.2. *Let $\chi \in X$ with $m \geq 1$, and let $L \cong$*

$\begin{pmatrix} \pi^{m-1} \\ e\pi^{m-1} & e^{-1}\pi^{m-1}\delta \end{pmatrix}$. Then $\chi(L) < 0$ if $\delta = \lambda$ and $\chi(L) > 0$ if $\delta = 0$.

Proof. If $e \neq 1$ set $J \cong L \oplus H_{m-1}$. By Lemma 1.2 and the uniqueness of $\mathcal{A}(kJ)$, $J \cong \begin{pmatrix} \pi^{m-1} \\ \pi^{m-1} & \pi^{m-1}\delta \end{pmatrix} \oplus H_{m-1}$. Thus we may reduce to the case in which $e = 1$. And since $\chi(L) = 2^{2f(m-1)}\chi_{(m-1)} \begin{pmatrix} 1 \\ 1 & \lambda \end{pmatrix}$, we assume $m = 1$.

If $\delta = 0$ use Proposition 3.4. Assume $\delta = \lambda$. Fix a basis $\{p_1, \dots, p_f\}$ for Ω over $\{0, 1\}$. With a_1, \dots, a_f running through $\{0, 1\}$, $\chi(L) = \sum_{a_1} \dots \sum_{a_f} [\chi((a_1p_1 + \dots + a_fp_f)^2\lambda)]$

$$[1 + \chi(p_1^2 + a_1p_1^2 + \dots + a_fp_1p_f)] \dots [1 + \chi(p_f^2 + a_1p_1p_f + \dots + a_fp_f^2)] .$$

Since $m = 1$, at least one element in each basis for Ω must be mapped to -1 by χ . Since $\lambda \notin \mathcal{P}\Omega$, it follows from elementary linear algebra that if $\{a_1, \dots, a_f\} \subseteq \{0, 1\}$, $\{(a_1p_1 + \dots + a_fp_f)^2, p_1^2 + a_1p_1^2 + \dots + a_fp_1p_f, \dots, p_f^2 + a_1p_1p_f + \dots + a_fp_f^2\}$ contains a basis for Ω . Thus $\chi(L) < 0$.

PROPOSITION 5.3. Let L be an s -modular lattice with $\text{rank } L \geq 4$ and $q(L) = e(\pi^u\mathfrak{o}^2 + \pi^v\mathfrak{o}^2)$. Write $M \cong \begin{pmatrix} \pi^s \\ e\pi^u & 0 \end{pmatrix}$ and $N \cong \begin{pmatrix} \pi^s \\ e\pi^v & 0 \end{pmatrix}$. For $\chi \in X$, $\chi(L) = 0$ iff $\chi(M) = 0$ or $\chi(N) = 0$.

Proof. Write $L \cong \begin{pmatrix} \pi^s \\ e\pi^u & c \end{pmatrix} \oplus \begin{pmatrix} \pi^s \\ e\pi^v & e^{-1}\pi^{2s-v}\delta' \end{pmatrix} \oplus H_s$ with $c = e^{-1}\pi^{2s-u}\delta + e^{-1}\pi^v\mathfrak{a}^2$ as in Lemma 1.2.

Sufficiency is obvious, so assume $\chi(L) = 0$. Set $M' \cong \begin{pmatrix} \pi^s \\ c & 0 \end{pmatrix}$ and $N' \cong \begin{pmatrix} \pi^s \\ e^{-1}\pi^{2s-v}\delta' & 0 \end{pmatrix}$. By Propositions 3.2 and 3.4 $\chi(N') \neq 0$. Thus at least one of $\chi(M)$, $\chi(N)$, and $\chi(M')$ is zero. If $\chi(M') = 0$, by Proposition 3.4 either $\chi(M) = 0$ or $\chi(N) = 0$.

THEOREM 5.4. Let L_1 and L_2 be s -modular lattices. $L_1 \cong L_2$ iff $\chi(L_1) = \chi(L_2)$ for all $\chi \in X(2s - \max(u_{L_1}, u_{L_2}) + 1)$.

Proof. Necessity is obvious, so we assume $\chi(L_1) = \chi(L_2)$ for $\chi \in X(2s - \max(u_{L_1}, u_{L_2}) + 1)$. By Proposition 5.1, $u_{L_1} = u_{L_2}$. Call the common value u . By Propositions 3.4 and 5.3, $\chi(L_i) = \text{sgn}(\chi(L_i))2^{f(m+s)(\text{rank } L_i)/2}$ when $m = 2s - u + 1$. Thus $\text{rank } L_1 = \text{rank } L_2$. Adjoining the appropriate hyperbolic plane to L_1 and L_2 and applying Lemma 1.2 and Proposition 5.1, $Q(L_1) = q(L_1) = q(L_2) = Q(L_2) = e(\pi^u\mathfrak{o}^2 + \pi^v\mathfrak{o}^2)$ and

$$L_i \cong \begin{pmatrix} \pi^s \\ e\pi^u & e^{-1}\pi^{2s-u}\delta_i + e^{-1}\pi^{u_i}\mathfrak{a}_i^2 \end{pmatrix} \oplus \begin{pmatrix} \pi^s \\ e\pi^v & e^{-1}\pi^{2s-v}\delta'_i \end{pmatrix} \oplus H_s$$

with a_i a unit or zero, $w_i \geq v$, and $w_i + v \equiv 0 \pmod{2}$. By showing that $\text{ord}(\pi^{w_1}a_1^2 + \pi^{w_2}a_2^2) > 2s - u$ and $\delta_1 = \delta_2$, it will follow from [6, Lemma 4.1] that $\begin{pmatrix} \pi^s \\ e\pi^u & e^{-1}\pi^{2s-u}\delta_1 + e^{-1}\pi^{w_1}a_1^2 \end{pmatrix} \cong \begin{pmatrix} \pi^s \\ e\pi^u & e^{-1}\pi^{2s-u}\delta_2 + e^{-1}\pi^{w_2}a_2^2 \end{pmatrix}$. If we also show that $\delta'_1 = \delta'_2$, we will have $L_1 \cong L_2$.

By [6, Theorem 4.6],

$$\begin{aligned} L_1 \oplus L_2 \cong & \begin{pmatrix} \pi^s \\ e\pi^u & e^{-1}\pi^{2s-u}(\delta_1 + \delta_2) \end{pmatrix} \oplus \begin{pmatrix} \pi^s \\ e\pi^u & e^{-1}\pi^w a^2 \end{pmatrix} \\ & \oplus \begin{pmatrix} \pi^s \\ e\pi^v & e^{-1}\pi^{2s-v}(\delta'_1 + \delta'_2) \end{pmatrix} \oplus H'_s \end{aligned}$$

where $\pi^w a^2 = \pi^{w_1}a_1^2 + \pi^{w_2}a_2^2$ with $w \geq v$, $w + v \equiv 0 \pmod{2}$, and a is a unit or zero. Let $K \cong \begin{pmatrix} \pi^s \\ e\pi^u & e^{-1}\pi^{2s-u}(\delta_1 + \delta_2) \end{pmatrix}$, $M \cong \begin{pmatrix} \pi^s \\ e\pi^u & e^{-1}\pi^w a^2 \end{pmatrix}$, and $N \cong \begin{pmatrix} \pi^s \\ e\pi^v & e^{-1}\pi^{2s-v}(\delta'_1 + \delta'_2) \end{pmatrix}$. For $\chi \in X$ with $m = 2s - u$, $\chi(K \oplus M \oplus N \oplus H'_s) = \chi(L_1 \oplus L_2) > 0$ and $\chi(K \oplus H_s) > 0$. Thus $\text{sgn}(\chi(M)) = \text{sgn}(\chi(N))$.

Claim. If $\text{ord}(e^{-1}\pi^w a^2) \leq 2s - u$, then there exists $\chi', \chi'' \in X$ with $m(\chi') = m(\chi'') = 2s - u$ such that $\chi'(M) > 0$ and $\chi''(M) < 0$.

Justification. Since $w + u \equiv 1 \pmod{2}$, $\text{ord}(e^{-1}\pi^w a^2) \neq 2s - u$. Write $J \cong \begin{pmatrix} \pi^{2s-u-1} \\ e\pi^{w-1} & e^{-1}\pi^{2s-u-1}a^2 \end{pmatrix}$, $J' \cong \begin{pmatrix} \pi^{2s-u-1} \\ e\pi^{w-1} & e^{-1}\pi^{2s-u+1}a^2 \end{pmatrix}$, $J'' \cong \begin{pmatrix} \pi^{2s-u-1} \\ 0 & e^{-1}\pi^{2s-u-1}a^2 \end{pmatrix}$, and $J''' \cong \begin{pmatrix} \pi^{2s-u-1} \\ e\pi^{w-1} & 0 \end{pmatrix}$. $J \oplus J' \cong J'' \oplus J'''$. There exists $\chi' \in X$ with $m = 2s - u$ such that $\chi'(J') > 0$ and $\chi'(J'' \oplus J''') > 0$. Hence $\chi'(J) > 0$. Taking $P \cong \begin{pmatrix} \pi^{2s-u-1} \\ e\pi^w & 0 \end{pmatrix}$, by an argument like that given in the proof of Proposition 3.4 (3), $\chi''(J) \leq 0$ for some $\chi'' \in X$ with $m = 2s - u$. Now compare M with J by Lemma 4.1 to establish the claim.

It is easily seen that $\chi'(N) = \chi''(N)$. Thus $\text{ord}(e^{-1}\pi^w a^2) > 2s - u$ and $\delta'_1 = \delta'_2$.

By another application of Lemma 4.1, $\delta_1 = \delta_2$.

6. Classification when $L = K(L)$. If $\text{rank } L = 1$, then $L \cong (e\pi^u)$. Otherwise $\text{rank } L = 2$, in which case $\text{rank}_{s_2} q(L) = 2$ and $L \cong (e\pi^u) \oplus (e\pi^v)$.

THEOREM 6.1. Let $L = K(L)$, $M = K(L)$, and $\text{rank } L \leq \text{rank } M$.

Write $w = \begin{cases} \max(u_L, u_M) & \text{if } \text{rank } L = \text{rank } M = 1 \\ \max(u_L, u_M, v_M) & \text{if } \text{rank } L = 1 \text{ and } \text{rank } M = 2 \\ \max(u_L, v_L, u_M, v_M) & \text{if } \text{rank } L = \text{rank } M = 2 \end{cases}$

- (1) If $\chi(L) = \chi(M)$ for $\chi \in X(w+1)$, then $\text{rank } L = \text{rank } M$.
 (2) For $L \cong (e\pi^{u_L})$ and $M \cong (e'\pi^{u_M})$, $L \cong M$ if and only if $e\pi^{u_L} \in e'\pi^{u_M}k^2$ and $\chi(L) = \chi(M)$ for all $\chi \in X(w)$.
 (3) If $\text{rank } L = \text{rank } M = 2$, then $L \cong M$ if and only if $\chi(L) = \chi(M)$ for all $\chi \in X(w)$.

Proof. Write $M \cong (e'\pi^{u_M})$ or $M \cong (e'\pi^{u_M}) \oplus (e'\pi^{v_M})$ and $L \cong (e\pi^{u_L})$ or $L \cong (e\pi^{u_L}) \oplus (e\pi^{v_L})$ depending on the case at hand. Assume that $\chi(L) = \chi(M)$ for all $\chi \in X(w)$ and that $u_L \leq u_M$. Setting $L' \cong (e\pi^{u_L})$ and using (3.2) and Proposition 3.4, $u_M = u_L$.

(1) Assume $\text{rank } M = 2$. Then v_M exists, and comparing L' and $(e'\pi^{v_M})$ with appropriate lattices of rank 2, we see that $\chi(L') \neq 0$ for some χ with $m = v_M + 1$ and $\chi(e'\pi^{v_M}) = 0$ for all χ with $m = v_M + 1$. Hence L' and L are not isometric, and $\text{rank } L = 2$.

(2) Since $u_L = u_M$, this is obvious.

(3) Necessity is obvious. $L(a; \pi^w) = M(a; \pi^w)$ for all $a \in \mathfrak{o}$. By a straightforward calculation $q(L) = q(M)$, and by [6, Lemma 1.2] $v_L = v_M$. It now follows that $L \cong M$.

7. Classification when $L \neq K(L)$.

PROPOSITION 7.1. *Let $L = (\sum_{i=0}^t L_i) \oplus K(L)$ be a saturated decomposition for the normal lattice L . For any $\chi \in X$ such that $\chi(L_0) = 0$, $\chi(L_0^\perp) \geq 0$.*

Proof. Suppose $\chi(L_0) = 0$ and $\chi(L_0^\perp) \neq 0$ for some $\chi \in X$. Using the hypothesis that the given decomposition is saturated and [6, Definition 5.1], there exists a unit e such that $Q(L_j) = e(\pi^{u(s(j))}\mathfrak{o}^2 + (\pi^{v(s(j))}\mathfrak{o}^2))$ for $j = 0, 1$. By Propositions 5.3 and 3.4, $u(s(0)) < m$ $2s(0) - u(s(0))$.

We show that $m \leq v(s(1))$. Then $m \leq v(s(i))$ and hence $\chi(L_i) \geq 0$ for $1 \leq i \leq t$. Since $\chi(K(L)) \geq 0$, we then have $\chi(L_0^\perp) > 0$.

Assume $v(s(1)) < m$. Since $\chi(L_1) \neq 0$ and $u(s(1)) < m < 2s(1) - u(s(1))$, $2s(0) - v(s(0)) \leq 2s(1) - v(s(1)) \leq m$. So $\chi \begin{pmatrix} \pi^{s(0)} & \\ e\pi^{u(s(0))} & 0 \end{pmatrix} = 0$. By [6, Definition 5.1], one of $u(s(1))$ and $v(s(1))$ equals either $u(s(0))$ or $u(s(0)) + 2$. Since $\begin{pmatrix} \pi^{s(0)} & \\ e\pi^{u(s(0))} & 0 \end{pmatrix} \oplus \begin{pmatrix} \pi^{s(1)} & \\ e\pi^{u(s(0))} & 0 \end{pmatrix} \cong \begin{pmatrix} \pi^{s(0)} & \\ 0 & 0 \end{pmatrix} \oplus \begin{pmatrix} \pi^{s(1)} & \\ e\pi^{u(s(0))} & 0 \end{pmatrix}$, $\chi \begin{pmatrix} \pi^{s(1)} & \\ e\pi^{u(s(0))} & 0 \end{pmatrix} = 0$; and by Lemma 3.3, if $s(1) \leq m$, $\chi \begin{pmatrix} \pi^{s(1)} & \\ e\pi^{u(s(0))+2} & 0 \end{pmatrix} = 0$. Thus $\chi(L_1) \neq 0$ implies $s(1) > m$. Hence $s(1) > m \geq 2s(1) - v(s(1)) \geq s(1) - 1$. So $v(s(1)) \geq m$.

PROPOSITION 7.2. *Let $L = (\sum_{i=0}^t L_i) \oplus K(L) = (\sum_{i=0}^t M_i) \oplus K(L)$ be canonical decompositions of a normal lattice L . If $u(j) =$*

$u(j-1) + 2$ and $v(j) = v(j-1) + 2$ for some j with $I < j \leq T$, write $\sum_{i=0}^{j-I-1} (\Delta(kL_i) + \Delta(kM_i)) = \beta^2\pi^{-1} + \delta + \mathcal{P}k$ with $\beta \in \Omega[\pi^{-1}]$ and $\delta \in \{0, \lambda\}$. Then β and δ are uniquely determined by the two decompositions and $\text{ord}(\beta^2\pi^{-1} + \delta) \geq u(j) + v(j) - 2j - 1$.

Proof. The uniqueness of β and δ follows from [6, Lemma 1.1].

By [6, Theorems 2.2 and 2.3 and Lemma 1.2], a reduction can be made to the case in which the two canonical decompositions are complete and the second is obtained from the first by an elementary lattice transformation.

Denote $u(j-1)$, $v(j-1)$, and $j-I$ by u , v , and w , respectively. Let T_r denote the elementary lattice transformation used to obtain the second decomposition from the first. Only the cases for which $r = 4$ and $r = 9$ need be considered.

Case 1. $r = 9$. If the two planes effected by T_9 are of the same modularity, the result holds trivially. If not, a reduction can be made to the case in which T_9 acts on L_{w-1} and L_w . Then for some $c, d \in Q(L_{w-1})$, $c', d' \in Q(L_w)$, and $\alpha \in \mathfrak{o}$, T_9 sends $\begin{pmatrix} \pi^{j-1} \\ c & d \end{pmatrix} \oplus \begin{pmatrix} \pi^j \\ c' & d' \end{pmatrix}$ to $\begin{pmatrix} \pi^{j-1} \\ c + \alpha^2 c' & d \end{pmatrix} \oplus \begin{pmatrix} \pi^j \\ c' & d' + \alpha^2 \pi^2 d \end{pmatrix}$ and leaves the other planes fixed.

By a direct computation, $\sum_{i=0}^{w-1} (\Delta(kL_i) + \Delta(kM_i)) = \alpha^2 c' d \pi^{-2j+2} + \mathcal{P}k = \beta^2\pi^{-1} + \delta + \mathcal{P}k$ with $\text{ord}(\beta^2\pi^{-1} + \delta) \geq u(j) + v(j) - 2j - 1$, where $\beta \in \Omega[\pi^{-1}]$ and $\delta \in \{0, \lambda\}$.

Case 2. $r = 4$. If the plane altered by the application of T_4 has modularity greater than $j-1$, the result follows. Otherwise there exists a sequence of complete decompositions from the first given decomposition to the second, each of which can be obtained from its predecessor by either an application of T_9 or an application of T_4 to a plane of modularity $j-1$. Assume T_4 is applied to a plane of modularity $j-1$. For some $c, d \in Q(L_{w-1})$, $c' \in Q(K(L))$, and $\alpha \in \mathfrak{o}$, T_4 sends $\begin{pmatrix} \pi^{j-1} \\ c & d \end{pmatrix}$ to $\begin{pmatrix} \pi^{j-1} \\ c + \alpha^2 c' & d \end{pmatrix}$ and leaves all else fixed. Now $c' \in Q(L_w)$; so the proof follows exactly as in Case 1.

NOTATION. Let L and M be lattices of the same type (I, T, R) , j be a rational integer such that $I < j < T$, and $H = \sum_{i=1}^T H_i$ with H_i i -hyperbolic of rank 8. $L \oplus H$ and $M \oplus H$ are normal lattices of type $(I, T, R + R(H))$; and if $(\sum_{i=0}^t L_i) \oplus K(L \oplus H)$ and $(\sum_{i=0}^{t'} M_i) \oplus K(M \oplus H)$ are canonical decompositions of $L \oplus H$ and $M \oplus H$ respectively, then $t = t' = T - I$ and $s(L_i) = s(M_i) = I + i$ for $0 \leq i \leq t$. Write

$$\begin{aligned} \text{Ord}(L, M, j) &= \min \{ \text{ord}(\beta^2 \pi^{-1} + \delta) \mid \beta^2 \pi^{-1} + \delta + \mathcal{P}k \\ &= \sum_{i=0}^{j-I-1} (\Delta(kL_i) + \Delta(kM_i)) \end{aligned}$$

for some canonical decompositions $(\sum_{i=0}^{T-I} L_i) \oplus K(L \oplus H)$ and $(\sum_{i=0}^{T-I} M_i) \oplus K(M \oplus H)$ of $L \oplus H$ and $M \oplus H$ respectively.

LEMMA 7.3. Let $L \cong \begin{pmatrix} \pi^s & \\ a & c \end{pmatrix}$ and M be an $s+1$ -modular lattice such that $\text{rank } M \geq 8$ and $\pi^2 q(L) \subseteq q(M)$. If $a + a' \in q(M)$, then $L \oplus M \cong \begin{pmatrix} \pi^s & \\ a' & c \end{pmatrix} \oplus M'$, where M' is $s+1$ -modular and $q(M') = q(M)$.

The proof is a straightforward manipulation of lattices.

LEMMA 7.4. Let s, w , and w' be nonnegative integers such that either $w' = w$ or $w' = w + 2$; and let e be a unit. For any $\chi \in X$ with $m > s$ and $\chi \begin{pmatrix} \pi^s & \\ e\pi^w & 0 \end{pmatrix} = 0$, $\chi \begin{pmatrix} \pi^{s+1} & \\ e\pi^{w'} & 0 \end{pmatrix} = 0$.

Proof. If $w' = w + 2$, use Lemma 3.3. If $w' = w$, write $\begin{pmatrix} \pi^s & \\ e\pi^w & 0 \end{pmatrix} \oplus \begin{pmatrix} \pi^{s+1} & \\ e\pi^{w'} & 0 \end{pmatrix} \cong \begin{pmatrix} \pi^s & \\ 0 & 0 \end{pmatrix} \oplus \begin{pmatrix} \pi^{s+1} & \\ e\pi^{w'} & 0 \end{pmatrix}$.

THEOREM 7.5. For lattices L and M , $L \cong M$ if and only if

- (1) L and M are of the same type (I, T, R) ;
- (2) For $I \leq i \leq T$, $q(L(i) \oplus H_i) = q(M(i) \oplus H_i)$. That is, the i th norm groups of L and M are the same.
- (3) $\chi(L) = \chi(M)$ for all $\chi \in X(2T - u(T) + 2)$;
- (4) If for some j with $I < j \leq T$, $u(j) = u(j-1) + 2$ and $v(j) = v(j-1) + 2$, then $\text{Ord}(L, M, j) \geq u(j) + v(j) - 2j - 1$;
- (5) If $K(L) \cong (d_1)$ and $K(M) \cong (d_2)$, then $d_1 \in d_2 k^2$.

Proof. Necessity. Condition (4) follows from [6, Lemma 5.2] and Proposition 7.2. The others hold trivially.

Sufficiency. Replacing L and M by $L \oplus (\sum_{i=I}^T H_i)$ and $M \oplus (\sum_{i=I}^T H_i)$ respectively, we may assume L and M are normal lattices. Proceed by induction on $n = \text{length } d = \text{length } M$.

Part I. $n = 1$. Then $T = I$. Write $u(I) = u$ and $v(I) = v$.

Case 1. $\text{Rank}(K(L)) = 0$. Assume $K(M) \neq 0$. Then $u = u_{K(M)}$ by normality; and for any unit e and any χ with $m = 2T - u + 1$, $\chi(e\pi^{u_{K(M)}}) = 0$. Thus $\chi(L) = \chi(M) = 0$ for such χ . But $\chi(L) = 0$ by Propositions 3.4 and 5.3. Therefore $K(M) = 0$. It now follows from

Theorem 5.4 that $L \cong M$.

Case 2. $\text{Rank}(K(L)) = 1$. By normality, $K(L) \cong (e\pi^u)$ for some unit e . Write $L \cong L_0 \oplus K(L)$ with $q(L_0) = e(\pi^u \mathfrak{p}^2 + \pi^v \mathfrak{p}^2)$. There exists $\chi \in X$ with $m = 2I - v + 3$ and $\chi(L) \neq 0$. Since $\chi(e\pi^v) = 0$, $\text{rank}(K(M)) \neq 2$. Using condition (5), $K(M) \cong (e\pi^u)$. Using [6, Lemma 4.5, Theorem 4.6, and $LT_0 d$], $L \cong \begin{pmatrix} \pi^I & \\ e\pi^v & e^{-1}\pi^{2I-v}\delta \end{pmatrix} \oplus H_I \oplus (e\pi^u)$ and $M \cong \begin{pmatrix} \pi^I & \\ e\pi^v & e^{-1}\pi^{2s-v}\delta' \end{pmatrix} \oplus H_I' \oplus (e\pi^u)$. By an argument already used on several occasions (see e.g., Theorem 5.4) $L \cong M$.

Case 3. $\text{Rank}(K(L)) = 2$. $K(L) \cong (e\pi^u) \oplus (e\pi^v)$ for some unit e . By an argument similar to the one given in Case 2, $L \cong H_I \oplus (e\pi^v) \oplus (e\pi^v) \cong M$.

Part II. Assume $n > 1$ and that the result holds for normal lattices of length $n - 1$. By [6, Lemma 5.3], L and M admit saturated decompositions $(\sum_{i=0}^{n-1} L_i) \oplus K(L)$ and $(\sum_{i=0}^{n-1} M_i) \oplus K(M)$ respectively. Our strategy is to show the following:

(7.1) For $j = 0, 1$, there exists $s(j)$ -modular lattices L_j^* and M_j^* with $q(L_j^*) = q(L_j)$ and $q(M_j^*) = q(M_j)$ such that $L_0 \oplus L_1 \cong L_0^* \oplus L_1^*$, $M_0 \oplus M_1 \cong M_0^* \oplus M_1^*$, and $L_0^* \cong M_0^*$.

Then $(L_0^*)^\perp$ and $(M_0^*)^\perp$ are normal lattices of length $n - 1$ which satisfy conditions (1), (2), (4), and (5). For any $\chi \in X(2T - u(T) + 2)$ such that $\chi(L_0^*) \neq 0$, $\chi((L_0^*)^\perp) = \chi((M_0^*)^\perp)$. If $\chi \in X(2T - u(T) + 2)$ and $\chi(L_0^*) = 0$, then using Lemma 7.4 and Proposition 5.3, $\chi((L_0^*)^\perp) = 0 = \chi((M_0^*)^\perp)$. Thus condition (3) is satisfied. The result follows easily by induction.

We verify (7.1) by cases. But first write $L_0 \cong L'_0 \oplus L''_0 \oplus H_I$ where $L'_0 \cong \begin{pmatrix} \pi^I & \\ e\pi^{u(I)} & e^{-1}\pi^{2I-u(I)}\delta + e^{-1}\pi^{v(I)}\alpha^2 \end{pmatrix} \oplus \begin{pmatrix} \pi^I & \\ e\pi^{v(I)} & e^{-1}\pi^{2I-v(I)}\delta' \end{pmatrix}$ and $L''_0 \cong \begin{pmatrix} \pi^I & \\ e\pi^{u(I)} & 0 \end{pmatrix} \oplus \begin{pmatrix} \pi^I & \\ e\pi^{v(I)} & 0 \end{pmatrix}$ with $\delta, \delta' \in \{0, \lambda\}$, $\alpha \in \mathfrak{o}$, and e a unit such that $q_{s(j)}(L) = e(\pi^{u(s(j))}\mathfrak{p}^2 + \pi^{v(s(j))}\mathfrak{p}^2)$ for $j = 0, 1$.

Case A. $v(I + 1) = v(I)$. By [6, Definition 5.1], $u(I + 1) = u(I)$ or $u(I) + 2$. Applying Lemma 7.3 twice, $L_0 \oplus L_1 \cong L_0^* \oplus L_1^*$ with $L_0^* \cong L''_0 \oplus H_I'$. M_0^* and M_1^* exist with $M_0^* \cong L''_0 \oplus H_I'$, and these lattices satisfy (7.1).

Case B. $v(I + 1) = v(I) + 1$. Then $u(I + 1) = v(I) = u(I) + 1$. If $u(I) < I$, the result follows as in Case A. Suppose $u(I) = I$. $L_0^* = L_0 \cong \begin{pmatrix} \pi^I & \\ \pi^I & \pi^I\delta \end{pmatrix} \oplus H_I'$ and $M_0^* \cong \begin{pmatrix} \pi^I & \\ \pi^I & \pi^I\delta_1 \end{pmatrix} \oplus H_I'$ and the result follows.

Case C. $v(I+1) = v(I) + 2$. If $v(I+1) \leq I+1$ and $u(I+1) = u(I)$, $L_0^* \cong L_0'' \oplus H_I'$.

If $v(I+1) \leq I+1$ and $u(I+1) = u(I) + 2$, write $a = a_1 + a_2$ with $a_1 \in \Omega$ and $a_2 \in \mathfrak{o}(1)$. In this case $L_0^* \cong \left(e\pi^{u(I)} \pi^I e^{-1}\pi^{v(I)} a_1^2 \right) \oplus L_0'' \oplus H_I'$ and $M_0^* \cong \left(e\pi^{u(I)} \pi^I e^{-1}\pi^{v(I)} \delta^2 \right) \oplus L_0'' \oplus H_I'$. By condition (4) $a_1 = c$.

We are left with the situation in which $v(I+1) = I+2$. If $u(I+1) = u(I)$, take $L_0^* \cong \left(e\pi^{v(I)} \pi^I e^{-1}\pi^{2I-v(I)} \delta' \right) \oplus L_0'' \oplus H_I'$ and $M_0^* \cong \left(e\pi^{v(I)} \pi^I e^{-1}\pi^{2s-v(I)} \delta_1 \right) \oplus L_0'' \oplus H_I'$. If $u(I+1) = u(I) + 2$, by the argument used in the case where $v(I+1) = v(I) + 2 \leq I+1$ and $u(I+1) = u(I) + 2$, $L_0^* \cong \left(e\pi^{u(I)} \pi^I e^{-1}\pi^{v(I)} a_1^2 \right) \oplus \left(e\pi^{v(I)} \pi^I e^{-1}\pi^{2I-v(I)} \delta' \right) \oplus L_0'' \oplus H_I$ and $M_0^* \cong \left(e\pi^{u(I)} \pi^I e^{-1}\pi^{v(I)} a_1^2 \right) \oplus \left(e\pi^{v(I)} \pi^I e^{-1}\pi^{2I-v(I)} \delta_1 \right) \oplus L_0'' \oplus H_I$ with $a_1 \in \Omega$. We need only show that $\delta' = \delta_1$. The difficulty here is in checking that there exists $\chi \in X$ with $m = I+1$ and $\chi(L) \neq 0$. Let r be the largest integer such that $0 \leq r \leq n-1$ and $u(I+r) < I+1$. Since $u(I) < I+1$, r exists. $u(I+r) + (I+1) \equiv 0 \pmod{2}$. So there exists χ with $m = I+1$ and $\chi \left(e_{s(r)} \pi^{u(s(r))} \pi^{s(r)} \right) \neq 0$. It follows that $\chi(L_r) \neq 0$ and by induction on r , $\chi(\sum_{i=0}^r L_i) \neq 0$. Thus $\chi(L) \neq 0$.

REFERENCES

1. C. Arf, *Untersuchungen über quadratische formen in körpern der charakteristik 2*, I. J. Reine Angew. Math., **183** (1940), 148-167.
2. ———, *Über arithmetische äquivalenz quadratischer formen in potenzreihenkörpern über einem vollkommen körper der charakteristik 2*, Rev. Fac. Sci. Univ. Istanbul, **8** (1943), 297-327.
3. R. Jacobowitz, *Gauss sums and the local classification of hermitian forms*, Amer. J. Math., **90** (1968), 528-552.
4. O. T. O'Meara, *Local characterization of integral quadratic forms by Gauss sums*, Amer. J. Math., **79** (1957), 687-709.
5. ———, *Introduction to Quadratic Forms*, Springer-Verlag, Berlin (1963).
6. C.-H. Sah, *Quadratic forms over fields of characteristic 2*, Amer. J. Math., **82** (1960), 812-830.

Received December 30, 1975 and in revised form February 18, 1976. The results of this paper are contained in the author's doctoral thesis. The author wishes to express her appreciation to her advisor, Professor Ronald Jacobowitz.

BEREA COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 65, No. 1

September, 1976

David Lee Armacost, <i>Compactly cogenerated LCA groups</i>	1
Sun Man Chang, <i>On continuous image averaging of probability measures</i>	13
J. Chidambaraswamy, <i>Generalized Dedekind ψ-functions with respect to a polynomial. II</i>	19
Freddy Delbaen, <i>The Dunford-Pettis property for certain uniform algebras</i>	29
Robert Benjamin Feinberg, <i>Faithful distributive modules over incidence algebras</i>	35
Paul Froeschl, <i>Chained rings</i>	47
John Brady Garnett and Anthony G. O'Farrell, <i>Sobolev approximation by a sum of subalgebras on the circle</i>	55
Hugh M. Hilden, José M. Montesinos and Thomas Lusk Thickstun, <i>Closed oriented 3-manifolds as 3-fold branched coverings of S^3 of special type</i>	65
Atsushi Inoue, <i>On a class of unbounded operator algebras</i>	77
Peter Kleinschmidt, <i>On facets with non-arbitrary shapes</i>	97
Narendrakumar Ramanlal Ladhawala, <i>Absolute summability of Walsh-Fourier series</i>	103
Howard Wilson Lambert, <i>Links which are unknottable by maps</i>	109
Kyung Bai Lee, <i>On certain g-first countable spaces</i>	113
Richard Ira Loeb, <i>A Hahn decomposition for linear maps</i>	119
Moshe Marcus and Victor Julius Mizel, <i>A characterization of non-linear functionals on W_1^p possessing autonomous kernels. I</i>	135
James Miller, <i>Subordinating factor sequences and convex functions of several variables</i>	159
Keith Pierce, <i>Amalgamated sums of abelian l-groups</i>	167
Jonathan Rosenberg, <i>The C^*-algebras of some real and p-adic solvable groups</i>	175
Hugo Rossi and Michele Vergne, <i>Group representations on Hilbert spaces defined in terms of ∂_b-cohomology on the Silov boundary of a Siegel domain</i>	193
Mary Elizabeth Schaps, <i>Nonsingular deformations of a determinantal scheme</i>	209
S. R. Singh, <i>Some convergence properties of the Bubnov-Galerkin method</i>	217
Peggy Strait, <i>Level crossing probabilities for a multi-parameter Brownian process</i>	223
Robert M. Tardiff, <i>Topologies for probabilistic metric spaces</i>	233
Benjamin Baxter Wells, Jr., <i>Rearrangements of functions on the ring of integers of a p-series field</i>	253
Robert Francis Wheeler, <i>Well-behaved and totally bounded approximate identities for $C_0(X)$</i>	261
Delores Arletta Williams, <i>Gauss sums and integral quadratic forms over local fields of characteristic 2</i>	271
John Yuan, <i>On the construction of one-parameter semigroups in topological semigroups</i>	285