

Pacific Journal of Mathematics

**FIXED POINTS OF LOCALLY CONTRACTIVE AND
NONEXPANSIVE SET-VALUED MAPPINGS**

PETER K. F. KUHFITIG

FIXED POINTS OF LOCALLY CONTRACTIVE AND NONEXPANSIVE SET-VALUED MAPPINGS

PETER K. F. KUHFITTING

Let (M, d) be a complete metric space and $S(M)$ the set of all nonempty bounded closed subsets of M . A set-valued mapping $f: M \rightarrow S(M)$ will be called (uniformly) *locally contractive* if there exist ε and λ ($\varepsilon > 0, 0 < \lambda < 1$) such that $D(f(x), f(y)) \leq \lambda d(x, y)$ whenever $d(x, y) < \varepsilon$ and where $D(f(x), f(y))$ is the distance between $f(x)$ and $f(y)$ in the Hausdorff metric induced by d on $S(M)$. It is shown in the first theorem that if M is "well-chained," then f has a fixed point, that is, a point $x \in M$ such that $x \in f(x)$. This fact, in turn, yields a fixed-point theorem for locally nonexpansive set-valued mappings on a compact star-shaped subset of a Banach space. Both theorems are extensions of earlier results.

1. **Locally contractive set-valued mappings.** Following Assad and Kirk [1] we shall define Z as follows: if $r > 0$ and $Y \in S(M)$, let

$$Z(r, Y) = \{x \in M: \text{dist}(x, Y) < r\}.$$

Then for $A, B \in S(M)$ we define

$$D(A, B) = \inf \{r: A \subset Z(r, B) \text{ and } B \subset Z(r, A)\}.$$

Also noted in [1] are two lemmas:

LEMMA 1. *If $A, B \in S(M)$ and $x \in A$, then for each positive number α there exists $y \in B$ such that*

$$d(x, y) \leq D(A, B) + \alpha.$$

LEMMA 2. *Let $\{X_n\}$ be a sequence of sets in $S(M)$, and assume that $\lim_{n \rightarrow \infty} D(X_n, X_0) = 0$ ($X_0 \in S(M)$). Then if $x_n \in X_n$ ($n = 1, 2, \dots$) and $\lim_{n \rightarrow \infty} x_n = x_0$, it follows that $x_0 \in X_0$.*

Finally, suppose M is well-chained in the sense that for every $\varepsilon > 0$ and $x, y \in M$ there exists an ε -chain, that is, a finite set of points

$$x = y_0, y_1, \dots, y_n = z$$

(n may depend on both x and z) such that $d(y_i, y_{i+1}) < \varepsilon$ ($i = 0, 1, \dots, n - 1$).

THEOREM 1. *Suppose (M, d) is a complete well-chained metric space and $S(M)$ the set of all nonempty bounded closed subsets of M . If $f: M \rightarrow S(M)$ is locally contractive, then f has a fixed point.*

Proof. Assume that $\varepsilon < 1$ and let $x_0, y_0 \in M$ such that $d(x_0, y_0) < \varepsilon$. Then

$$D(f(x_0), f(y_0)) \leq \lambda d(x_0, y_0).$$

Now choose a positive number $\eta < \varepsilon - \lambda\varepsilon < 1$. Let x_1 be any element in $f(x_0)$; then there exists by Lemma 1 an element $y_1 \in f(y_0)$ such that

$$d(x_1, y_1) \leq D(f(x_0), f(y_0)) + \eta.$$

Hence

$$d(x_1, y_1) < \lambda\varepsilon + \eta < \lambda\varepsilon + \varepsilon - \lambda\varepsilon = \varepsilon.$$

Next, let $x_2 \in f(x_1)$; then there exists $y_2 \in f(y_1)$ such that

$$\begin{aligned} d(x_2, y_2) &\leq D(f(x_1), f(y_1)) + \eta^2 \\ &\leq \lambda d(x_1, y_1) + \eta^2. \end{aligned}$$

In general, for $n > 0$

$$d(x_n, y_n) \leq D(f(x_{n-1}), f(y_{n-1})) + \eta^n,$$

and we can show by induction that

$$(1) \quad d(x_n, y_n) < \lambda^n \varepsilon + \lambda^{n-1} \eta + \lambda^{n-2} \eta^2 + \cdots + \eta^n.$$

Indeed,

$$\begin{aligned} &\lambda^n \varepsilon + \lambda^{n-1} \eta + \lambda^{n-2} \eta^2 + \cdots + \eta^n \\ &< \lambda^n \varepsilon + \lambda^{n-1} (\varepsilon - \lambda\varepsilon) + \lambda^{n-2} (\varepsilon - \lambda\varepsilon)^2 + \cdots + (\varepsilon - \lambda\varepsilon)^n \\ &\leq \lambda^n \varepsilon + \lambda^{n-1} (\varepsilon - \lambda\varepsilon) + \lambda^{n-2} (\varepsilon - \lambda\varepsilon) + \cdots + (\varepsilon - \lambda\varepsilon) \\ &= \lambda^n \varepsilon + (\lambda^{n-1} \varepsilon - \lambda^n \varepsilon) + (\lambda^{n-2} \varepsilon - \lambda^{n-1} \varepsilon) + \cdots + (\varepsilon - \lambda\varepsilon) \\ &= \varepsilon. \end{aligned}$$

So if (1) is valid for $n = N > 0$, let $x_{N+1} \in f(x_N)$; then there exists $y_{N+1} \in f(y_N)$ such that

$$\begin{aligned} d(x_{N+1}, y_{N+1}) &\leq D(f(y_N), f(y_N)) + \eta^{N+1} \leq \lambda d(x_N, y_N) + \eta^{N+1} \\ &< \lambda(\lambda^N \varepsilon + \lambda^{N-1} \eta + \lambda^{N-2} \eta^2 + \cdots + \eta^N) + \eta^{N+1} \\ &= \lambda^{N+1} \varepsilon + \lambda^N \eta + \lambda^{N-1} \eta^2 + \cdots + \lambda \eta^N + \eta^{N+1}. \end{aligned}$$

Using this information we now construct a sequence in M as follows: let $y_{0,0}$ be an arbitrary element in M and let $y_{1,0} \in f(y_{0,0})$.

Consider the ε -chain

$$y_{0,0}, y_{0,1}, \dots, y_{0,n} = y_{1,0} \in f(y_{0,0}),$$

so that $d(y_{0,i}, y_{0,i+1}) < \varepsilon$ ($i = 0, 1, \dots, n - 1$). Since $y_{1,0} \in f(y_{0,0})$, we may choose $y_{1,1} \in f(y_{0,1})$ such that

$$(2) \quad d(y_{1,0}, y_{1,1}) \leq D(f(y_{0,0}), f(y_{0,1})) + \eta.$$

Similarly, since $y_{1,1} \in f(y_{0,1})$, choose $y_{1,2} \in f(y_{0,2})$ such that

$$d(y_{1,1}, y_{1,2}) \leq D(f(y_{0,1}), f(y_{0,2})) + \eta.$$

Continuing along the ε -chain, since $y_{1,n-1} \in f(y_{0,n-1})$, there exists $y_{1,n} = y_{2,0} \in f(y_{0,n})$ (i.e., $y_{2,0} \in f(y_{1,0})$) such that

$$d(y_{1,n-1}, y_{1,n}) \leq D(f(y_{0,n-1}), f(y_{0,n})) + \eta.$$

Consequently,

$$d(y_{1,0}, y_{2,0}) = d(y_{1,0}, y_{1,n}) \leq \sum_{i=0}^{n-1} d(y_{1,i}, y_{1,i+1}) < n(\lambda\varepsilon + \eta).$$

Next, referring to (2), since $y_{2,0} \in f(y_{1,0})$, there exists $y_{2,1} \in f(y_{1,1})$ for which

$$d(y_{2,0}, y_{2,1}) \leq D(f(y_{1,0}), f(y_{1,1})) + \eta^2,$$

and for $y_{2,n-1} \in f(y_{1,n-1})$, we have $y_{2,n} = y_{3,0} \in f(y_{1,n})$ (i.e., $y_{3,0} \in f(y_{2,0})$) such that

$$d(y_{2,n-1}, y_{2,n}) \leq D(f(y_{1,n-1}), f(y_{1,n})) + \eta^2.$$

Proceeding in this manner, and making use of (1), we get (for $m > 0$)

$$d(y_{m,l}, y_{m,l+1}) < \lambda^m\varepsilon + \lambda^{m-1}\eta + \lambda^{m-2}\eta^2 + \dots + \eta^m$$

($l = 0, 1, \dots, n - 1$). Now let $z_m = y_{m,0}$, so that $z_m \in f(z_{m-1})$, $m = 1, 2, \dots$, and $z_{m+1} = y_{m+1,0} = y_{m,n}$. Then

$$\begin{aligned} d(z_m, z_{m+1}) &\leq \sum_{l=0}^{n-1} d(y_{m,l}, y_{m,l+1}) \\ &< n(\lambda^m\varepsilon + \lambda^{m-1}\eta + \lambda^{m-2}\eta^2 + \dots + \eta^m). \end{aligned}$$

To show that $\{z_m\}$ is a Cauchy sequence, let $\beta = \max(\lambda, \eta)$. Then

$$d(z_m, z_{m+1}) < n(m + 1)\beta^m,$$

and for $0 < i < j$

$$\begin{aligned}
 d(z_i, z_j) &\leq \sum_{k=i}^{j-1} d(z_k, z_{k+1}) \\
 &< n \sum_{k=i}^{j-1} (k+1)\beta^k \\
 &\leq n \sum_{k=i}^{\infty} (k+1)\beta^k .
 \end{aligned}$$

It is easily checked that $d(z_i, z_j) \rightarrow 0$ as $i \rightarrow \infty$, implying that $\{z_m\}$ is a Cauchy sequence, which converges to some $z \in M$ by the completeness of M .

Finally, since $z_m \in f(z_{m-1})$ and $z_m \rightarrow z$, $f(z_{m-1}) \rightarrow f(z)$ and, by Lemma 2, $z \in f(z)$.

REMARK 1. Nadler [4] proved a similar theorem by a different method under the additional assumption that each $f(x)$ is compact.

2. **Locally nonexpansive set-valued mappings.** Let X be a Banach space and C a subset of X . A mapping $T: C \rightarrow S(C)$ will be called *locally nonexpansive* if there exists $\varepsilon > 0$ such that

$$D(Tx, Ty) \leq \|x - y\| ,$$

whenever $\|x - y\| < \varepsilon$ and where D is again the distance in the Hausdorff metric induced by d on $S(M)$ (as usual, $d(x, y) = \|x - y\|$ for all $x, y \in X$).

THEOREM 2. *Let X be a Banach space and C a compact star-shaped subset of X . If $T: C \rightarrow S(C)$ is locally nonexpansive, then there exists a point $x \in C$ such that $x \in Tx$.*

Proof. Let c be the star-center of C and let $\{k_n\}$ be an increasing sequence of real numbers converging to 1. Define $U_n: C \rightarrow S(C)$ by

$$U_n x = (1 - k_n)c + k_n T x ,$$

where $k_n T x = \{k_n y: y \in T x\}$. Let $z, y \in C$ such that $\|z - y\| < \varepsilon$. Then $D(Tz, Ty) \leq \|z - y\|$. Now for any two elements $z' \in Tz$ and $y' \in Ty$

$$\|(1 - k_n)c + k_n z' - (1 - k_n)c - k_n y'\| = k_n \|z' - y'\| .$$

Hence

$$D(U_n z, U_n y) \leq k_n \|z - y\| .$$

Consequently, U_n has a fixed point $x_n \in C$ by Theorem 1. Since C is

compact, there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ converging to some $x \in C$, and because T is continuous,

$$Tx_{n_i} \longrightarrow Tx .$$

Now

$$\begin{aligned} \text{dist}(x_{n_i}, Tx_{n_i}) &\leq D(U_{n_i}x_{n_i}, Tx_{n_i}) \\ &= D((1 - k_{n_i})c + k_{n_i}Tx_{n_i}, Tx_{n_i}) \longrightarrow D(Tx, Tx) \text{ as } i \longrightarrow \infty . \end{aligned}$$

Thus

$$\text{dist}(x, Tx) = 0 ,$$

which implies that $x \in Tx$, Tx being closed.

Theorem 2 and its point-to-point analogue generalize an earlier theorem due to Dotson [2]:

COROLLARY. *A nonexpansive self-mapping of a compact star-shaped subset of a Banach space has a fixed point.*

REMARK 2. Edelstein [3] has shown that a locally contractive (nonexpansive) point-to-point mapping need not be globally contractive (nonexpansive). On convex sets, however, a locally nonexpansive mapping is nonexpansive.

REFERENCES

1. N. A. Assad and W. A. Kirk, *Fixed point theorems for set-valued mappings of contractive type*, Pacific J. Math., **43** (1972), 553-562.
2. W. G. Dotson, Jr., *Fixed point theorems for nonexpansive mappings on star-shaped subsets of Banach spaces*, J. London Math. Soc., (2) **4** (1972), 408-410.
3. M. Edelstein, *An extension of Banach's contraction principle*, Proc. Amer. Math. Soc., **12** (1961), 7-10.
4. S. B. Nadler, Jr., *Multi-valued contraction mappings*, Pacific J. Math., **30** (1969), 475-488.

Received August 12, 1975.

MILWAUKEE SCHOOL OF ENGINEERING

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)

University of California
Los Angeles, California 90024

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT

University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM

Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of your manuscript. You may however, use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.),
8-8, 3-chome, Takadanobaba, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1975 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 65, No. 2

October, 1976

Andrew Adler, <i>Weak homomorphisms and invariants: an example</i>	293
Howard Anton and William J. Pervin, <i>Separation axioms and metric-like functions</i>	299
Ron C. Blei, <i>Sidon partitions and p-Sidon sets</i>	307
T. J. Cheatham and J. R. Smith, <i>Regular and semisimple modules</i>	315
Charles Edward Cleaver, <i>Packing spheres in Orlicz spaces</i>	325
Le Baron O. Ferguson and Michael D. Rusk, <i>Korovkin sets for an operator on a space of continuous functions</i>	337
Rudolf Fritsch, <i>An approximation theorem for maps into Kan fibrations</i>	347
David Sexton Gilliam, <i>Geometry and the Radon-Nikodym theorem in strict Mackey convergence spaces</i>	353
William Hery, <i>Maximal ideals in algebras of topological algebra valued functions</i>	365
Alan Hopenwasser, <i>The radical of a reflexive operator algebra</i>	375
Bruno Kramm, <i>A characterization of Riemann algebras</i>	393
Peter K. F. Kuhfittig, <i>Fixed points of locally contractive and nonexpansive set-valued mappings</i>	399
Stephen Allan McGrath, <i>On almost everywhere convergence of Abel means of contraction semigroups</i>	405
Edward Peter Merkes and Marion Wetzel, <i>A geometric characterization of indeterminate moment sequences</i>	409
John C. Morgan, II, <i>The absolute Baire property</i>	421
Eli Aaron Passow and John A. Roulier, <i>Negative theorems on generalized convex approximation</i>	437
Louis Jackson Ratliff, Jr., <i>A theorem on prime divisors of zero and characterizations of unmixed local domains</i>	449
Ellen Elizabeth Reed, <i>A class of T_1-compactifications</i>	471
Maxwell Alexander Rosenlicht, <i>On Liouville's theory of elementary functions</i>	485
Arthur Argyle Sagle, <i>Power-associative algebras and Riemannian connections</i>	493
Chester Cornelius Seabury, <i>On extending regular holomorphic maps from Stein manifolds</i>	499
Elias Sai Wan Shiu, <i>Commutators and numerical ranges of powers of operators</i>	517
Donald Mark Topkis, <i>The structure of sublattices of the product of n lattices</i>	525
John Bason Wagoner, <i>Delooping the continuous K-theory of a valuation ring</i>	533
Ronson Joseph Warne, <i>Standard regular semigroups</i>	539
Anthony William Wickstead, <i>The centraliser of $E \otimes_{\lambda} F$</i>	563
R. Grant Woods, <i>Characterizations of some C^*-embedded subspaces of $\beta\mathbb{N}$</i>	573