

Pacific Journal of Mathematics

**WEIERSTRASS POINTS OF PRODUCTS OF RIEMANN
SURFACES**

ROBERT F. LAX

WEIERSTRASS POINTS OF PRODUCTS OF RIEMANN SURFACES

R. F. LAX

Ogawa has defined sets of Weierstrass points of a holomorphic vector bundle on a compact complex manifold. We generate nontrivial examples of such sets of Weierstrass points by considering the canonical bundle on a product of Riemann surfaces.

In the first section, we review Ogawa's definition and some classical facts about Weierstrass points on Riemann surfaces. In §2, we prove our theorems and consider an example to illustrate the proofs. Finally, we remark that a connection between Weierstrass points on a Riemann surface and fixed points of a periodic automorphism does not seem to extend to higher dimensions.

We wish to thank Pierre Conner for helpful conversations and Roy Ogawa for useful communications.

1. Let M denote a connected, compact complex manifold of (complex) dimension n . Let E denote a holomorphic vector bundle on M of rank q . Let $J^k(E)$, $k = 0, 1, \dots$, denote the holomorphic vector bundle of k -jets of E (cf. [7]). Put $R_k = \text{rank } J^k(E) = q \cdot (n+k)!/n!k!$. Suppose that $\Gamma(E)$, the vector space of global holomorphic sections of E , is of dimension $d > 0$. Consider the trivial bundle $M \times \Gamma(E)$ and the map

$$j_k : M \times \Gamma(E) \rightarrow J^k(E)$$

which at a point $P \in M$ takes a section to its k -jet at P . Put $\mu = \min(d, R_k)$.

DEFINITION. (cf. [6, 3]). For $1 \leq r \leq \mu$, let $W'_k(E)$ denote the reduced closed analytic subspace of M defined by the vanishing of the exterior power $\Lambda^{\mu-r+1} j_k$.

The points of $W'_k(E)$ are those $P \in M$ such that the rank of $j_{k,P}$ is at most $\mu - r$.

PROPOSITION 1. *Either $W'_k(E)$ is empty or each component has codimension at most $r(|d - R_k| + r)$ in M .*

Proof. [2, Proposition 4].

Next, we need to review some facts from the classical theory of

Weierstrass points. We refer the reader to [1] for details. Let C denote a compact Riemann surface of genus $g > 1$ and let P be a point on C . Let t denote a local coordinate at P on C (so $t(P) = 0$). Suppose the sequence of gaps at P is $1, s_2, s_3, \dots, s_g$. Then we may choose a basis $\omega_1, \dots, \omega_g$ of holomorphic differentials on C such that, writing $\omega_j = f_j(t)dt$ locally at P , we have that $f_1(0) = 1$ and the order of f_j at P is $s_j - 1$ for $j = 2, \dots, g$. We will call such a basis of holomorphic differentials *special with respect to P* .

Let K denote the canonical bundle on C . Then the matrix of the map $j_k: C \times \Gamma(K) \rightarrow J^k(K)$ locally at P with respect to the above basis $\{\omega_j\}$ is

$$[f_j^{(i)}(t)] \begin{array}{l} i = 0, \dots, k \\ j = 1, \dots, g \end{array}$$

where $f_j^{(i)}(t)$ denotes the i^{th} derivative of f_j with respect to t . Note that, by our choice of basis, when we evaluate this matrix at P we get a lower triangular matrix. The next proposition follows easily from the form of this matrix and the choice of our basis $\{\omega_j\}$.

PROPOSITION 2. *Suppose $k \leq g - 1$. Then $P \in W_k^1(K)$ if and only if $s_j > j$ for some $j = 2, 3, \dots, k + 1$.*

2. Let C_i be a compact Riemann surface of genus $g_i > 1$, $i = 1, \dots, n$. Denote by K_i the canonical bundle on C_i . Put $X = C_1 \times C_2 \times \dots \times C_n$ and let K denote the canonical bundle on X . Then X is a connected, compact complex manifold of dimension n and $\dim_C \Gamma(K) = \prod_{i=1}^n g_i$. Put $R_k = \text{rank } J^k(K)$.

THEOREM 1. *Suppose $k \leq \min_{1 \leq i \leq n} \{g_i - 1\}$. Let $\mathbf{P} = (P_1, \dots, P_n) \in X$. Then $\mathbf{P} \in W_k^1(K)$ if and only if $P_i \in W_k^1(K_i)$ for some $i = 1, \dots, n$.*

Proof. The notation in the general case is very complicated. We will prove the theorem for the case $n = 2$. It is not hard to see that the general case may be demonstrated by a completely similar argument with no new ideas necessary.

So, let C and D be compact Riemann surfaces of genera $g > 1$ and $h > 1$ respectively, and suppose, without loss of generality, that g is greater than or equal to h . Let K_C (resp. K_D) denote the canonical bundle on C (resp. D). Put $X = C \times D$ and let $(P, Q) \in X$. Let t (resp. u) denote a local coordinate at P on C (resp. at Q on D). Let $\alpha_i = \varphi_i(t)dt$, $i = 1, \dots, g$, denote the basis of holomorphic differentials on

C special with respect to P and let $\beta_j = \psi_j(u)du$, $j = 1, \dots, h$, denote the basis of holomorphic differentials on D special with respect to Q .

Let $\pi_1: X \rightarrow C$ and $\pi_2: X \rightarrow D$ denote the respective projection maps. Put

$$\omega_{ij} = \pi_1^* \alpha_i \wedge \pi_2^* \beta_j \quad \begin{array}{l} i = 1, \dots, g \\ j = 1, \dots, h. \end{array}$$

Then the ω_{ij} are a basis of holomorphic 2-forms on X and locally at (P, Q) we may write $\omega_{ij} = \varphi_i(t)\psi_j(u)dt \wedge du$.

Let K denote the canonical bundle on X and suppose $0 \leq k \leq h - 1$. (Note then that $R_k < h^2 \leq gh$.) Consider the map $j_k: X \times \Gamma(K) \rightarrow J^k(K)$. Denote by $D^{l,m}$ the differential operator $\partial^{l+m}/\partial t^l \partial u^m$. The entries of the matrix of j_k locally at (P, Q) are then $D^{l,m}(\varphi_i(t)\psi_j(u))$, where $1 \leq i \leq g$, $1 \leq j \leq h$, and where l, m are nonnegative integers such that $l + m \leq k$. It is not hard to see that after suitably ordering the basis elements of $\Gamma(K)$ and $J_{(P,Q)}^k(K)$ the matrix of j_k when evaluated at (P, Q) is a lower triangular matrix with diagonal entries

$$D^{l,m}(\varphi_{i+1}(t)\psi_{m+1}(u))|_{(0,0)} = \varphi_{i+1}^{(l)}(0) \cdot \psi_{m+1}^{(m)}(0).$$

More precisely, we order the operators $D^{l,m}$ "lexicographically in each degree"; i.e. $D^{l,m}$ comes before $D^{l',m'}$ if $l + m < l' + m'$ or if $l + m = l' + m'$ and $l > l'$. Similarly, ω_{ij} comes before $\omega_{i'j'}$ if $i + j < i' + j'$ or if $i + j = i' + j'$ and $i > i'$.

Now, the rank of this matrix at (P, Q) is less than R_k (the maximum possible rank) if and only if $\varphi_{i+1}^{(l)}(0) = 0$ for some $l = 0, 1, \dots, k$ or $\psi_{m+1}^{(m)}(0) = 0$ for some $m = 0, 1, \dots, k$. But, by Proposition 2, $\varphi_{i+1}^{(l)}(0) = 0$ for some $l, 0 \leq l \leq k$, if and only if $P \in W_k^1(K_C)$ and $\psi_{m+1}^{(m)}(0) = 0$ for some $m, 0 \leq m \leq k$, if and only if $Q \in W_k^1(K_D)$. Hence $(P, Q) \in W_k^1(K)$ if and only if $P \in W_k^1(K_C)$ or $Q \in W_k^1(K_D)$.

THEOREM 2. *With the notation of Theorem 1, suppose $k > \min_{1 \leq i \leq n} \{g_i - 1\}$ and $R_k \leq \prod_{i=1}^n g_i$. Then $W_k^1(K) = X$.*

Proof. Again, we will prove this only for $n = 2$. With notation as in the proof of Theorem 1, we have that the matrix of j_k when evaluated at (P, Q) will be a lower triangular matrix with a term of the form

$$D^{0,k}(\varphi_i(t)\psi_h(u))|_{(0,0)} = \varphi_i(0)\psi_h^{(k)}(0)$$

with $l > 1$, as the last entry on the diagonal. But $\varphi_i(0) = 0$ for $l > 1$, so the mapping j_k fails to have maximal rank at every point of X . Hence $W_k^1(K) = X$.

To illustrate the proofs of the above theorems, we consider the following example. With notation as in Theorem 1, we suppose $g = h = 4$. We order the basis $\{\omega_{ij}\}$ of holomorphic 2-forms on X as follows: $\omega_{11}, \omega_{21}, \omega_{12}, \omega_{31}, \omega_{22}, \omega_{13}, \omega_{41}, \omega_{32}, \omega_{23}, \omega_{14}, \omega_{42}, \omega_{33}, \omega_{24}, \omega_{43}, \omega_{34}, \omega_{44}$. The matrix of j_4 evaluated at (P, Q) is a 15×16 lower triangular matrix with diagonal entries: $1, \varphi'_2(0), \psi'_2(0), \varphi''_3(0), \varphi'_2(0)\psi'_2(0), \psi''_3(0), \varphi'''_4(0), \varphi''_3(0)\psi'_2(0), \varphi'_2(0)\psi''_3(0), \psi'''_4(0), 0, 0, 0, 0, 0$.

It is then clear that the conclusions of Theorems 1 and 2 hold.

3. Let C be a compact Riemann surface of genus $g > 1$. Let σ be a periodic automorphism (conformal homeomorphism) of C of order n . Put $C^* = C/\langle\sigma\rangle$ and let g^* denote the genus of C^* . In [8], Schoeneberg proves the following theorem (also cf. [4]):

THEOREM. *Let P denote a fixed point of σ . Then P is a Weierstrass point of C if $g^* \neq [g/n]$, where $[x]$ denotes the greatest integer in x .*

We remark here that this result does not seem to generalize to higher dimensions. Indeed, consider $C \times C$, an algebraic manifold of geometric genus g^2 . Let $C(2)$ denote the second symmetric product of C with itself; i.e. $C(2) = C \times C/S_2$. Then $C(2)$ is an algebraic manifold of geometric genus $g(g-1)/2$ [5]. Note that $g(g-1)/2 < [g^2/2]$. Now, the set of fixed points of $C \times C$ under the action of S_2 is the diagonal, while, by Theorem 1, the nontrivial set of Weierstrass points of the canonical bundle on $C \times C$, the set $W_{g-1}^1(K_{C \times C})$, consists of all points (P, Q) such that either P or Q is a Weierstrass point of C . Thus not all fixed points are Weierstrass points in this case. We do not see any good way of generalizing Schoeneberg's Theorem to higher dimensions.

REFERENCES

1. R. C. Gunning, *Lectures on Riemann surfaces*, Princeton Math. Notes, Princeton Univ. Press, Princeton, N. J., 1966.
2. R. F. Lax, *On the dimension of varieties of special divisors*, Trans. Amer. Math. Soc., **203** (1975), 141-159.
3. ———, *On the existence of Ogawa's Weierstrass points*, J. London Math. Soc., (to appear).
4. J. Lewittes, *Automorphisms of compact Riemann surfaces*, Amer. J. Math., **85** (1963), 734-752.
5. I. G. MacDonald, *Symmetric products of an algebraic curve*, Topology, **1** (1962), 319-343.
6. R. H. Ogawa, *On the points of Weierstrass in dimensions greater than one*, Trans. Amer. Math. Soc., **184** (1973), 401-417.
7. R. S. Palais, *Seminar on the Atiyah-Singer Index Theorem*, Ann. of Math. Studies 57, Princeton Univ. Press, Princeton, N. J., 1965.
8. B. Schoeneberg, *Über die Weierstrasspunkte in den Körpern der elliptischen Modulfunktionen*, Abh. Math. Sem. Univ. Hamburg, **17** (1951), 104-111.

Received April 9, 1976. Partially supported by NSF Grant No. MPS 75-07552.

Pacific Journal of Mathematics

Vol. 66, No. 1

November, 1976

Helen Elizabeth. Adams, <i>Factorization-prime ideals in integral domains</i>	1
Patrick Robert Ahern and Robert Bruce Schneider, <i>The boundary behavior of Henkin's kernel</i>	9
Daniel D. Anderson, Jacob R. Matijevic and Warren Douglas Nichols, <i>The Krull intersection theorem. II</i>	15
Efraim Pacillas Armendariz, <i>On semiprime P.I.-algebras over commutative regular rings</i>	23
Robert H. Bird and Charles John Parry, <i>Integral bases for bicyclic biquadratic fields over quadratic subfields</i>	29
Tae Ho Choe and Young Hee Hong, <i>Extensions of completely regular ordered spaces</i>	37
John Dauns, <i>Generalized monoform and quasi injective modules</i>	49
F. S. De Blasi, <i>On the differentiability of multifunctions</i>	67
Paul M. Eakin, Jr. and Avinash Madhav Sathaye, <i>R-endomorphisms of $R[[X]]$ are essentially continuous</i>	83
Larry Quin Eifler, <i>Open mapping theorems for probability measures on metric spaces</i>	89
Garret J. Etgen and James Pawlowski, <i>Oscillation criteria for second order self adjoint differential systems</i>	99
Ronald Fintushel, <i>Local S^1 actions on 3-manifolds</i>	111
Kenneth R. Goodearl, <i>Choquet simplexes and σ-convex faces</i>	119
John R. Graef, <i>Some nonoscillation criteria for higher order nonlinear differential equations</i>	125
Charles Henry Heiberg, <i>Norms of powers of absolutely convergent Fourier series: an example</i>	131
Les Andrew Karlovitz, <i>Existence of fixed points of nonexpansive mappings in a space without normal structure</i>	153
Gangaram S. Ladde, <i>Systems of functional differential inequalities and functional differential systems</i>	161
Joseph Michael Lambert, <i>Conditions for simultaneous approximation and interpolation with norm preservation in $C[a, b]$</i>	173
Ernest Paul Lane, <i>Insertion of a continuous function</i>	181
Robert F. Lax, <i>Weierstrass points of products of Riemann surfaces</i>	191
Dan McCord, <i>An estimate of the Nielsen number and an example concerning the Lefschetz fixed point theorem</i>	195
Paul Milnes and John Sydney Pym, <i>Counterexample in the theory of continuous functions on topological groups</i>	205
Peter Johanna I. M. De Paepe, <i>Homomorphism spaces of algebras of holomorphic functions</i>	211
Judith Ann Palagallo, <i>A representation of additive functionals on L^p-spaces, $0 < p < 1$</i>	221
S. M. Patel, <i>On generalized numerical ranges</i>	235
Thomas Thornton Read, <i>A limit-point criterion for expressions with oscillatory coefficients</i>	243
Elemer E. Rosinger, <i>Division of distributions</i>	257
Peter S. Shoenfeld, <i>Highly proximal and generalized almost finite extensions of minimal sets</i>	265
R. Sirois-Dumais and Stephen Willard, <i>Quotient-universal sequential spaces</i>	281
Robert Charles Thompson, <i>Convex and concave functions of singular values of matrix sums</i>	285
Edward D. Tymchatyn, <i>Some n-arc theorems</i>	291
Jang-Mei Gloria Wu, <i>Variation of Green's potential</i>	295