

Pacific Journal of Mathematics

QUOTIENT-UNIVERSAL SEQUENTIAL SPACES

R. SIROIS-DUMAIS AND STEPHEN WILLARD

QUOTIENT-UNIVERSAL SEQUENTIAL SPACES

R. SIROIS-DUMAIS AND S. WILLARD

We produce 2^c mutually nonhomeomorphic countable sequential spaces. These are used

(1) to answer in the negative the following question of Michael and Stone [4]: is every regular T_1 -space which is a quotient of some separable metric space and a continuous image of the space \mathbf{P} of irrationals a quotient of \mathbf{P} ?

(2) to characterize c (with or without the continuum hypothesis) as the smallest cardinal κ with the property that a metric space of cardinality κ exists of which every sequential space of cardinality $\leq \kappa$ is a quotient.

1. Introduction. We let Q denote the space of rationals, \mathbf{P} the space of irrationals, \mathbf{R} the real line, and c the cardinality of \mathbf{R} . For any set X , the cardinality of X is denoted $|X|$.

We begin with the basic construction, which will be applied in the sequel in two different directions. Denote by Y the set $[Q \times (Q - \{0\})] \cup \{\infty\}$ and, for $E \subseteq \mathbf{R}$, denote by τ_E the quotient topology induced on Y by the obvious map from the subspace $[Q \times (Q - \{0\})] \cup (E \times \{0\})$ of $\mathbf{R} \times \mathbf{R}$. The set Y endowed with the topology τ_E will be denoted Y_E . Note that Y_E is a countable, regular, T_1 -space which is, by construction, the quotient of a separable metric space. (Thus, see [3], Y_E is both an \aleph_0 -space and a k -space.)

2. Quotients of \mathbf{P} . In [4], Michael and Stone establish that every metrizable continuous image of \mathbf{P} is a quotient of \mathbf{P} . The question is raised there whether this result can be extended to nonmetrizable images of \mathbf{P} , that is, whether a regular T_1 -space which is at the same time a quotient of some separable metric space and a continuous image of \mathbf{P} must be a quotient of \mathbf{P} . The construction of §1 provides the negative answer. To see this, first note that the countable discrete space (hence, every countable space) is a continuous image of \mathbf{P} (collapse each interval $(n, n + 1)$ to a point). It follows that each space Y_E is a regular T_1 -space which is a continuous image of \mathbf{P} and a quotient of some separable metric space. But:

THEOREM. *Not every space Y_E is a quotient of \mathbf{P} .*

Proof. If E and F are distinct subsets of \mathbf{R} , the topologies τ_E and τ_F on Y are different, one containing a set containing ∞ which does not belong to the other.

Now let S be the set of all surjections $f: \mathbf{P} \rightarrow Y$ such that each $f^{-1}(y)$, $y \in Y$, is closed in \mathbf{P} , and let Φ be the set of all $\phi: Y \rightarrow 2^{\mathbf{P}}$, where $2^{\mathbf{P}}$ denotes the collection of closed subsets of \mathbf{P} . Then $f \rightarrow f^{-1}$ is a one-one map from S into Φ ; since $|\Phi| = \mathfrak{c}^{\mathfrak{c}} = \mathfrak{c}$, we have $|S| \leq \mathfrak{c}$. Let J be the set of all T_1 topologies τ on Y such that (Y, τ) is a quotient image of \mathbf{P} . Then each $\tau \in J$ is generated by some $f \in S$, so $|J| \leq \mathfrak{c}$. Since $|\{\tau_E \mid E \subseteq \mathbf{R}\}| = 2^{\mathfrak{c}}$, and since each τ_E is T_1 , it follows that (Y, τ_E) is not a quotient of \mathbf{P} for some $E \subseteq \mathbf{R}$.

NOTES. (1) From the above, it is easily seen that there are $2^{\mathfrak{c}}$ nonhomeomorphic spaces Y_E , at most \mathfrak{c} of which can be quotients of \mathbf{P} . This result can be sharpened, with some difficulty. In fact, Y_E is a quotient of \mathbf{P} iff E is an analytic subset of \mathbf{R} .

(2) If, in the construction of Y , the set $Q \times (Q - \{0\})$ is replaced by a discrete space, say $\{(k/n, 1/n) \mid k, n \in \mathbf{N}\}$, the spaces Y_E which result still work, and have now the additional property that each has only one nonisolated point.

3. Quotient-universal sequential spaces. Let κ be an infinite cardinal and let $S(\kappa)$ denote the collection of all sequential spaces of cardinality $\leq \kappa$. A sequential space S is *quotient-universal** for $S(\kappa)$ if $S \in S(\kappa)$ and every $T \in S(\kappa)$ is a quotient of S . We are particularly interested in the existence of metrizable quotient-universal spaces for $S(\kappa)$.

Whenever $\kappa^{\mathfrak{c}} = \kappa$, the disjoint union of κ copies of the converging sequence will serve as a metrizable quotient-universal space for $S(\kappa)$. In particular, there is a metrizable quotient-universal space for $S(\mathfrak{c})$. In this section, we use the construction of §1 to demonstrate that, whether or not the continuum hypothesis is true, \mathfrak{c} is the smallest cardinal for which this is true. In fact, we exhibit a countable sequential space which is not a quotient of any metric space of cardinality $< \mathfrak{c}$.

LEMMA. *There exists a subset E of \mathbf{R} with $|E| = \mathfrak{c}$ which contains no uncountable closed subset of \mathbf{R} .*

Proof. Let $\{C_\alpha \mid \alpha < \mathfrak{c}\}$ be a transfinite enumeration of the \mathfrak{c} uncountable closed subsets of \mathbf{R} . Pick p_0 and q_0 in C_0 with $p_0 \neq q_0$. If p_α and q_α have been chosen in C_α for $\alpha < \beta$ so that all p_α and q_α are distinct, choose p_β and q_β in C_β so that $p_\beta \neq q_\beta$ and p_β, q_β are distinct from all p_α, q_α for $\alpha < \beta$. This is possible since any uncountable closed subset of \mathbf{R} has cardinal \mathfrak{c} so that $C_\beta - \{p_\alpha, q_\alpha \mid \alpha < \beta\} \neq \emptyset$.

* The term "universal" has been preempted by those who study spaces with a given property P which contain as subspaces every space (of appropriate cardinality or weight) having property P . See, for example, [2], [5] and [6].

Let $E = \{p_\alpha \mid \alpha < \mathfrak{c}\}$. Then $|E| = \mathfrak{c}$ and E contains no uncountable closed subset of \mathbf{R} since $q_\alpha \in C_\alpha - E$ for each α .

Let $E \subseteq \mathbf{R}$ be the set of the lemma. Let M_E denote the subspace $[Q \times (Q - \{0\})] \cup (E \times \{0\})$ of $\mathbf{R} \times \mathbf{R}$. Recall that Y_E is the quotient of M_E obtained by collapsing $E \times \{0\}$ to a single point e . Let $q: M_E \rightarrow Y_E$ be the quotient map.

Y_E is a countable sequential space, but:

THEOREM. Y_E is not the quotient of any metric space of cardinality $< \mathfrak{c}$.

Proof. Suppose there is a quotient map f of S onto Y_E , where S is a metric space and $|S| = \kappa < \mathfrak{c}$. For each $p \in E$, let $\sigma_p = (x_{p1}, x_{p2}, \dots)$ be a sequence in $Q \times (Q - \{0\})$ such that

$$|x_{pn} - (p, 0)| \leq \min \left\{ \frac{1}{n}, |x_{p(n-1)} - (p, 0)| \right\}.$$

Recall that q denotes the quotient map of M_E onto Y_E . For each n , let

$$z_{pn} = q(x_{pn})$$

and denote by η_p the sequence (z_{p1}, z_{p2}, \dots) in Y_E . Now $\eta_p \rightarrow e$. Hence, since f is a hereditary quotient map, there exists some $b_p \in f^{-1}(e)$ and a sequence $\sigma_p = (s_{p1}, s_{p2}, \dots)$ in $S - f^{-1}(e)$ such that $\sigma_p \rightarrow b_p$ and $f(\sigma_p) = \eta_p$. Let

$$f^{-1}(e) = \{x_\alpha \mid \alpha < \kappa\}$$

and, for $\alpha < \kappa$, let

$$A_\alpha = \{p \in E \mid b_p = x_\alpha\}.$$

We claim some A_α must contain a sequence (p_i) converging to some element of $\mathbf{R} - E$. For otherwise $C1_{\mathbf{R}}(A_\alpha) \subset E$ for each $\alpha < \kappa$, whence E is the union of fewer than \mathfrak{c} closed sets. But since $|E| = \mathfrak{c}$, one of these would be an uncountable closed set in E , contradicting the construction of E .

Without loss of generality, say A_1 contains a sequence (p_i) which is closed and discrete in E . Then the sequence $\eta_{p_i} = (z_{p_i1}, z_{p_i2}, \dots)$ converges to e , for each i , and the sequence $\delta_{p_i} = (s_{p_i1}, s_{p_i2}, \dots)$ converges to x_1 , for each i . A diagonal sequence $(s_{p_1n_1}, s_{p_2n_2}, \dots)$ with $n_k \geq k$ for each k will then converge to x_1 . Then $(z_{p_1n_1}, z_{p_2n_2}, \dots)$ converges to e . Hence $(x_{p_1n_1}, x_{p_2n_2}, \dots)$ must have a cluster point in M_E .

But $|x_{p_k n_k} - (p_k, 0)| \leq |x_{p_k k} - (p_k, 0)| \leq 1/k$, so any cluster point of $(x_{p_1 n_1}, x_{p_2 n_2}, \dots)$ in M_E would be a cluster point of $((p_1, 0), (p_2, 0), \dots)$, which is impossible by choice of the p_i .

We conclude with some observations on extension of the result above.

(1) As noted in §2, there are 2^c mutually nonhomeomorphic spaces Y_E . Since there are at most c quotients of any single countable sequential space, there can exist no quotient-universal space (metrizable or not) for $S(\aleph_0)$. It is at least consistent with the usual (Zermelo-Fraenkel) axioms for set theory (with Choice) that this result extends to all cardinals $\kappa < c$, for Martin's axiom entails $2^\kappa < 2^c$ for $\kappa < c$.

(2) Let $M(\kappa)$ denote the collection of metrizable spaces of cardinal $\leq \kappa$. The space Q of rationals is a (metrizable) quotient-universal space for $M(\aleph_0)$, while the disjoint union of c copies of the converging sequence is a quotient-universal space for $M(c)$. For cardinals κ between \aleph_0 and c little is known. Baumgartner ([1]) has shown that it is consistent with Zermelo-Fraenkel set theory with choice that all \aleph_1 -dense subsets of \mathbf{R} are order-isomorphic. (A subset A of \mathbf{R} is \aleph_1 -dense if whenever $a < b$ in \mathbf{R} , $(a, b) \cap A$ has cardinal \aleph_1 .) If this is the case, then every separable metric space M of cardinal $\leq \aleph_1$ is a quotient of the unique \aleph_1 -dense subset D of \mathbf{R} . For M is a quotient of $M \times D$, while ([7], Theorem 76) $M \times D$ is homeomorphic to a subset of \mathbf{R} and hence, by Baumgartner's result, to D .

REFERENCES

1. James E. Baumgartner, *All \aleph_1 -dense sets of reals can be isomorphic*, Fund. Math., **79** (1973), 103–106.
2. Jonsson, *Universal Relational Systems*, Math. Scand., **4** (1956), 193–208.
3. E. Michael, *\aleph_0 -spaces*, J. Math. Mech., **15** (1966), 983–1002.
4. E. Michael and A. H. Stone, *Quotients of the space of irrationals*, Pacific J. Math., **28** (1969), 629–633.
5. R. Pol, *There is no universal totally disconnected space*, Fund. Math., **79** (1973), 265–267.
6. W. Sierpinski, *Sur les espaces métriques universels*, Fund. Math., **33** (1945), 123–136.
7. ———, *General Topology*, 2nd ed., U. of Toronto Press, 1956 (translated by C. C. Krieger).

Received March 12, 1976 and in revised form May 12, 1976.

UNIVERSITY OF ALBERTA

Pacific Journal of Mathematics

Vol. 66, No. 1

November, 1976

Helen Elizabeth. Adams, <i>Factorization-prime ideals in integral domains</i>	1
Patrick Robert Ahern and Robert Bruce Schneider, <i>The boundary behavior of Henkin's kernel</i>	9
Daniel D. Anderson, Jacob R. Matijevic and Warren Douglas Nichols, <i>The Krull intersection theorem. II</i>	15
Efraim Pacillas Armendariz, <i>On semiprime P.I.-algebras over commutative regular rings</i>	23
Robert H. Bird and Charles John Parry, <i>Integral bases for bicyclic biquadratic fields over quadratic subfields</i>	29
Tae Ho Choe and Young Hee Hong, <i>Extensions of completely regular ordered spaces</i>	37
John Dauns, <i>Generalized monoform and quasi injective modules</i>	49
F. S. De Blasi, <i>On the differentiability of multifunctions</i>	67
Paul M. Eakin, Jr. and Avinash Madhav Sathaye, <i>R-endomorphisms of $R[[X]]$ are essentially continuous</i>	83
Larry Quin Eifler, <i>Open mapping theorems for probability measures on metric spaces</i>	89
Garret J. Etgen and James Pawlowski, <i>Oscillation criteria for second order self adjoint differential systems</i>	99
Ronald Fintushel, <i>Local S^1 actions on 3-manifolds</i>	111
Kenneth R. Goodearl, <i>Choquet simplexes and σ-convex faces</i>	119
John R. Graef, <i>Some nonoscillation criteria for higher order nonlinear differential equations</i>	125
Charles Henry Heiberg, <i>Norms of powers of absolutely convergent Fourier series: an example</i>	131
Les Andrew Karlovitz, <i>Existence of fixed points of nonexpansive mappings in a space without normal structure</i>	153
Gangaram S. Ladde, <i>Systems of functional differential inequalities and functional differential systems</i>	161
Joseph Michael Lambert, <i>Conditions for simultaneous approximation and interpolation with norm preservation in $C[a, b]$</i>	173
Ernest Paul Lane, <i>Insertion of a continuous function</i>	181
Robert F. Lax, <i>Weierstrass points of products of Riemann surfaces</i>	191
Dan McCord, <i>An estimate of the Nielsen number and an example concerning the Lefschetz fixed point theorem</i>	195
Paul Milnes and John Sydney Pym, <i>Counterexample in the theory of continuous functions on topological groups</i>	205
Peter Johanna I. M. De Paepe, <i>Homomorphism spaces of algebras of holomorphic functions</i>	211
Judith Ann Palagallo, <i>A representation of additive functionals on L^p-spaces, $0 < p < 1$</i>	221
S. M. Patel, <i>On generalized numerical ranges</i>	235
Thomas Thornton Read, <i>A limit-point criterion for expressions with oscillatory coefficients</i>	243
Elemer E. Rosinger, <i>Division of distributions</i>	257
Peter S. Shoenfeld, <i>Highly proximal and generalized almost finite extensions of minimal sets</i>	265
R. Sirois-Dumais and Stephen Willard, <i>Quotient-universal sequential spaces</i>	281
Robert Charles Thompson, <i>Convex and concave functions of singular values of matrix sums</i>	285
Edward D. Tymchatyn, <i>Some n-arc theorems</i>	291
Jang-Mei Gloria Wu, <i>Variation of Green's potential</i>	295